

THE GEOLOGICAL EVOLUTION OF AFRICA

Paul H.G.M. Dirks

School of Geosciences, University of the Witwatersrand, Johannesburg, South Africa

Tom G. Blenkinsop

School of Earth Sciences, James Cook University, Townsville, Australia

Hielke A. Jelsma

De Beers Consolidated Mines Ltd., Johannesburg, South Africa

Keywords: African geology, Pan African orogen, Archaean cratons, African Rift, hot spots, plume activity

Contents

1. Introduction
2. The Archean between 3800-2550 MA: Formation of Cratons
 - 2.2. The Central Africa (Congo-Tanzania) Craton
 - 2.3. The Southern Africa Craton
 - 2.4. An Archean Passive Margin Sequence and Foreland Basin
 - 2.5. East Sahara or Nile Craton
 - 2.6. Malagasy Shield
3. Paleoproterozoic Growth of Archean Cratonic Blocks: the Eburnian
 - 3.1. Rifts and Passive Margins
 - 3.1.1. The Southern Africa Craton
 - 3.2. Active Continental Margins and Accretion of Volcanic Arcs
 - 3.2.1. The Southern Africa Craton
 - 3.2.2. The West Africa Craton: the Birrimian
 - 3.2.3. The SW Congo Craton
 - 3.3. Continental Collision
 - 3.3.1. Southern Africa Craton: the Magondi-Okwa-Kheis Belts
 - 3.3.2. The NW Congo Craton: the Gabon Belt
 - 3.3.3. Collision of the Congo and Tanzania Cratons: the Ubendian-Ruzizian-Ruwenzori Belts
 - 3.3.4. Collision along the SE Tanzania Craton: the Usagaran Belt
4. Mesoproterozoic Continental Break-up and Growth
 - 4.1. Rifts and Passive Margins
 - 4.1.1. The Central Africa Craton: the Kibaran Belt, a Failed Rift
 - 4.2. Active Continental Margins and Accretion of Volcanic Arcs
 - 4.2.1. The S Southern Africa Craton: the Sinclair-Namaqua-Natal Belts
 - 4.3. Continental Collision
 - 4.3.1. Collision of the Central and Southern Africa Cratons: the Irumide-Zambezi-Lurio Belts
 - 4.4. Kibaran Mantle Plume Activity
5. Neoproterozoic Continental Break-up and Growth
 - 5.1. Rifts and Passive Margins

- 5.1.1. Rift-drift of the Proto-South Atlantic (Adamaster) Ocean: West Congo-Kaoko-Gariiep-Saldania Belts
- 5.1.2. Rifts and Proto-oceanic Basins in Southern Africa: Damara-Lufilian-Zambezi Belts
- 5.1.3. Passive Margins along the West Africa Craton
- 5.2. Active Continental Margins, Accretion of Volcanic Arcs and Continental Collision
 - 5.2.1. The East African Orogen
 - 5.2.2. West Congo-Kaoko-Gariiep-Saldania Belts
 - 5.2.3. The Trans-Saharan Belt
 - 5.2.4. The Mauritanide-Bessaride-Rokelide Belts
 - 5.2.5. The Damara-Lufilian-Zambezi Belts
 - 5.2.6. The Oubangide Belt
- 6. The Phanerozoic Evolution of Africa
 - 6.1. Paleozoic Continental Margins of Gondwana (570-290 Ma)
 - 6.1.1. The Cape Supergroup, South Africa
 - 6.1.2. The North African Shelf
 - 6.2. Hercynian Collision: the Gondwanide Orogens
 - 6.2.1. Cape Fold Belt
 - 6.2.2. Mauritanide Belt
 - 6.3. Rifting of Gondwana: The Karoo basins (290-180 Ma)
 - 6.3.1. The Karoo Rift Basins
 - 6.3.2. The Karoo Foreland Basins and Sags
 - 6.4. Development of a Stable Tethys Shelf in North Africa (280-110 Ma)
 - 6.5. Break up of Gondwana
 - 6.5.1. E African Margin Development (165 Ma-Recent)
 - 6.5.2. North Atlantic Ocean, NW African Margin Development (180 Ma-Recent)
 - 6.5.3. South Atlantic Ocean, W African Margin Development (150 Ma-Recent)
 - 6.6. Rift-sag Basins of North, West and Central Africa (140 Ma-Recent)
 - 6.7. East African Rift System 30 Ma-Recent and Development of Red Sea Margin
 - 6.8. Collision between Europe and Africa (90 Ma-Recent)
 - 6.8.1. Atlas Mountains
 - 6.9. Hot Spots and Sags: the Story of a Stagnant Continent: 30 Ma to Recent
- Glossary
- Bibliography
- Biographical Sketches

Summary

Seven Archaean cratonic nuclei characterized by granite-greenstone terrains form the foundation of Africa. During the Neoproterozoic and Paleoproterozoic these Archaean nuclei merged into three major cratons referred to as the West Africa, Central Africa (Congo-Tanzania) and Southern Africa (Zimbabwe-Kaapvaal) cratons.

The Paleoproterozoic geology of the African Plate is dominated by sedimentation, volcanism, and tectonism between 2.3 and 1.8 Ga. These events are collectively known as the Eburnian orogeny and have been recognized in the Southern, Central and West Africa cratons.

The worldwide 1.35-0.9 Ga cascade of continental collisions that constructed the supercontinent Rodinia is known in Africa as the Kibaran orogeny. Effects of this can be found along the margins of the Central and Southern Africa Cratons. Kibaran events did not affect the West African Craton.

The break-up of Rodinia at 0.85 Ga led to a fan-shaped aggregation of pre-Gondwana continental blocks including East Gondwana, West Gondwana and the intervening African cratonic blocks. These blocks were separated by oceanic basins, which were consumed and closed between 0.85-0.55 Ga during events referred to as the Pan-African Orogeny.

By 0.55 Ga the Gondwana Supercontinent had formed with Africa at its center. For the next 350 Ma this continent remained in existence and Africa only experienced sedimentation along the Gondwana margin or in intra-continental rift-sag basins. Orogenic activity during this period was limited to the NW and S extremities of the continent. From the late Jurassic onwards, Gondwana rifted resulting in the African plate as we now know it, with most tectonic activity controlled by extension and hot spot activity.

1. Introduction

The African continent preserves evidence for major crust-forming events dating back to 3.8 Ga. These events represent cycles of continental break-up and growth, which have been recognized worldwide and can be largely explained in a plate-tectonic context, within the confines of partly overlapping Wilson cycles. In Africa the main orogenic episodes are shown in Table 1.

<i>Orogeny</i>	<i>Age</i>	<i>Main outcome</i>
Paleoarchean	3.55-3.15 Ga	Formation of early Archean cratonic cores (Kaapvaal, Tokwe)
Mesoarchean	3.15-2.75 Ga	Accretionary growth of Kaapvaal, Zimbabwe, Congo, Tanzania cratons and the Man and Reguibat shields
Neoarchean	2.75-2.55 Ga	Stabilisation of Kaapvaal, Zimbabwe, Congo, Tanzania cratons and the Man and Reguibat shields; merging of Kaapvaal and Zimbabwe cratons as Southern Africa Craton
Eburnian	2.2-1.8 Ga	Growth of the West Africa Craton along an active accretionary margin (Birimian). Merging of the Congo and Tanzania cratons in the Central Africa Craton. Passive margin development and orogenesis along the W margin of Central and Southern Africa cratons.
Kibaran	1.4-0.85 Ga	Merging of Southern and Central Africa cratons as part of the Rodinia supercontinent
Pan-African	0.85-0.5 Ga	Merging of all cratonic fragments to form the Gondwana supercontinent to which Africa is central
Hercynian	0.45-	Limited collision and tectonic activity along the NW

	0.25 Ga	and S margins of the African plate
Alpine	0.12-0 Ga	Subduction of African plate under Eurasia and formation of the Atlas mountains

Table 1: Summary of the main orogenic episodes in Africa
All figures with this text have been largely adapted from Choubert & Faure-Muret (1990).

2. The Archean between 3800-2550 MA: Formation of Cratons

Seven major cratonic nuclei form the foundation of Africa: the Kaapvaal craton, the Zimbabwe Craton, the Tanzania Craton, the Congo Craton, the Man Shield, the Reguibat Shield and the elusive (largely covered and reworked) East Sahara or Nile Craton. During the Neoproterozoic and Paleoproterozoic these Archean nuclei merged into three major cratons which will be referred to as the West Africa, Central Africa (Congo-Tanzania) and Southern Africa (Zimbabwe-Kaapvaal) cratons (de Wit and Ashwal, 1997; Trompette, 1994; Thomas et al., 1993). Archean terranes in Africa are shown in Fig. 1.




Figure 1: Archean and Paleoproterozoic (Ubendian) terranes (3800-1750 Ma)

2.1. The West Africa Craton

The West Africa Craton (1.5 million km² Archean; 3.0 million km² Paleoproterozoic) in NW Africa consists of the Reguibat and (Leo-) Man Shields in the N and S of the craton, respectively, that are separated by the Neoproterozoic to Paleozoic Taoudeni basin. In both shields, Archean rocks are exposed in the western parts and separated from Paleoproterozoic rocks to the east by major shear zones, referred to as the Sassandra Fault in the Man Shield (Ivory Coast) and the Zednes Fault in the Reguibat Shield (Mauritania) (Attoh & Ekwueme, 1997).

Age dating and paleomagnetic reconstructions suggest that the West Africa Craton and the Guyana Shield (Amazonian Craton) in South America formed a single domain in the Palaeoproterozoic. A possible Archean suture may be found along the SW margin of the craton (Sierra Leone, Liberia) where a linear belt of Archean granulite-facies supracrustal and meta-igneous rocks (Kasila Group) occurs, marked by a shallowly W-dipping mylonite zone. This area has been overprinted by deformation related to the Pan-African Rokelide Belt (Williams & Culver, 1988).

The Archean rocks of the **Reguibat Shield** include TTG-type migmatitic orthogneiss and metavolcanic and metasedimentary belts with ironstones, marbles, ultramafic rock, and amphibolite (e.g. Tasiast belt, Amsaga area in Mauritania), greywacke and pelite. The gneisses have ages of 3.52 Ga, 3.42 Ga and 2.84 Ga and are intruded by 2.99-2.83 Ga charnockitic gneiss. The shield has been affected at 2.74 Ga by polyphase folding and high-grade metamorphism followed by the emplacement of late-tectonic granites at 2.73-2.72 Ga. Retrograde amphibolite-grade mineral assemblages occur near the main shear zones (Attoh & Ekwueme, 1997; Potrel et al., 1998).

The oldest component of the **Man Shield** consists of banded TTG gneiss (> 3.0 Ga) overlain by greenstone belts with (ultra)mafic metavolcanics, banded ironstone, phyllite, greywacke and quartzite (e.g. Sula and Nimini belts in Sierra Leone) and intruded by 2.97-2.78 Ga granites (Attoh & Ekwueme, 1997).

2.2. The Central Africa (Congo-Tanzania) Craton

The Congo-Tanzania Craton is a region of pre-2.5 Ga continental crust, located in central Africa and includes the Tanzania and Congo cratons, linked by the Archean Uganda Basement Gneiss and West Nile Complex (> 2.8 Ga Kilo-Moto Terrane).

The **Tanzania Craton** (0.5 million km²; Tanzania, SW Kenya, SE Uganda) consists of the >2.9 Ga Dodoma Terrane in the south and the younger Lake Victoria Terrane in the north. The Dodoma Terrane comprises high-grade TTG-type orthogneiss, mafic-ultramafic rocks, quartzite and phyllite.

The Lake Victoria Terrane consists of tectonized greenstone belts intruded by large amounts of granite. The greenstones consist of 2.81-2.70 Ga bimodal volcanics and flysch (Nyanzian Group) overlain by 2.68-2.63 Ga volcanics and clastic sediments (Nzega and Kakamega Groups). Basalt, andesite, shoshonitic rhyolite and conglomerate (Rongo Group) unconformably overlie the older sequences. Granite was emplaced

during several stages at 2.72-2.68 Ga (early-syntectonic), 2.64 Ga (late-syntectonic) and 2.58-2.56 Ga (the post tectonic Western Granite Complex). A 2.53 Ga platform sequence was deposited on the NE and N parts of the craton (Kisii, Buganda Groups) (Borg & Krogh, 1999; Pinna et al., 1996, 2000).

Archean outcrops in the **Congo Craton** include the Chaillu - Gabon blocks (SW Cameroon, Equatorial Guinea, Gabon, Congo), the Zaire Block (NE DR Congo, W Uganda), the Angola Block and the Kasai Block (NE Angola, S DR Congo).

The **Chaillu - Gabon blocks** contain 3.19-3.12 Ga granite-greenstones deformed and metamorphosed at high metamorphic grades at ~ 3.15 Ga. They are intruded by 2.95-2.85 Ga calc-alkaline tonalites and granites, associated with intermediate to felsic volcanics (2.97-2.94 Ga), metamorphosed at low metamorphic grades. Ultrabasic rocks intruded at 2.78 Ga, followed by late-orogenic granites between 2.8-2.5 Ga before stabilization of the block (Feybesse et al., 1998).

The **Zaire Block** can be divided into the Bomu (amphibolite-gneiss) Complex and Gangan greenstone belts in the far W, the West Nile Complex in the N, and the Upper-Zaire Granitoid Massif with Kibalian greenstone belts in the S. The Kibalian Group greenstones are thought to rest unconformably on older gneiss and consist of a mafic volcanic-dominated lower portion intruded by 2.8 Ga tonalite, and a bimodal volcano-sedimentary upper sequence intruded by 2.45 Ga old granite, similar to greenstones found in the Tanzania Craton. The West Nile Complex consists of high-grade gneiss with remnant mafic greenstone segments. The gneiss complex can be traced across S Sudan and central Uganda into the Tanzania craton and may form a link (orogenic belt ?) between both cratons. The > 3.0 Ga Gangan Group greenstones are thought to rest unconformably on still older gneisses of the Bomu Complex. The Gangan greenstones form distinct belts of slightly metamorphosed quartzite, slate, jaspilite, talc schist and dolerite. The 3.4-3.0 Ga Bomu Complex extends into the Central African Republic and is composed of migmatitic gneiss with mafic and metasedimentary schist inclusions. Deformation has been recorded at ~ 3.3-3.0 Ga and 3.0 Ga (Borg & Shackleton, 1997; Lavreau, 1982, 1984).

The Archean of the **Angola Block** was largely reworked during the Eburnian event. Archean gneiss was metamorphosed at ~2.8 Ga and intruded by granite between 2.83-2.60 Ga. They are covered by volcano-sedimentary rocks (Jamba Group in the N; Utende-Chela Supergroup in the S) of poorly constrained age (2.8-2.2 Ga). In the SW, Archean granite-greenstone terrains occur (de Carvalho, 1983). De Wit (2001, unpublished data) reports 3.5 and 2.7 Ga ages and a 2.0 Ga overprint, indicating that the Angola Block is more complicated than previously known.

The **Kasai Block** consists of tonalitic gneiss and old granodiorite (3.5-3.3 Ga) associated with supracrustals, gabbro-norite-anorthosite complexes and amphibolite, metamorphosed at granulite facies around 2.8 Ga during emplacement of 2.9-2.8 Ga charnockite and 2.83 Ga granite. A second event is associated with widespread calc-alkaline granite emplacement and migmatization between 2.7-2.6 Ga.

2.3. The Southern Africa Craton

The Southern Africa Craton comprises the Zimbabwe Craton to the N and the Kaapvaal Craton to the S, merged across the high-grade gneisses of the Limpopo Belt.

The 3.55-2.58 Ga **Zimbabwe Craton** (0.3 million km², Zimbabwe, NE Botswana, W Mozambique) contains a 3.55-3.35 Ga, central gneissic nucleus (Tokwe Segment) affected by regional deformation (N-S grain) > 3.35 Ga. After it stabilized, ~3.0 Ga stable shelf sedimentation occurred along its W margin. Subsequent tectono-magmatic events added granite and greenstones including clastic and deep-water sediments at ~2.9-2.8 Ga (Belingwean, Lower Bulawayan Groups) and ~2.72-2.64 Ga (Upper Bulawayan, Shamvaian Groups). Greenstone forming events are either explained in continental rift- or flood basalt settings with deposition on older basement, or in active continental margin settings involving subduction-accretion and back-arc rifting. Important Archean unconformities have been described at the base of greenstone sequences in the Belingwe greenstone belt (Wilson et al., 1995; Jelsma & Dirks, 2002).

Thermal and isostatic stabilization of the craton occurred at 2.6 Ga with the emplacement of late-orogenic monzogranite (Chilimanzi Suite) followed by crustal relaxation, and emplacement of the Great Dyke layered mafic-ultramafic complex at 2.58 Ga. A 2.62-2.56 Ga granulite terrane occurs along the N margin of the Zimbabwe Craton. The terrane represents Archean lower crust brought to the surface during Kibaran and Pan-African events in the adjacent Zambezi belt (Jelsma & Dirks, 2002).

The **Kaapvaal Craton** (1.2 million km², South Africa, Botswana, Lesotho, Swaziland) originated from an early nucleus that contains the Barberton greenstone belt, Natal Terrane and Ancient Gneiss Complex with the oldest known crustal fragment in Africa (Ngwane Gneiss, 3.64 Ga).

In the Barberton greenstone belt, oceanic, 3.49-3.46 Ga (ultra)mafic volcanic and plutonic rocks experienced 3.45-3.42 Ga deformation-metamorphism and TTG-type plutonism as oceanic crust and 3.55-3.52 Ga tonalitic gneiss were obducted onto an active arc-trench-like terrain. An early continental margin may have formed to the S (3.4 Ga, Mkhondo-Mahamba Groups, Swaziland). At 3.3-3.2 Ga, renewed tectono-magmatism resulted in crustal thickening with clastic sedimentation (Fig Tree-Moodies Groups), evolving to late strike-slip shear and post-tectonic granite emplacement between 3.15-3.07 Ga (Brandl & de Wit, 1997).

By 3.1 Ga, a large part of the Kaapvaal Craton had stabilized and a thick volcano-sedimentary pile accumulated on this proto-cratonic block. Along its N margin, the ~3.2-2.97 Ga Pietersburg-Giyani and Murchison Terranes represent juvenile oceanic- and island-arc crust that accreted with the proto-craton between 3.2-2.8 Ga. West of the proto-craton, the ~3.1-2.9 Ga Amalia, Colesburg and Kraaipan Terranes accreted along the Colesberg Lineament (Brandl & de Wit, 1997; McCourt, S., 1995).

The Zimbabwe and Kaapvaal cratons were juxtaposed at ~2.6 Ga across the ENE trending **Limpopo Belt** that is separated from the cratons by major thrusts (North Limpopo Thrust Zone in the N; Hout River Shear Zone in the S). The Limpopo Belt has been interpreted as an Archean collisional orogen and comprises the North Marginal

Zone (reworked Zimbabwe craton), Central Zone and South Marginal Zone (reworked Kaapvaal Craton). The marginal zones have been interpreted as metamorphosed deep-crustal equivalents of granite-greenstones in the adjacent cratons. The Central Zone is separated from the marginal zones by the dextral Triangle Shear Zone to the north and the sinistral Palala shear zone to the south. The Central Zone preserves no evidence of greenstone precursors, but is characterized by platform sediments (provenance 3.8-3.3 Ga) deposited on a basement of 3.25-3.17 Ga orthogneiss (Sand River gneiss). Deformation occurred at high-grades at ~3.2-3.1 Ga (Central Zone only), 2.65-2.55 Ga (collision) and 2.05-1.95 Ga (transpression).

2.4. An Archean Passive Margin Sequence and Foreland Basin

A large section of the Kaapvaal craton stabilized by 3.1 Ga, and acted as basement to the thick volcano-sedimentary sequences of the co-evolving (> 2.94 Ga) Pongola and (3.12-2.71 Ga) Witwatersrand basins. In the Witwatersrand basin, the sequence comprises the lowermost 3.12-3.07 Ga Dominion Group volcanics, overlain by the Witwatersrand Supergroup sediments. Sedimentation took place during two cycles, with rocks of the lower cycle forming on a passive continental margin.

Sediments of the upper cycle are rich in gold and accumulated between 2.84-2.71 Ga in a foreland basin. Early rifting evolved to post-rift subsidence and flexural loading due to thrusting along the SE and NW margins of the basin. The evolution of the Witwatersrand Basin came to an end with the vast outpouring of the Ventersdorp Group rift-basin volcanics at ~ 2.71 Ga (Brandl & de Wit, 1997; Coward et al., 1995).

2.5. East Sahara or Nile Craton

The poorly known East Sahara or Nile Craton stretches E-W from the Hoggar Massif (Algeria) to the Western Desert (Egypt). It is largely covered by younger rocks and reworked by Pan-African events. Rocks from the Uweinat Massif (Egypt) are as old as 2.63 Ga, but the bulk of the pre-Pan-African Nd model ages reported for the craton are Proterozoic. Provenance ages and xenocryst ages from Egypt and Sudan indicate significant episodes of crustal growth during the Proterozoic at 2.5-2.4 Ga, 2.1-1.9 Ga, 1.7-1.2 Ga and 1.0-0.8 Ga (Stern et al., 1994).

2.6. Malagasy Shield

The Malagasy Shield (Madagascar) is divided in two parts by the NW-SE trending Bongolava-Ranotsara shear zone with high-grade Pan-African gneiss affecting Mesoproterozoic and possibly Archean protoliths to the S. To the N of the shear zone, low- and medium-grade greenstone-gneiss terrains occur separated by metasedimentary rocks and granitoid.

The greenstone belt stratigraphy has been divided into metasedimentary/paragneiss dominated (Androyen, Graphite Systems), and greenstone dominated (Vohibory System) sequences. A migmatite tonalite gneiss in the sequence is dated at 3.19 Ga, whilst 2.52-2.49 Ga, granitic orthogneiss intruded the greenstone (Tucker et al., 1997).

-
-
-

TO ACCESS ALL THE 36 PAGES OF THIS CHAPTER,
Visit: <http://www.eolss.net/Eolss-sampleAllChapter.aspx>

Bibliography

Abouachami, W., Boher, M., Michard, A. & Albarede, F., (1990). A major 2.1 Ga event of mafic magmatism in West Africa: An early stage of crustal accretion. *Journal of Geophysical Research* **95**, 17605-17629.

Attoh, K. & Ekwueme, B.N., (1997). The West African Shield. In: de Wit, M. and Ashwal, L.D. (eds.) *Greenstone belts*. Oxford University Press. 517-528.

Attoh, K., Dallmeyer, R.D. & Affaton, P., (1997). Chronology of nappe assembly in the Pan-African Dahomeyide orogen, West Africa: evidence from ⁴⁰Ar/³⁹Ar mineral ages. *Precambrian Research* **82**, 153-171.

Boher, M., Abouchami, W., Michard, A., Albarede, F. & Arndt, N., (1992). Crustal growth in West Africa at 2.1 Ga. *Journal of Geophysical Research*, **97**, B1, 345-369.

Boote, D.R.D., Clark-Lowes, D.D. & Traut, M.W., (1998). Palaeozoic petroleum systems of North Africa. *Geological Society of London Special Publications* **132**, 7-68. [Contains a good overview of petroleum basins in N Africa].

Borg, G. & Shackleton, R.M., (1997). The Tanzania and NE-Zaire Cratons. In: de Wit, M. and Ashwal, L.D. (eds.) *Greenstone belts*. Oxford University Press, 608-619.

Borg, G. & Krogh, T., (1999). Isotopic age data of single zircons from the Archaean Sukamaland greenstone belt, Tanzania. *Journal of African Earth Sciences*, **29**, 301-312.

Bossière, G., Bonkougou, I., Peucat, J.-J. & Pupin, J.-P., (1996). Origin and age of Paleoproterozoic conglomerates and sandstones of the Tarkwaian Group in Burkina Faso, West Africa. *Precambrian Research*, **80**, 153-172.

Brandl, G. & de Wit, M.J., (1997). The Kaapvaal Craton. In: de Wit, M. and Ashwal, L.D. (eds.) *Greenstone belts*. Oxford University Press, 581-607.

Brede, R., Hauptman, M. & Herbig, H.B., (1992). Plate tectonics and the intracratonic mountain ranges in Morocco. *Geologische Rundschau*, **81**, 127-141.

Brown, R.W. Gallagher, K., Gleadow A.J. & Summerfield, M.A., (2000). Morphotectonic evolution of the South Atlantic margins of Africa and South America. In: M.A. Summerfield (Ed.) *Geomorphology and global tectonics*, 255-281.

Burke, K.(1996). The African Plate. *S. Afr. J. Geol.*, **99**, 341-409. [A comprehensive overview of tectonic events affecting the African plate over the past 50 Ma].

Burke, K.C. & Dewey, J.F., (1972). Orogeny in Africa. In: Dessauvage, T.F., Whiteman, A.J., *African Geology*. University of Ibadan Press, 583-608. [A classic].

Cahen, L. Snelling, N.J., Delhal, J. & Vail, J.R., (1984). *The geochronology and evolution of Africa*. Clarendon Press, 512p [Presents a very good data base with regards older references to African geology].

Carvalho, H., Fernandez, A. & Vialette, Y. (1987). The Kibaran cycle in Angola, a discussion. *Geological Journal*, **22**, 85-102.

Castaing, C., Triboulet, C., Feybesse, J.L. & Chevremont, P., (1993). Tectonometamorphic evolution of Ghana, Togo and Benin in the light of the Pan-African/Brasiliano orogeny. *Tectonophysics*, **218**, 323-342

Choubert, G. & Faure-Muret, A., (1990). *Carte Geologique Internationale de l'Afrique*. Commission for the geological map of the world, CCGM & UNESCO, Paris. [This map represents the most comprehensive tectonic compilation of the continent].

Costa, M., Cadoppi, P., Sacchi, R. & Fanning, C.M., (1994). U-Pb SHRIMP dating of zircons from Mozambique gneiss. *Boll. Soc. Geol. Italiana*, **113**, 173-178.

Coward, M.P., Purdy, E.G., Ries, A.C. & Smith, D.G., (1999). The distribution of petroleum reserves in basins of the South Atlantic margins. In: Cameron, N.R., Bate, R.H. & Clure, V.S. (eds). The oil and gas habitats of the South Atlantic. *Geol. Soc. Lond. Spec. Publ.*, **153**, 101-131.

Coward, M.P., Spencer, R.M. & Spencer, C.E., (1995). Development of the Witwatersrand Basin, South Africa. In: Coward, M.P., Ries, A.C. (eds). Early Precambrian Processes. *Geol. Soc. Spec. Publ.*, **95**, 243-269.

Cox, K.G., (1992). Karoo igneous activity and the early stages of break-up of Gondwanaland. In: Storey, B.C., Alabaster, T. & Pankhurst, R.J. (eds), Magmatism and the causes of continental break-up. *Geol. Soc. Spec. Publ.* **68**, 137-148.

Culver, S.J., (1988). Plate boundaries in west Africa and their implications for Pangean continental fit. *Geology*, **16**, 322-325.

Dallmeyer, R.D. & Villeneuve, M., (1987). ⁴⁰Ar/³⁹Ar mineral age record of polyphase tectono-thermal evolution in the southern Mauritanide orogen, southeastern Senegal. *GSA Bulletin* **98**, 602-611.

Daly, M.C., (1986). The intracratonic Irumide Belt of Zambia and its bearing on collisional orogeny during the Proterozoic of Africa. In: Coward, M.P., Ries, A.C. (eds.) Collision Tectonics. *Geol. Soc. Spec. Publ.*, **19**, 321-328.

Daly, M.C., (1988). Crustal shear zones in central Africa: a kinematic approach to Proterozoic tectonics. *Episodes* **11**, 5-11.

Dalziel, I.W.D., Mosher, S. & Gahagan, L.M., (2000). Laurentia-Kalahari collision and the assembly of Rodinia. *Journal of Geology*, **108**, 499-513.

De Carvalho, H., (1983). Notice explicative préliminaire sur la géologie de l'Angola. *Sep. Garcia de Orta, Ser. Geol., Lisboa*, **6**, 15-30.

de Wit, M. & Ashwal, L.D. (eds.) (1997). *Greenstone belts*. Oxford University Press. [Excellent overview of greenstone terrains in Africa].

Dirks, P.H. G. M. & Sithole, T. (1999). Eclogites in the Makuti gneisses of Zimbabwe: implications for the tectonic evolution of the Zambezi Belt in southern Africa. *Journal of Metamorphic Geology*, **17**, 593-612.

Dirks, P.H.G.M., Jelsma, H.A., Vinyu, M. & Munyanyiwa, H. (1998). The structural history of the Zambezi belt in northeast Zimbabwe; evidence for crustal extension during the early Pan-African. *S. Af. J. Geol.* **10**, 1-16.

- Doglioni, C., Gueguen, E., Harabaglia, P. & Mongelli, F., (1999). On the origin of west-directed subduction zones and applications to the western mediterranean. In: Durand, B. Jolivet, L. Horvath, F. & Seranne, M. (eds). *The mediterranean basins: Tertiary extension within the Alpine Orogen*. *Geol. Soc. Lond. Spec. Publ.*, **156**, 541-561.
- Doumbia, S., Pouclet, A., Kouamelan, A., Peucat, J.J., Vidal, M. & Delor, C., (1998). Petrogenesis of juvenile-type Birimian (Paleoproterozoic) granitoids in central Côte-d'Ivoire, West Africa: geochemistry and geochronology. *Precambrian Research*, **87**, 33-63.
- Ferré, E., Délérès, J., Bouchez, J.-L., Lar, A.U. & Peucat, J.-J., (1996). The Pan-African reactivation of Eburnean and Archaean provinces in Nigeria: structural and isotopic data. *Journal of the Geological Society, London*, **153**, 719-728.
- Feybesse, J.-L., Johan, V., Triboulet, C., Guerrot, C., Mayaga-Mikolo, F., Bouchot, V. & Eko N'dong, J., (1998). The West Central African belt: a model of 2.5-2.0 Ga accretion and two-phase orogenic evolution. *Precambrian Research*, **87**, 161-216.
- Feybesse, J.-L. & Milesi, J.-P., (1994). The Archaean/Proterozoic contact zone in West Africa: a mountain belt of decollement thrusting and folding on a continental margin related to 2.1 Ga convergence of Archaean cratons? *Precambrian Research*, **69**, 199-227.
- Gabert, G. (1984). Structural-lithological units of Proterozoic rocks in east Africa, their base cover and mineralisation. In: Klerkx, J. & Michot, J. (eds) *African geology*. Musée Royal de L'Afrique centrale, Tervuren, 11-22.
- Goscombe, B., Armstrong, R. & Barton, J.M. (2000). Geology of the Chewore inliers, Zimbabwe: constraining the mesoproterozoic to Paleozoic evolution of the Zambezi belt. *Journal of African Earth Sciences* **30**, 589-627.
- Green, J.C., (1992). Proterozoic Rifts. In: *Developments in Precambrian Geology 10, Proterozoic Crustal Evolution*, Ed. K.C. Condie, Elsevier, Amsterdam, 97-149.
- Gurnis, M., Mitrovica, J.X., Ritsema, J. & van Heijst, H.-J., (2000). Constraining mantle density structure using geological evidence of surface uplift rates: the case of the African Superplume. *Geochemistry, Geophysics, Geosystems*, **1**, pages?
- Hanson, R., (2001). Proterozoic tectonic evolution and geochronology of Southern Africa. In: Windley (ed). *Proterozoic East Gondwana: Supercontinent assembly and break-up*. *Geological Society of London Special Publications*, Under review. [Excellent overview of the geochronological evolution of Southern Africa with numerous references].
- Hanson, R.E., Martin, M.W., Bowring, S.A. & Munyanyiwa, H., (1998). U-Pb zircon age for the Umkondo dolerites, eastern Zimbabwe: 1.1 Ga large igneous province in southern Africa-East Antarctica and possible Rodinia correlations. *Geology*, **26**, 1143-1146.
- Hanson, R.E., Wilson, T.J. & Munyanyiwa, H., (1994). Geologic evolution of the Neoproterozoic Zambezi Orogenic Belt in Zambia. *Journal of African Earth Sciences*, **18**, 135-150.
- Hartley, R., Watts, A.B. & Fairhead, J.D., (1996). Isostasy of Africa. *Earth and Planetary Science Letters*, **137**, 1-18.
- Hirdes, W., Davis, D.W., Lüdtke, G. & Konan, G., (1996). Two generations of Birimian (Paleoproterozoic) volcanic belts in northeastern Côte d'Ivoire (West Africa): consequences for the "Birimian controversy". *Precambrian Research*, **80**, 173-191.
- Hoffman, P.F., (1999). The break-up of Rodinia, birth of Gondwana, true polar wander and the snowball Earth. *Journal of African Earth Sciences*, **28**, 17-33.

Holzer, L., Frei, R., Barton, J. M. & Kramers, J. D. (1998). Unravelling the record of successive high grade events in the Central Zone of the Limpopo Belt using Pb single phase dating of metamorphic minerals. *Precambrian Research* **87**, 87-115.

Hunter, D.R. (ed.) (1981). *Precambrian of the Southern Hemisphere*. Elsevier, Amsterdam, 882p.

Jelsma, H.A. & Dirks, P.H.G.M., (2002). Neoproterozoic tectonic evolution of the Zimbabwe Craton. In Fowler, C.M.R., Ebinger, C.J. & Hawkesworth, C.J. (eds). *The Early Earth: Physical, Chemical and Biological Development. Geological Society of London, Special Publications* **199**, 183-211 [Comprehensive overview of the Zimbabwe Craton with data base of all high precision age data].

Johnson & Oliver, (2000). Mesoproterozoic oceanic subduction, island arc formation and the initiation of back-arc spreading in the Kibaran belt of central southern Africa. Evidence from the ophiolite terrane, Chewore Inliers, northern Zimbabwe. *Precambrian Research* **103**, 125-146.

Jolivet, L. & Faccenna, C., (2000). Mediterranean extension and the Africa-Eurasia collision. *Tectonics* **19**, 1095-1106.

Kampunzu, A.B., Tembo, F., Matheis, G., Kapenda, D. & Huntsman-Mapila, P., (2000). Geochemistry and tectonic setting of mafic igneous units in the Neoproterozoic Katangan basin central Africa: implications for Rodinia break-up. *Gondwana Research* **3**, 125-153.

Karner, G.D. & Driscoll, N.W., (1999). Tectonic and stratigraphic development of the West African and eastern Brazilian Margins: insights from quantitative basin modelling. In: Cameron, N.R., Bate, R.H. & Clure, V.S. (eds). *The oil and gas habitats of the South Atlantic. Geological Society of London, Special Publications* **153**, 101-131.

Khan, M.A. et al., (1999). The lithospheric structure of the Kenya Rift as revealed by wide-angle seismic measurements. *Geological Society of London, Special Publications* **164**, 257-269.

Klerkx, J., Liegeois, J-P., Lavreau, J & Claessens, W., (1987). Crustal evolution of the northern Kibaran belt. Eastern and central Africa. In: Kroener, A. (ed.). *Proterozoic lithospheric evolution. American Geophysical Union, Geodynamics series*, **17**, 217-223.

Kreuser, T., (1995). Rift to drift evolution in Permian-Jurassic basins of East Africa. In: Lambiase, J.J. (ed). *Hydrocarbon habitat in rift basins. Geol. Soc. Spec. Publ.* **80**, 297-315.

Kroener, A., (1981). *Precambrian plate tectonics*. Elsevier, Amsterdam, pp. 781. [A good starting point].

Kröner, A. Sacchi, R., Jaekel, P. & Costa, M., (1997). Kibaran magmatism and Pan-African granulite metamorphism in Northern Mozambique: single zircon ages and regional implications. *Journal of African Earth Sciences* **25**, 467-484.

Lavreau, J., (1982). Etude géologique du Haut-Zaïre: genèse et évolution d'un segment lithosphérique Archaïque. Musée Royal de l'Afrique Centrale – Tervuren, Belgique. Annales, Série in-8, *Sciences Géologiques*, no. **88**.

Lavreau, J., (1984). Vein and stratabound gold deposits of northern Zaïre. *Mineralium Deposita*, **19**, 158-165. [Provides useful references to this economically important but otherwise little known terrain].

Mapeo, R. B. M. , Armstrong, R.A. & Kampunzu, A. B., (2001). SHRIMP U-Pb geochronology of gneisses from the gweta borehole, NE Botswana: implications for the Paleoproterozoic Magondi belt in southern Africa. *Geol. Mag.*, 138/3, 299-308

Mattauer, M., Tapponier, P., & Proust, F., (1977). Sur le mécanisme de formation des chaînes intracontinentales. L'exemple des chaînes atlastiques du Maroc. *Bull. Soc. Geol. France*, **7**, 521-526.

Mayer, A., Signigoi, S. & Morais, E., (2000). The Kunene gabbro-anorthosite complex: coalescence of crystal mush intrusions during the early Kibaran. *Am. Geoph. Union Trans.*, **81**, 1248-1249.

McCourt, S., (1995). The crustal architecture of the Kaapvaal crustal block, South Africa, between 3.5 and 2.0 Ga. *Mineralium Deposita*, **30**, 89-97.

McCourt, S., Hilliard, P. Armstrong, R.A. & Munyanyiwa, H., (2001). SHRIMP U-Pb zircon geochronology of the Hurungwe granite northwest Zimbabwe: Age constraints on the timing of the Magondi orogeny and implication for the correlation between the Kheis and Magondi belts. *S. Af. J. Geology*.

Miller, R. McG., (1983). Evolution of the Damara orogen of SW Africa/Namibia. *Geol. Soc. S.Afr. Spec. Publ.* **11**, 115-138.

Morley, C.K., (1995). Developments in the structural geology of rifts over the last decade and their impact on hydrocarbon exploration. In: Lambiase, J.J. (ed.) Hydrocarbon habitat in rift basins. *Geol. Soc. Spec. Publ.* **80**, 1-32.

Notice Explicative de la Carte Géologique du Zaïre au 1/2 000 000, 1974. Département des Mines, République du Zaïre.

Oliver, G. J., (1994). Mid-crustal detachment and domes in the central zone of the Damara orogen, Namibia. *J. Af. Earth Sci.* **19**, 331-344.

Paquette, J-L. & Nedelec, A., (1998). A new insight into Pan-African tectonics in the East-West Gondwana collision zone by U-Pb zircon dating of granites from central Madagascar. *Earth and Planetary Science Letters* **155**, 45-56.

Pinna, P., Cocherie, A., Thieblemont, D., Feybesse, J.-L. & Lagny, P., (1996). Évolution géodynamique du Craton Est-Africain et déterminisme géologique. *Chronique de la Recherche Minière*, **525**, 33-43.

Pinna, P., Cocherie, A., Thieblemont, D. & Jezequel, P., (2000). The Kisii Group of western Kenya: an end-Archaean (2.53 Ga) late orogenic volcano-sedimentary sequence. *Journal of African Earth Sciences*, **30**, 79-91.

Pinna, P., Jourde, G., Calvez, J.Y., Mroz, J.P. & Marques, J.M., (1993). The Mozambique Belt in northern Mozambique: Neoproterozoic (1100-850 Ma) crustal growth and tectogenesis, and superimposed Pan-African (800-550 Ma) tectonism. *Precambrian Research*, **62**, 1-59. [A comprehensive overview].

Porada, H. (1989). Pan African rifting and orogenesis in Southern to Equatorial Africa and Eastern Brazil. *Precambrian Research* **44**, 103-136.

Porada, H. & Berhorst, V., (2000). Towards a new understanding of the Neoproterozoic-Early Paleozoic Lufilian and northern Zambezi belts in Zambia and the Democratic Republic of Congo. *Journal of African Earth Sciences* **30**, 727-771.

Potrel, A., Peucat, J.J. & Fanning, C.M., (1998). Archean crustal evolution of the West African Craton: example of the Amsaga Area (Reguibat Rise), U-Pb and Sm-Nd evidence for crustal growth and recycling. *Precambrian Research*, **90**, 107-117.

Reid, D.I., Welke, H.J., Erlank, A. J. & Moyes, A., (1987). The Orange River Group: a major Proterozoic calcalkaline volcanic belt in the western Namaqua Province, southern Africa. In: T.C. Phaoraoh, R.D. Becknsale and D. Rickard (Ed.), Geochemistry and Mineralization of Proterozoic Volcanic Suites. *Geological Society of London, Special Publications* **33**, 327-346.

Sacchi, R., Cadoppi, P. & Costa, M., (2000). Pan-African reactivation of the Lurio segment of the Kibaran Belt system: a reappraisal from recent age determinations in northern Mozambique. *Journal of African Earth Sciences*, **30**, 629-639.

Salem, M.J., Mouzughy, A.J. & Hammuda, O.S., (1996). *The geology of the Sirt basin*. Elsevier, Amsterdam.

Selley, R.C., (1997). *African basins. Sedimentary basins of the world 3*. Elsevier, Amsterdam, 320p. [Provides a good overview of post Cambrian depositional environments].

Shackleton, R.M., (1993). Tectonics of the Mozambique Belt in East Africa. In: Prichard, H.M., Alabaster, T., Harris, N.B.W., Neary, C.R. (eds). Magmatic Processes and Plate Tectonics. *Geological Society Special Publication*, **76**, 345-362.

Shackleton, R.M., (1996). The final collision zone between East and West Gondwana: where is it? *Journal of African Earth Sciences*, **23**, 271-287.

Stern, R. J., (1994). Arc assembly and continental collision in the neoproterozoic east African orogen: Implications for the consolidation of Gondwanaland. *Ann. Rev. Earth Plante. Sci.* **22**: 319-351. [An excellent overview paper].

Stern, R.J., Kröner, A., Bender, R., Reischman, T. & Dawoud, A.S., (1994). Precambrian basement around Wadi Halfa, Sudan: a new perspective on the evolution of the East Saharan Craton. *Geologische Rundschau* **83**, 564-577.

Stowe, C.W., Hartnady, C.J.H. & Joubert, P., (1984). Proterozoic tectonic provinces of southern Africa. *Precambrian Research*, **25**, 229-231.

Thomas, R.J., Von Veh, M.W. & McCourt, S., (1993). The tectonic evolution of southern Africa: an overview. *Journal of African Earth Sciences*, **16**, 5-24.

Thorweihe, U. & Schandelmeier H. (eds) (1993). *Geoscientific Research in NE Africa*, Balkema, Rotterdam, 776p.

Treloar, P.J., (1988). The geological evolution of the Magondi mobile belt, Zimbabwe. *Precambrian Research*, **38**, 55-73

Trompette, R., (1994). *Geology of Western Gondwana*. Balkema, Rotterdam, pp. 776, 350p. [Provides comprehensive overviews of relatively little known tectonic terrains along the W margin of the African plate including Angola].

Trouw, R.A. & de Wit, M.J., (1999). Relation between the Gondwanide Orogen and contemporaneous intracratonic deformation. *Journal of African Earth Sciences*, **28**, 203-213.

Tucker, R.D., Hadke, M.J., Hamilton, M.A., Ashwal, L.D., LeGrange, M. & Rambeloson, R.A., (1997). U-Pb geochronology and isotope geochemistry of the Archean granite-greenstone belts of Madagascar. *Journal of Geology*.

Turner, B., (1999). Tectonostratigraphic development of the Upper Karoo foreland basin: orogenic unloading versus thermally-induced Gondwana rifting. *Journal of African Earth Sciences*, **28**, 215-238.

Veevers, J.J., Cole, D.I. & Cowan, E.J., (1994). Southern Africa, Karoo basin and cape Fold Belt. *Geol. Soc. Am. Mem.*, **184**, 223-279.

Villeneuve, M. & Dallmeyer, R.D., (1987). Geodynamic evolution of the Mauritinide, Bassaride and Rokelide orogens (West Africa). *Precambrian Research* **37**, 19-28.

Weil, A.B. Van der Voo, R., Mac Niocaill, C. & Meert, J.G., (1998). The Proterozoic supercontinent Rodinia: Palaeomagnetically derived reconstructions for 1100 to 800 Ma. *Earth and Planetary Science Letters* **154**, 13-24.

White, R. & McKenzie, D. (1989). Magmatism at rift zones: the generation of volcanic continental margins and flood basalts. *Journal of Geophysical Research* **94**, 7685-7729.

Williams, H.R. & Culver, S.J., (1982). The roselides of West Africa – Pan African aulacogen or Back-arc basin? *Precambrian Research* **18**, 261-273.

Williams, H.R. & Culver, S.J., (1988). Structural terranes and their relationships in Sierra Leone. *Journal of African Earth Sciences* **7**, 473-477.

Wilson, J.F., Nesbitt, R.W. & Fanning, C.M., (1995). Zircon geochronology of Archaean felsic sequences in the Zimbabwe craton: a revision of greenstone stratigraphy and a model for crustal growth. In: Coward, M.P. & Ries, A.C. (eds) Early Precambrian processes. *Geological Society Special Publications*, **95**, 109-126.

Wilson, M. & Giraud, R., (1992). Magmatism and rifting in Western and Central Africa from Late Jurassic to Recent times. *Tectonophysics* **213**, 1-23.

Wilson, M., (1992). Magmatism and continental rifting during the opening of the South Atlantic Ocean: a consequence of Lower Crataceous super-plume activity? In: Storey, B.C., Alabaster, T. & Pankhurst, R.J. (eds), Magmatism and the causes of continental break-up. *Geol. Soc. Spec. Publ.* **68**, 241-255.

Windley, B.F. (1992). Proterozoic collisional and accretionary orogens. In: *Developments in Precambrian Geology 10, Proterozoic Crustal Evolution*, Ed. K.C. Condie, Elsevier, Amsterdam, 419-445. About the Mauritanides:

Windley, B.F., (1990). *The evolving continents*. John Wiley & Sons, 3rd edition, pp 399.

Biographical Sketches

Paul Dirks is originally from the Netherlands, but received his PhD from Melbourne University in Australia. He has since worked in many countries around the world including China and Antarctica. After almost eight years as lecturer and later professor in the geology department of the University of Zimbabwe in Harare, he took up the position of head of the School of Geosciences at the University of Witwatersrand, Johannesburg in 2002. His principle research interests are structural and metamorphic geology and metallogeny of greenstone belts and high grade gneiss terrains, applied GIS and tectonics. In recent years he has been involved extensively in prospectivity modeling for precious metals in Archaean terrains.

Tom Blenkinsop is originally from the UK but has worked in Africa for many years, initially in Sierra Leone and since 1989 in Zimbabwe where he was Head of Department of Geology at the University of Zimbabwe until his departure to James Cook University, Townsville in 2002. His latest publications comprise investigations of fractal dimensions of quartz grain boundaries and mineral deposits, tectonic controls on materialization, intrusion and folding of granites during regional shortening as well as intrusion-related deformation in granites.

Hielke Jelsma obtained his PhD at the Free University Amsterdam, working on greenstone belts in Zimbabwe. From 1990 until 2000 he lectured at the University of Zimbabwe after which he moved to the Department of Geological Sciences at the University of Cape Town, where he worked as a research associate on problems related to the evolution of the Kaapvaal Craton including kimberlite genesis. In 2003 he took up a position in the Exploration Division of De Beers Consolidated Mines. His main research interests are the tectonic evolution, structural geology and metallogeny of Archean cratons and orogenic belts.