

DESERT RECLAMATION AND MANAGEMENT OF DRY LANDS: FERTILITY ASPECTS

Isam I. Bashour

Faculty of Agricultural and Food Sciences, American University of Beirut, Bliss Street, Beirut, Lebanon.

Keywords: Arid, alkaline (sodic) soils, calcareous, gypsiferous, salt affected soils, salt fertilizer index, semi-arid

Contents

1. Introduction
2. Fertilization of Dry Lands
 - 2.1. Fertilization of Sandy Soils
 - 2.1.1 Properties
 - 2.1.2. Methods of Fertilization
 - 2.2. Fertilization of Clayey Soils
 - 2.3. Fertilization of Calcareous Soils
 - 2.3.1 Properties
 - 2.3.2 Fertilizer Application
 - 2.3.3 Time of Nutrient Application
 - 2.4. Fertilization of Salt Affected Soils
 - 2.4.1 Properties
 - 2.4.2 Quantifying the Salt Effect on Soils
 - 2.4.3 Salt Index of Fertilizers
 - 2.4.4 Managing Salt Affected Soils
3. Conclusions
- Glossary
- Bibliography
- Biographical Sketch

Summary

Arid soils belong to different soil types, depending on parent material and climate, but they share many characteristics regarding their fertility. Arid soils generally support sparse vegetation, because of a shortage of water. They are characterized by a neutral to slightly alkaline pH, have a good structure and are well supplied with K, Ca, Mg, S, B and Mo as a result of not being subjected to leaching. Clearly, their production potential is often considerable, provided sufficient amounts of water are available.

Fertilization with nitrogen (N) is almost always necessary; phosphorus (P) is often needed, especially in calcareous and sandy soils, and fertilization with potassium (K) is required in coarse-textured soils, especially for intensive cultivation and for K-loving crops. Due to the high light intensity in arid regions and limited zinc (Zn) availability, Zn deficiency becomes very common. Supply of iron (Fe) depends to a great extent on the soil pH and lime content. Micronutrients such as copper (Cu), manganese (Mn), boron (B) and molybdenum (Mo) are rarely needed for cropping in medium to heavy

textured soils. Sandy soils in arid regions often require fertilization with most essential nutrients, especially in irrigated agriculture. Acidifying irrigation water has shown to be beneficial and has given good production results in calcareous arid soils.

1. Introduction

Desertification refers to a decline of the biological productivity of arid and semi-arid lands, caused by stress conditions that can be natural or man-made. These conditions, if continued for a long period, lead to ecological and soil degradation.

The primary processes of soil deterioration are: erosion, salinization, and chemical, physical and biological degradation. If these processes continue for a long time they create irreversible changes in the soil and vegetation ecosystem and may lead to the conversion of productive land into a desert. Fertilizer applications can to some extent stop these processes and help in reclaiming the land. For fertilization of salt affected soils, the moisture level should be kept near field capacity to dilute the salt concentration in soil solution. Salt index fertilizers should be used, and fertilizers applied in small quantities in several doses during the season.

Arid soils are the product of several soil forming factors, e.g. parent rock, topography and climate. Water and wind erosion are common in arid regions, and gypsum and salts are present in the soil at variable depths. The factors that affect the fertility levels of arid regions can be summarized as follows:

- Topography: land surrounded by hills receives runoff water and deposits from elevated areas.
- Soil Depth: the depth of the soil profile and eventual presence of an impermeable layer in the soil profile, or of loose sand.
- Soil physical properties: such as soil texture, stoniness and permeability.
- Salinity level: accumulation of salts or gypsum in the soil affects soil productivity and limit plant growth.
- Lime content: soils developed from calcareous rocks contain various levels of CaCO_3 that may affect soil characteristics and management.
- Soil chemical properties: the inherent chemical properties such as clay type, CEC, organic matter content, and plant nutrient levels.

Soils of arid and semi-arid regions are usually low in organic matter. Hence, they are deficient in nitrogen. Since leaching is low, usually these soils are rich in cations and have a basic pH. Their cation exchange capacity (CEC) depends on soil texture, clay type and organic matter level. Sand occupies a major portion of soil components, and the basic soil pH values lead to the precipitation of P, Fe, Zn, Cu and Mn, mostly in forms that are not available for plant uptake.

2. Fertilization of Dry Lands

Soils in dry regions are usually exposed to a hot climate, scanty rainfall and limited leaching. Hence, the potential biomass production is high, provided water is available in satisfactory amounts. The quantity of nutrients available for recycling via plant and

animal residues is not sufficient to compensate for the amounts removed in agricultural products, even in low-productivity situations. Therefore, the use of mineral fertilizers is required for good production.

Soils derived from sandstone become sandy in texture, and those derived from limestone become calcareous. Some dry lands are affected by salts and become saline, sodic or saline-sodic. Each of these soil types (sandy soils, calcareous soils or salt affected soils) requires special management and fertilization.

-
-
-

TO ACCESS ALL THE 21 PAGES OF THIS CHAPTER,
[Click here](#)

Bibliography

Balba, A. (1995). *Management of Problem Soils in Arid Ecosystems*. CRC Lewis Publishers, New York, USA. [A general description of properties of arid soils and explanations of important soil management practices. It covers the desertification problem and how to combat it].

Bashour I. and Nimah M. (1998). *Users Needs for Quality Fertilizer: Types and Formulations*. In: Proceedings Regional Workshop on Guidelines for Efficient Fertilizers Use Through Modern Irrigation, Cairo, Egypt 14 – 16, Dec, 1998, pp 77 – 90. [An overview of the properties and formulations of the fertilizers that are usually applied via the irrigation system].

Bashour I. (2003). *Fertilizer Consumption in the New East: Low Potassium Rates?* In: A.E. Johnston, ed: Potassium and Water Management in West Asia and North Africa, International Potash Institute, Basel, Switzerland. pp 20 – 25. [Review of fertilizer production and consumption in Near Eastern countries with comparison to other countries].

Miller, R and D. Gardiner (1998). *Soils in Our Environment*. 8th ed. Prentice Hall Inc, Upper Saddle River, New Jersey 07458, USA. [A textbook covering a complete spectrum of soil topics; taxonomy, formation, properties, problems, pollution and precision agriculture].

Sys, C. (1975). *Land Evaluation in Europe*. FAO Soils Bulletin 29, FAO, Rome. (Describes European soil types and evaluation systems).

Tabet, A. (1975). *The Effect of Total and Active Calcium Carbonate in the Soil on the Retention of Water and Some Nutrient Elements*. M.Sc. Thesis, Faculty of Agricultural and Food Sciences, American University of Beirut, Beirut, Lebanon. [An account of research on the effect of calcium carbonate on the retention of water, P, K, Fe, Mn, Zn, and Cu in four calcareous clayey soils in Lebanon].

Tisdale, S., Nelson W., Beaton J. and Havlin J. (1993). *Soil Fertility and Fertilizers* 5th ed. Prentice Hall, Upper Saddle River, New Jersey 07458, USA. [Textbook used in many universities to teach students the principles of soil fertility and nutrient management].

NN (1975). *Western Fertilizer Handbook*. California Fertilizer Association, 2222 Watt Avenue, Sacramento, California 95825, USA. [A concise reference explaining technical properties of different fertilizers, grades and formulations. A good source of information for growers and the fertilizer industry].

Wiedenfeld R. (1997). *Sugarcane Responses to N Fertilizer Application on Clay Soils*. Journal of the American Society of Sugar Cane Technologists, Vol. 17, 1997. [An account of research on sugarcane fertilization and response to nitrogen application conducted on clay soils in Texas, USA].

Biographical Sketch

Isam Bashour is an Associate Professor and Chairperson of the Land and Water Resources Department, Faculty of Agricultural and Food Sciences, American University of Beirut, Lebanon. He holds a PhD in Soil Science and Plant Nutrition (1977) from the University of California-Davis.

He has been active for more than twenty five years in teaching, research and consultation in the fields of soil fertilization, reclamation and crop management in arid and semi-arid regions. He developed and formulated acidic fertilizers in different forms: granular, suspension, liquid and soluble powder. His research has mainly focused on soil fertility of various crops and fruit trees with emphasis on fertigation. He also has written many technical reports to FAO, NGOs and national agencies in the Near East.

UNESCO – EOLSS
SAMPLE CHAPTERS