

PAST GLOBAL CRISES

J. Hladil

Institute of Geology, Academy of Sciences of the C.R., Prague, Czech Republic

Keywords: Biocrises, environmental crises, mass extinctions, recoveries, life strategies, evolution, past climatic changes, geochemical anomalies, sediments, bolide impacts, background settings, triggering mechanisms, interactively amplified effects, cascade of causal effects, fossil record, Earth history, Phanerozoic

Contents

1. Introduction to Past Global Crisis
 - 1.1 Particularity of each Crisis and How to Recognize it
 - 1.2 Common Methods for Determining of Past Global Crises
 - 1.3 Duration - A Dating of Global Crises
 - 1.4 Main Differences among the Global Crises
 - 1.5 Geological Record of Global Crises in Sedimentary Beds
 - 1.6 Environmental Perturbations before and after the Global Crisis
 - 1.7 Rare Events or True Catastrophes
 - 1.8 Strong Climatic Control and Dominant Role of the Complex System Behavior
 2. Processes and Effects Connected with Crises
 - 2.1 Fossil Biota, Mass Extinctions and Recoveries
 - 2.2 Geochemical Anomalies and Past Global Crises
 - 2.3 Geological Conditions During the Crises: Sediments and Volcanic Products
 - 2.4 Shock Deformations after Bolide Impacts and other Catastrophic Markers
 3. General Aspects of Natural Global Crises and Interest of Mankind
 - 3.1 Background Setting and Triggering Mechanisms: How did the Crises Work?
 - 3.2 Past Global Crises: A Challenge to Investigation
- Glossary
Bibliography
Biographical Sketch

Summary

Occurrence of global crises is an integral part of the evolution of Earth's biota. These major depressions of life have both aspects of catastrophes, and chances conducive for innovation, which is continued by further evolution. These crises were usually connected with environmental changes. Although these crises are as old as the origin of life, the main information refers to Phanerozoic, between 542 million years ago and the present, when numerous fossils witness to diversification of abundant organisms. Six major crises of Phanerozoic life have been recognized:

- (a) Late Ordovician,
- (b) Late Devonian,
- (c) Terminal Permian,
- (d) Late Triassic,
- (e) Terminal Cretaceous, and

(f) Late Cenozoic to the present, a crisis, which is in its earliest phase.

The largest datasets are related to the late Devonian crisis, the terminal Permian crisis and the terminal Cretaceous crisis, 375, 251, and 65 million years ago respectively. Besides these major global crises, more than one hundred globally traced crises seem to be recorded in the fossil record. Less than one third of these have names according to index fossils, geographical areas or stratigraphic time units, for example, the *murchisoni*, Kacak, or Cenomanian–Turonian crises in Silurian, Devonian, and Cretaceous, respectively.

The global crises of past were triggered by any strong perturbations of the physical, chemical, and biological conditions, which interrupted slowly changing background processes in the pre-existing natural systems on a global or near global scale. The global crisis usually displays a complex structure of developing entities in time and areas. Thus the application of simple models has usually only limited value. The strongest depressions of life after the global crises may persist up to hundreds of thousands of years. However, the first signals of these events and the aftermath of such crises may span several millions of years. The evidence of these global crises in Earth's history is mostly based on fossil organisms, geochemical anomalies and perturbations in sedimentary records. Very close links exist between bio-crises and global change of climate. In sedimentary record, the climatic changes are often reflected by oscillations of sea level that especially reduce extension of shallow shelf seas that strongly affected the feedbacks between biota and environment. Rare, catastrophic events (bolide impacts or huge volcanic eruptions) may contribute to these cataclysms. However, because their distribution displays a weak correlation, if any, with the timing of major global crises, it seems that these rare events need not necessarily result in significant changes to biotic systems. Studies of past ecosystems during crises can provide new insights for environmental management (such as long-term, so-called secular trends) or, more generally, into the behavior of complex systems.

1. Introduction to Past Global Crisis

1.1 Particularity of each Crisis and How to Recognize it

The relatively fast evolution of Earth's life caused that each major crisis of the past was realized on different structure of organisms and had to meet another conditions. In addition, the very complex interactions in ecosystems have also characteristics of chaos, where reactions or survival solutions cannot be copied from one crisis to another but they are usually only canalized to the most probable types of reaction. The global crises gradually emerge from background evolutionary processes and are structured both in time and areas. The main characteristics of culminating stages of these crises are extinction of many normal habitats and depression in diversity of life on a global or near global scale. Because of the structured beginning and aftermath of each global crisis, it is only on decision of each discipline and/or purpose of our studies to select a relevant point in time, and area as a proper start or end of the crisis

1.2 Common Methods for Determining of Past Global Crises

Paleontologists generally agree, that the global crisis starts, when 50% of organic species is extinct without direct replacement by other organisms. The termination of the crisis corresponds to the state, when the recovery processes re-create the diversity values comparable with the pre-crisis situation. Auxiliary boundaries are placed at both pre-existing and subsequent collapses of groups of organisms, as are documented in the fossil record. Similarly, the geochemists can agree that the start of the global crisis is related to the moment when the perturbation in content of some isotope or geochemical-marker (such as ^{13}C) exceeds two times the normally developing values. However, the delimitation of these formal boundaries has only marginal significance, especially in comparison with studies of the entire crisis, which has many links and feedbacks traceable both in time and areas. The global crises can also reach different magnitudes in connection with chances how deeply can proceed the destruction of the pre-crisis organic communities or destruction of the life environment. Each global crisis has a different magnitude. It means practically, how many groups of organisms was affected by step-wise extinction, one after another, or how large part of marine or continental areas was covered by unfavorable conditions (such as presence of anoxic or poisoned water in marine basins or extreme temperature or precipitation values on land).

1.3 Duration - A Dating of Global Crises

The duration of major global crises is also variable. The culminated depressions of diversity with relatively hidden processes of the reconstruction persisted for tens to hundreds of Ka (thousands of years). It departs with timing of singular catastrophes on the Earth, which have not longer aftermath than hundreds to several thousands of years. The dating of the critical events is based on standard stratigraphical methods, which are used in geology for determination of rock ages. These have either sequential, relative character (such as biostratigraphy with its biozones, gamma-ray or magneto-susceptibility stratigraphy, magneto-stratigraphy based on reversals of the Earth's magnetic field, standard chemo-stratigraphy, and stratigraphy based on the changing average contents of isotopes in global oceanic reservoir) or absolute character (the radiometric dating of rocks using the half-time of slowly but continuously transforming, natural radioactive isotopes). As become generally accepted during the last two decades of research, six major crises of Phanerozoic life can be distinguished:

- End-Ordovician, culminating according to most recent radiometric dating approximately 445 million years (Ma) ago,
- Late Devonian, 375 Ma,
- End-Permian, 251 Ma,
- End-Triassic, 200 Ma,
- End-Cretaceous, 65 Ma,
- Late Cenozoic (and the present), a crisis, which is in its early phase.

[International Commission on Stratigraphy - the International Stratigraphic Chart 2004/2008 - <http://www.stratigraphy.org/cheu.pdf>].

The base of the Phanerozoic era, its lowermost Cambrian period, has been recently correlated with the date 542 millions years ago, so that the first major biotic and environmental global crisis (End-Ordovician) emerged 97 millions of years, and the

other occurred irregularly with time scale, after intervals of 70, 124, 51, 135 and 265 Ma. Although the less visible ecological and taxonomical severity of Mid-Carboniferous, Early Ordovician and Jurassic/Cretaceous changes cause these three long or coupled periods, this succession may be indicative of 50-70 million years rhythm. However, a number of secondary wide regional to global crises have been documented in time spans between major crises, and these are at least one decimal order of magnitude shorter than the "Big Six of Crises. "

1.4 Main Differences among the Global Crises

The difference between the pre- and post-crisis life provides another view on the type of the crisis. Especially these crises from the big six that have long post-crisis depression (late Devonian Kellwasser or end-Permian crisis are connected with great changes in the genome and constructions of organisms. Some of the second order global crises are not connected with drastic changes in the composition or construction of successors. The low difference at a glance is connected with the timing of the crisis, which disturbs a major proliferate period in the evolution of life. In Earth's history, when basic part of the life is connected with marine environment, it is usually a time of rising sea levels, warming of the planet, and joining of formerly separated habitats. Middle Devonian Kacak crisis may serve as a good example. It resembles an extensive growth in economy when entire system profits of increased influx of energy and can construct bigger and more effective structures. In such conditions, the depressions are short-lived and aftermath in genome and constructions is small for no reason whatsoever. Conversely, if a similar crisis meet the times of long-term deterioration of life conditions then it usually ends with dramatic consequences for the collapsed structures. These systematic interactions are as obvious as in other systems (economy, system engineering).

1.5 Geological Record of Global Crises in Sedimentary Beds

Worsening of conditions for undisturbed evolution of life has also purely geological signals. Areas on marine shelves are emerged, and are truncated with erosion, so that the sections in sequences of sedimentary strata show significant unconformities covered by so-called condensed sediments, which reflects the low rate of sedimentation. Especially the sedimentological data suggest that many past global crises originated in connection with a fast and extreme oscillation of the sea level, which first consists of a very strong drop of the sea level and then of its strong but short-lived increase. Details of crisis-related intervals in marine sedimentary sequences show the following characteristics:

- The close pre-crisis set of the strata consists of gradually condensed sediments. Truncation of the bedding surfaces is as obvious as direct omission of significant parts of the sedimentary record. Increased proportion of re-worked rock debris contrasts with upward thinning sequence of the beds.
- The erosional or non-sedimentation sequence boundary represents either emergence (close to land) or low influx of rock debris, respective organic skeletal clasts (on the slope of the marine basins). This disconformity between the underlying and

overlying beds reflects the fast sea level lowering, including the subsequent transition from the lowstand to flooding phase. Horizons close to this sequence boundary are rich in condensed products of weathering, where aeolian silt corresponds to arid conditions but clay signalizes the humid episodes. Such sedimentary starving boundaries contain also re-worked volcanic ash, meteoritic material or other background sedimentation components. Typical is enrichment in iron, phosphate, transition metals and rare elements. Good event markers are also spikes in plots of contents of metals as well as increased radioactivity (uranium VI for example).

- The overlying sediments are often breccias or megabreccias. The rapidly flooded mass of coastal weathering products tends to slide into the basin. The overlying set of sediments is usually anoxic because the flooding mass of seawater tends to be stratified and accompanied fertilization of seawater is connected with increased burial of phytoplankton. Cherts and glauconite are common components of in these sediments. As blooms of autotrophic microorganisms consume preferentially the light carbon isotope ^{12}C , the crisis-relevant carbonates are conversely enriched in ^{13}C . However, the total production of carbonates near the crisis is very low, because the reef-building organisms usually collapsed and calcareous shelly organisms, including plankton, are replaced by organisms with silica, phosphate, and/or organic-walled organisms that are accustomed in low-energy, calm water environments.

1.6 Environmental Perturbations before and after the Global Crisis

Instability of the environment near the global crisis horizons in the geological record is documented by rising both the frequency and amplitudes in geochemical, and also geophysical perturbations. There is no reason whatsoever why we would not believe that it is a direct reflection of strongly oscillating climatic conditions, where cold and warm episodes rapidly fluctuate on large areas of the Earth's surface. Considering the energetic requirements of the modern life, where the main power is the energy of the Sun, there are good reasons to suggest that Phanerozoic biotic crises must be preferentially linked with the global climatic changes. According to geological and paleontological dating of the main crisis-related oscillations, many of them can hardly be explained by simple accumulation of the effects of changes in the eccentricity of the Earth's orbit, the tilt of its axis and the precession of the equinoxes. The cycling at about 100 thousands of years, which is recorded in sediments before the late Devonian crisis or in pelagic deposits on the floor of modern (Cenozoic) oceans, especially needs other explanation than the above mentioned Milankovitch parameters of insolation. One plausible explanation is accumulated dust in the stratosphere. This theory is based on the tilt of the Earth's orbit and its periodic contact with different parts of the solar zodiacal ring. Similarly, a contact with hypothetical, extra-solar stardust veils has been suggested as a possible reason for rapid decreases in insolation.

1.7 Rare Events or True Catastrophes

A special chapter of the global crises is connected with so-called rare events or real catastrophes. These catastrophes (bolide impacts or huge volcanic eruptions) may

contribute to biotic crises, but need not necessarily do so, since their temporal distribution displays little, if any, correlation with the timing of past global crises. Periodicity of these volcanic and bolide catastrophic events is slightly documented, and used, to be only hypothesized. It seems that they have for systematics of the major global crises of the past even less importance than orogenic and geotectonic events, which represents the significant endogenic factors on the Earth. Of course, if such catastrophes largely deteriorated the global environment in critical times (as was a reality after the late Devonian Kellwasser crisis, in Early-Middle Famennian, or during the Cretaceous–Tertiary crisis), it had to imply some significant deepening of the crisis and/or its aftermath. The effects of a huge collisions with bolides with similar or coupled extraordinary large volcanic eruptions are compared with a scenario of so-called nuclear winter when the expulsion of the dust into the stratosphere caused a freezing on Earth's surface as well as a consequent collapse of the light-dependent food chains both on land and in the sea. However, the absence of well-structured, gradual beginning of the crisis may serve as a good criterion for separation of the usual past global crises from these abruptly coming catastrophes. Although a big world catastrophe may imply an overkill of many organisms, the organic life can cope with the catastrophic situations much better than with complicated intrinsic problems of the global crises. It is because of the fact that organic communities meet more than not the regional wild fires, volcanic effusions, catastrophic flooding, cool periods, tsunami, and other regional destruction events. Some of them have special strategies how to survive bad climatic years (to burrow in soft rocks, to shelter in rock cavities, producing resistant cysts). Scarce organisms directly calculate on catastrophes since they are part of their long-term life, (for example, some plants in semi-arid environments cannot seed without a regional fire) and some biotic systems do the same, because the rejuvenation of their succession stages helps to their vitality. It is required by their adaptation to quasi-periodical dynamics in ecosystems.

1.8 Strong Climatic Control and Dominant Role of the Complex System Behavior

The climatic change represents a really dominant control on evolution of biocrises, because the connection with glacieustatic change of the sea level has been documented at many of the major crises. Usually, an extreme freezing was followed by an extreme melting of polar ice for no reason whatsoever. Another reason to believe this climatic control is the analysis of energy flows in the biosphere, where the inputs from energy of the Sun are incomparably higher than all other primary resources. However, there are many transformers of this energy (mostly paleogeographical, hydrospheric–atmospheric parameters, and also land coverage by vegetation). These transformers produce complex network of feedbacks, which regularly becomes unstable in times of the coming crises. Possible connection with fluctuating, proper sun's radiation is also promising subject of many present discussions. The connection between organisms and their proximal environment in hydrosphere and atmosphere is also very significant for the biocrises. Many people have understood the passive reaction of the life to be a mirror of the environmental change and that is not true at all. For one thing it not true because the organisms largely produce useful as well as detrimental agents (such as oxygen, humus, heat, and/or organic poisons) and for another thing it is not true because the organisms have to evolve in some canalized ways, where new modifications are based on quality

of their precursors, and many theoretical directions of evolutions are practically impossible. Therefore, we can hardly neglect the intrinsic factors in evolution of life. Many of partial links and feedbacks remain underestimated. It is no doubt, that all effective variables affect the background setting of emerging global crises to modify the chance for their triggering by little, partial mechanisms. This departs from the usual theorizing about the biocrises from the 1960s to 1980s, where a uniform view prevailed (i.e. where processes remain constant over vast stretches of time). The theories about the simple dependence of global biocrises on bolides, volcanoes or other types of the exclusive control appear to be at least doubtful in present times.

-
-
-

TO ACCESS ALL THE **36 PAGES** OF THIS CHAPTER,
[Click here](#)

Bibliography

Alvarez L. W., Alvarez W., Asaro F., and Michel H. (1980). Extraterrestrial cause for the Cretaceous-Tertiary Extinction. *Science* **208**(4448), 1095–1108. [This classic paper describes the iridium anomaly (spike) and suggests collisions with meteoroids, based on the K-T section at Gubbio, Italy.]

Bai S. L., Bai Z. Q., Ma X. P., Wang D. R., and Sun Y. L., eds. (1994). *Devonian Events and Biostratigraphy of South China*. 348 p. Peking University Press, Beijing, China. [This monograph deals with bio-events, chemo-events, Milankovitch cycles, and nickel-episodes of global crises with a large body of biostratigraphy.]

Berger A. and Loutre M. F. (1991). Insolation Values for the Climate of the Last 10 Million Years. *Quaternary Science Reviews* **10**, 297–317. [This paper provides calculated estimates of insolation on the Earth's surface. Visualization by curves was derived using the changing values of the eccentricity of the orbit, tilt of the axis, and precession of the equinoxes.]

Buggisch W., Piepjohn K., Thiedig F., and van Gosen W. (1994). A Middle Carboniferous Conodont Fauna from Blomstrandhalvoya (NW-Spitsbergen): Implications on the Age of Post-Devonian Karstification and the Svalbardian Deformation. *Polarforschung* **62**, 2–3 and 83–90. [This paper documents the mid-Carboniferous drop in sea level that has a chronological connection with the mid-Carboniferous glaciation in South America, Argentina, San Juan, and Mendoza.]

Cejchan P. and Hladil J., eds. (1997). *Final Conference Recoveries 97, Biotic Recoveries from Mass Extinctions*. The Abstract Book, 63 pp. The Field Trip Book, 38 pp. Malak Co. and Eurocongress Center, Czech Republic. [These booklets assembled the opinions of 98 authors, with additional contributors in the field trip book, in an effort to understand the diverse problems of biotic collapse and recovery.]

Chlupac I. and Kukul Z. (1988). Possible Global Events and the Stratigraphy of the Paleozoic, and of the Barrandian (Cambrian-Middle Devonian, Czechoslovakia). *Sbornik Geologických ved, Geologie* **43**, 83–146. [Expanded characteristics of several biocrises within the Devonian are provided, including general comparisons with their sedimentological and eustatic records.]

- Foote M. and Sepkoski J. J. Jr. (1999). Absolute Measures of the Completeness of the Fossil Record. *Nature* **398**, 415–417. [This paper characterizes, in plain form, the problems of inaccuracy and artificial grouping of data in the stratigraphic record.]
- Geldsetzer H. H. J. and Goodfellow W. D. (1987). Sulphur-isotope anomaly associated with the Frasnian-Famennian extinction, Medicine Lake, Alberta, Canada. *Geology* **15**, 393–396 p. [This paper documents perturbation in the $\delta^{34}\text{S}$ sulfur-isotope record.]
- Golovneva L. B. (1994). The flora of the Maastrichtian-Danian deposits of the Koryak Upland, Northeast Russia. *Cretaceous Research* **15**, 89–100. [Terrestrial vegetation on the opposite side of the globe from the Chicxulub crater shows little evidence, if any, for drastic and abrupt changes at the K-T boundary.]
- Hallam A. and Goodfellow W. D. (1990). Facies and geochemical evidence bearing on the end-Triassic disappearance of the Alpine reef system. *Historical Biology* **4**, 131–138. [This paper describes environmental and geochemical factors coincident with the rapid decrease of bio-diversity at the end of the Triassic system.]
- Hart M. B., ed. (1996). *Biotic Recovery from Mass Extinction Events, The Geological Society Special Publication* **102**, 463. London. [This volume summarizes data on many groups of organisms, including foraminifers. The main emphasis is on structures of extinction and recovery processes.]
- Hladil J. and Kalvoda J. (1993). Extinction and Recovery Successions of the Devonian Marine Shoals, Eifelian-Givetian and Frasnian-Famennian Events, Moravia and Bohemia. *Czech Geological Survey, Bulletin* **68**, 13–23. [This paper documents similarities of the Kacak and F-F crises, including anomalous drop—and—rise of sea level, and explains the differing results using the background of climate and biotic evolution. Contact with stardust is suggested as a possible cause of strongly decreased insulation.]
- Hurley N. F. and van der Voo R. (1990). Magneto-Stratigraphy, Late Devonian Iridium Anomaly, and Impact Hypothesis. *Geology* **18**, 291–294. [This paper disputes the bolide interpretation of the F-F iridium anomaly, showing that the duration of this condensed sedimentary sequence extends through several magnetozones.]
- Hsu K. J. (1982). Mass Mortality and its Environmental and Evolutionary Consequences. *Science*, **216**, p. 249–256. [A classic paper dealing with mass mortality, mass extinctions, and fossilization. The concept of the depleted or so-called Strangelove Ocean is derived from this publication.]
- Jablonski D. and Bottjer J. D. (1990). *Onshore-Offshore Trends in Marine Invertebrate Evolution*. In: R. M. Ross W. D. Allmon (eds) *Causes of Evolution—A Paleontological Perspective*. University of Chicago Press. pp. 21–75. [This contribution presents basic paleobiological facts about the dynamics and relationships of the fauna on a gently inclined sea floor.]
- Joachimski M. M. and Buggisch W. (1996). The Upper Devonian Reef Crisis—Insights From the Carbon Isotope Record, in: Reitner J., Neuweiler F., and Gunkel F., (eds), *Global and Regional Controls on Biogenic Sedimentation, Part I. Goettinger Arbeiten in der Geologie und Paleontologie* **2**, p. 365–370. [This paper summarizes data on the widespread occurrence of positive $\delta^{13}\text{C}$ anomalies at the F-F crisis, and includes feedback mechanisms among organic productivity, burial of carbon and atmospheric concentrations of carbon dioxide.]
- Kidwell S. M. (1998). Time-averaging in the marine fossil record: Overview of strategies and uncertainties. *Geobios* **30**, 977–995. [This paper is a conceptual and process analysis of the taphonomy of death assemblages in near-shore and shelf settings as age-mixtures formed over thousands to tens of thousands of years.]
- Krhovsky J. and Cejchan, P. (1995). Cascade of causally linked effects of rapid glaciation–de-glaciation events as possible cause of non-selectivity of mass extinctions. *GeoLines* **3**, p. 27–37. [This work presents references and solutions related to causally linked series of effects.]
- McGhee G.R.Jr., Sheehan P.M., Bottjer D.J. and Droser M.L. (2004). Ecological ranking of Phanerozoic biodiversity crises: ecological and taxonomic severities are decoupled. *Palaeogeography, Palaeoclimatology, Palaeoecology* **211**, 289–297. [The analyses of biota and ecological key-stone organisms indicate that an identification of taxa with high ecological value would be a good strategy to understand (or mitigate?) the ecological effects of the current ongoing loss of global biodiversity.]

McGhee G. R. (1990). The Frasnian-Famennian Mass Extinction Record in the Eastern United States. In: E. G. Kauffman and O. H. Walliser (eds) *Extinction Events in Earth History. Lecture Notes in Earth Sciences* **30**, 161–168. Springer Verlag, Berlin. [This paper examines the detailed evolution of brachiopods and mollusks, among other fauna, in thick sequences of the Frasnian Catskill delta.]

McKinney M. L. (1989). Periodic mass extinctions—product of biosphere growth dynamics? *Historical Biology* **2**, p. 273–287. [This paper provides approaches for the modeling of self-regulated oscillations and extinctions of larger ecosystems, continuing the ideas of Tappan-Loeblich.]

Muller R. A. and MacDonald G. J. (1995). Glacial Cycles and Orbital Inclination. *Nature* **377**, 107–108. [This paper deals with isotopical data and climate variations that show a 100 ka cycling that is caused by the previously ignored tilt of the Earth's orbital plane and not by Milankovitch eccentricity. As the orbit changes, it passes through different parts of the Sun's zodiacal ring, and encounters different regions of stardust.]

Oldfield F. (1999). The Past Global Changes (PAGES) Project: A Personal Perspective. *Quaternary Science Reviews* **18**, 317–320. [This reflection provides good information (and/or deliberation) about major threads and subprojects, methods of the investigation, character of results, as well as the applicability of results, and predictability of processes from this project.]

Playford P. E. and McLaren D. J. (1984). Iridium Anomaly in the Upper Devonian of the Canning Basin, Western Australia. *Science* **226**, 437–439. [This paper documents increased content of iridium and other metals in Early Famennian stromatolitic, and weathering zones on partly emergent, extinct Frasnian reefs.]

Racki G. (1998). Frasnian-Famennian biotic crisis—Under-evaluated tectonic control? *Paleogeography, Paleoclimatology, and Paleoecology* **141**, 177–198. [This provides an application of the Cathles-Hallam model of fluctuating tectonic stress among lithospheric plates, including the massive effusions of basalts that may change sea level and Earth environments. Comparisons are also made with the ideas of F. E. Suess, nineteenth century.]

Raup D. M. and Boyajian G. E. (1988). Patterns of Generic Extinction in the Fossil Record. *Paleobiology* **14**, 109–125. [Plots of global databases on extinctions of organisms are used to search for periodicities. A 26—million—year periodicity and others are postulated.]

Sepkoski J. J. Jr. (1992). Phylogenetic and Ecologic Patterns in the Phanerozoic History of Marine Biodiversity. In Eldredge N. (ed.) *Systematics, Ecology, and Biodiversity Crisis*, pp. 77–100. Columbia University Press, New York. [This essay includes all the classic Sepkoski conclusions based on a structural view of the problem.]

Schindler E. (1990). Die Kellwasser-Krise (Hohe Frasn-Stufe, Ober-Devon). *Goettingen Arbeiten fuer Geologie und Paleontologie* **46**, 1–115. [This monograph summarizes much data on the F-F crisis and defines in detail two phases—the Lower and Upper Kellwasser events.]

Shukolyukov A. and Lugmair G. W. (1998). Isotopic Evidence for the Cretaceous-Tertiary Impactor and Its Type. *Science* **282**, 927–929. [This paper carefully documents the probability that the Chicxulub impactor was a carbonaceous asteroid similar to those which supplied Earth with carbonaceous chondrites of different metamorphic types. The alternative that the bolide was a comet cannot be refuted. Enhanced flux of extraterrestrial ³He in sediments associated with the Chesapeake impact is also discussed.]

Strauss H. and Joachimski M. (2000). Response of the Ocean–Atmosphere System to Past Global Changes. *Chemical Geology, Special Issue* **162**. [This collection of papers discusses phosphorites, sulphur, and carbon isotopes of Neoproterozoic–Cambrian times. Topics include high $\delta^{34}\text{S}$ —values in near-shore areas as a result of overturn in ocean stratification, $\delta^{18}\text{O}$ evidence for Carboniferous and Terminal Permian glaciations, etc.]

Twitchett R. J. and Wignall P. B. (1996). Trace fossils and the aftermath of the Permo-Triassic mass extinction, evidence from Northern Italy. *Paleogeography, Paleoclimatology, and Paleoecology* **124**, 137–152. [This key study illustrates the change of ichnofossils during and after a major crisis. Regional and global controls are examined.]

Urbanek A. (1993). Biotic Crises in The History of Upper Silurian Graptoloids—A Paleobiological Model. *Historical Biology* **7**, 29–50. [An introductory study of Ordovician biocrises, mainly with

hypotheses about the open ocean environment, accompanied by a hypothesis about refugia in “hydrospheric–atmospheric whirls” far offshore.]

Valentine J. W. and Jablonski D. (1993). Fossil Communities, Compositional Variation At Many Time-Scales. In Ricklefs R. E. and Schluter D., eds. *Species Diversity in Ecological Communities—Historical and Geographical Perspectives*. University of Chicago Press. pp. 341–349. [This study exemplifies asynchronous and structured changes of communities with many characteristics of non-competitive diffusion.]

Walliser O. H., ed. (1995). *Global Events and Event Stratigraphy in the Phanerozoic*, 340 pp. Berlin: Springer Verlag. [This is a basic monograph reviewing the results of the international project IGCP 216: Global Bioevents and Mass Extinctions. Ammonite faunas and eustasy are emphasized, together with outlines of ideas about “aging” of systems, and organic groups.]

Wenzel B. C. and Joachimski M. M. (1996). Carbon and oxygen isotopic compositions of Silurian brachiopods (Gotland—Sweden): paleoceanographic implications. *Paleogeography, Paleoclimatology, and Paleoecology* **122**, 143–166. [This paper deals with the composition of Silurian seawater and indicates slightly different relationships to environmental controls than those observed in Devonian and younger rocks.]

Wilde P. and Berry W. B. N. (1984). De-stabilization of the oceanic density structure and its significance to marine extinction events. *Paleogeography, Paleoclimatology, and Paleoecology* **48**, 143–162. [This is a truly classic study focused on mixing in the oceans, overturns of anoxic water and other significant changes in oceanic thermal transfer.]

Biographical Sketch

Jindrich Hladil is engaged with basic research within the framework of the Academy of Sciences in Praha and is teaching as an associated professor at the Masaryk University in Brno. He was born in 1953 and consequently had been living and educated in Brno in Moravia for many years until he moved to Praha. He got through the entire local stairs of academic degrees in geology, paleontology and earth sciences - M. Sc. 1976, R. N. Dr. 1978, Ph. D. 1986 and D. Sc. 2000. His native language is Czech. After short assistant contracts on universities and in companies, a substantial part of life was connected with the Czech Geological Survey, i.e. with drilling survey, mapping of Czech and foreign terrains. Positions of chief stratigrapher and later also head of division can exemplify other relevant activities. Research experience is extremely diversified, being spread from subjects relevant to sedimentology, geochemistry, geophysics, carbonate petrology, stratigraphy, paleobasin analysis in orogens, deformation, and tectonics, but also paleontology, global crises of biota, or entire environmental systems. His special fad is the investigation of coral reefs, especially the tabulate and scleractinian corals. A major part from 200 published papers and communications deals with the Devonian, in both the theoretical, and the application levels.