

## LIMNOLOGY OF THE RIVER NILE

**Gamal M. El-Shabrawy and Mohamed E. Goher**

*National Institute of Oceanography and Fisheries, Fish Research Station, El-Khanater  
El-Kharia, Cairo, Egypt*

**Keywords:** Limnology, climatic change, human impacts, water utilization and River Nile

### Contents

1. Introduction
  2. Watershed of the Nile River
  3. Climate of the Nile Basin
  4. Hydrology and Limnology of the Nile basin
  5. Temperature
  6. Transparency
  7. Dissolved oxygen
  8. Nutrients
  9. Hydrophytes
  10. Phytoplankton
  11. Primary Productivity
  12. Zooplankton
  13. Macrobenthos
  14. Crabs and Shrimps
  15. Fish Fauna
  16. The Nile Birds
  17. A Limnological feature of the main Nile source Lakes
    - 17.1 Lake Tana
    - 17.2. Major Conservation Measures Needed for Lake Tana
    - 17.3 Lake Victoria
  18. Climate change and The River Nile
  19. Effect of Human Impacts on the River Nile
    - 19.1 Deforestation
    - 19.2 Pollution
    - 19.3 Stalination
  20. Utilization of the Nile Basin Water Resource
  21. The Nile Basin Initiative (NBI)
- Glossary  
Bibliography  
Biographical Sketches

### Summary

The Nile Basin covers an area of over 3.12 million km<sup>2</sup>, and a length of about 6800 km, making it the longest in the world. The Basin extends from 4° S to 31° N, stretching over different geographical, climatological and topographical regions. The hydrographical and hydrological characteristics vary greatly over the basin with

abundant rainfall in the headwaters and arid conditions in Sudan and Egypt. Fluctuations in both abiotic and biotic characteristic features of the Nile water are discussed. The river and its lakes are important fisheries resources; navigable waterways; the various dams are generating large amounts of power, irrigation source for increasing agricultural production, underdeveloped hydrocarbon deposits, and seats of ancient and modern civilizations in at least three zones of the Basin. This contributes to river pollution. The full utilization of the water resources of the Nile Basin is an essential prerequisite for the development of its agricultural and industrial potential, besides being basic to the survival of human and animal life.

## 1. Introduction

Water is an important resource for sustaining life. The uses of water are manifold, and include domestic uses, industrial uses such as the production of hydroelectricity, irrigation and animal husbandry. Flowing and stagnant water in water bodies such as rivers and lakes serve as reservoirs for waste disposal. Water security around the world continues to be threatened by population explosion and the rising standards of living, confirming that water is finite and cannot withstand all pressures to its quality, quantity and life-giving support. The intensified use of the world's water resources in the last 100 years has been hastened by technical developments, expansion of energy capture systems and the subtle and direct consequences of population growth at a scale unprecedented in human history. These developments occur amidst the natural variability in soil types, river drainage networks, and climate among the world's watersheds. Rivers are characterized by unidirectional current flow with a relatively high, average flow velocity ranging from 0.1 to 1 m s<sup>-1</sup>. The river flow is highly variable in time, depending on the climatic situation and the drainage pattern.

The Nile basin covers an area of over 3.12 million km<sup>2</sup>, and a length of about 6800 km, longest in the world. The basin extends from 4° S to 31° N, stretching over different geographical, climatological and topographical regions. Besides the two plateaus in Ethiopia and around the equatorial lakes (Victoria, Albert, Kayoga, Edward), the Nile Basin can be considered as a large flat plain, in particular the White Nile sub-basin. The Nile Basin is one of the world's most famous river basins. There is a fascination about the Nile River which has captured human imagination throughout history. Some five thousand years ago a great civilization emerged depending on the river and its annual flooding cycle. Unlike other World Rivers, the Nile is marked by the following characteristics. It passes from south to north; The Nile covers more than 35 latitudes stretching between its sources at the Equatorial Lakes and its mouth in the Mediterranean sea; The Nile water flows into a distance of 2700 kilometers between Atbara River and Mediterranean Sea without receiving any tributaries; the River Nile yield fluctuates from year to another, the lowest recorded yield reached 42 billion cubic meters, while the highest amount of yield reached 150 billion cubic meters. The Niles average annual yield throughout twentieth century is nearly 84 billion cubic meters at Aswan. In spite of its great length and large drainage basin (3,120,000 km<sup>2</sup>, or about 10% of Africa, and affecting 10 nations), it carries relatively little water. Yearly flows over the past century ranged from a low of 42 km<sup>3</sup> in the drought year of 1984 to a high of 120 km<sup>3</sup> for 1916 (Hulme, 1994). This relatively low flow for such a long river is because no water is added to it north of its confluence with the Atbara River, and much

is lost by evaporation. Most other great rivers join with other large streams as they approach the sea, joining their waters into an ever-swelling stream. Instead, the Nile wanders through the largest and most arid region on earth, the Sahara Desert.

The hydrographical and hydrological characteristics vary greatly over the basin with abundant rainfall in the headwaters and arid conditions in Sudan and Egypt. Therefore, although the watershed is large, the portion contributing to stream flow is almost half of the entire basin (only  $1.6 \times 10^6 \text{ km}^2$ ) due to the fact that north of  $18^\circ \text{N}$  latitude, rainfall is almost zero. Precipitation increases towards the headwaters to about 1,200 to 1,600  $\text{mm yr}^{-1}$  on the Ethiopian Plateau and in the region of the Equatorial lakes: Victoria, Albert, Kayoga, and Edward (Mohamed *et al.*, 2005). The seasonal pattern of rainfall follows the Inter-Tropical Convergence Zone (ITCZ), where the dry northeast winds meet the wet southwest winds and are forced upward causing water vapor to condense. The ITCZ follows the area of most intense solar heating and warmest surface temperature and reaches the northerly position of Ethiopian Plateau by late July. The southward shift of the ITCZ results in the retreat of the rainy season towards the central part of the basin after October. Therefore, the monthly distribution of precipitation over the basin shows one single but long wet season over the Ethiopian Plateau and two rainy seasons over the Equatorial Lakes Plateau (Mohamed *et al.*, 2005).

## 2. Watershed of the Nile River

Nile River, with an estimated length of over 6800 km, is the longest river flowing from south to north over 35 degrees of latitude (FAO, 1997). It was long thought that Lake Victoria was its ultimate source, the lake itself is fed by rivers that arise further south, the most important of which is the Kagera. Until recently, it was believed that its tributary, the Luvironza that springs in Tanzania at ca  $4^\circ \text{S}$  was the Nile's ultimate source. The Nile is the only permanent river that manages to cross the Sahara, the largest desert in the world, and reach the Mediterranean Sea, yet its early beginnings are in a Montana equatorial climate and it traverses a series of climatic zones before reaching its delta. Its basin orientation is unique among the major rivers in the world in that it runs almost perfectly from south to north, discharging at  $31^\circ \text{N}$ . Each climate zone which it crosses shows considerable variability in precipitation and run-off (Camberlin, 2009), but over more than half its length it receives less than 150 mm of rain per annum. Its basin is relatively narrow and small ( $3.12 \times 10^6 \text{ km}^2$ ) compared to that of most other large rivers of the world (the Congo, ca  $4 \times 10^6 \text{ km}^2$  according to Bailey, 1986; the Amazon, ca  $7 \times 10^6 \text{ km}^2$  according to Sioli, (1984).

The Nile basin covers the whole of Egypt and the Sudan, one third of Ethiopia, the whole of Uganda, and part of Kenya, Tanzania, Congo, Rwanda and Burundi (Tudorancea & Taylor, 2002). Conventionally, the Nile is divided into a number of sub-basins: the White or Equatorial Nile and its source lakes, the Blue Nile and Lake Tana, and the Main Nile. The River Atbara is often considered a separate, although small, sub-basin (Fig. 1).




Figure 1. The Nile Basin (Source: UNEP/DEWA/GRID-Geneva 1998-09 with copyright permission)

According to Eltahir (2004) the component parts that make up the Nile River's watershed are:

- (1) The Lake Victoria Basin
- (2) East African Lakes below Lake Victoria
- (3) The Bahr el Jebel and the Sudd
- (4) The Bahr el Ghazal Basin
- (5) The Sobat Basin and the Machar Marshes
- (6) The White Nile below Malakal
- (7) The Blue Nile and its Tributaries
- (8) The Atbara and Main Nile to Wadi Halfa
- (9) The Main Nile in Egypt.

The Lake Victoria Basin has an annual rainfall of 1151 mm contributing approximately 122 km<sup>3</sup>/yr to the flow. Its tributaries contribute approximately 276 mm or 22.4 km<sup>3</sup>/yr while evaporation accounts for 1116 mm or 107 km<sup>3</sup>/yr. The resulting outflow from this system is 311 mm or 39.8 km<sup>3</sup>/yr. This provides a relatively steady base flow for the river. The East African Lakes below Lake Victoria include Lake Albert, Lake Kyoga, and Lake Edward. Rainfall contributes to about 10.3 km<sup>3</sup>/yr its tributaries contribute about 10.6 km<sup>3</sup>/yr and 16.3 km<sup>3</sup>/yr is evaporated. Thus, the total resulting outflow is approximately 45 km<sup>3</sup>/yr. The outflow contribution to Nile is dominated by Lake Victoria. This region has also had a dramatic variation in flow level historically.

The Bahr el Jebel and the Sudd receive an annual rainfall of 871 mm while evaporation in the area is much higher at 2150 mm. This area which Nile reaches is the most complex having many seasonal inflows. The high levels of evaporation and transpiration come from the wide distribution of the river area of the water and from the large amounts of vegetation (i.e. *papyrus*). The Jonglei Canal was created in order to provide a more direct way of the water traveling through this region in order to stem the evaporation losses incurred in this area.

The Bahr el Ghazal Basin outflow to the White Nile is almost negligible which amounts to less than 3%. The upper basins have relatively high rainfall, but the river flow spills over into the many flood plain areas resulting in almost all lost to evaporation. The sediment load of these rivers is greater than lake-fed Bahr el Jebel and they have a higher potential for alluvial channels.

The Sobat Basin and the Machar Marshes are also a highly complex area. Most of the runoff develops in the mountains and foothills of Ethiopia. Pibor drains a wide area of plains, but only contributes significantly in times of high rainfall. These reaches also provide about half of the flow for the White Nile, and thus have relatively the same outflow as the Sudd.

The White Nile below Malakal drops 13 m over 840 km. The tributary inflows are sporadic and small while flood plain storage results in delay of outflow and increased loss by evaporation. The Jebel Aulia dam further raised upstream river levels after June 1937. Irrigation and evaporation have led to increased losses.

The Blue Nile and its Tributaries provide a greater part of flow of the Main Nile approximated at 60%. Limited information is known about this area's hydrology, especially in its upper basin within Ethiopia. Its reaches begin in the Ethiopian Plateau at elevations averaging 2000-3000 m, peaking at 4000 m. The terrain consists of very broken and hilly, grassy uplands, swamp valleys, and scattered trees.

The Blue Nile leaves and travels through series of cataracts through the Sudanese's Plains sloping westward from about 700 m. The region it passes through in the plains are covered with Savannah or thorn shrub. Its major tributaries are the Rahad and the Dinder. Along its length are the Roseires Dam (2.4 km<sup>2</sup>) and Sennar Dam (0.5 km<sup>2</sup>). It is the single largest contributor to sediments in the Main Nile, averaging at 140- million tons per year.

The Atbara and Main Nile to Wadi Halfa is the convergence of White Nile and the Blue Nile at Khartoum. The river Atbara is the only major tributary that exists after Khartoum. The river Atbara drains northern Ethiopia (68,800 km<sup>2</sup>) and the mountains north of Lake Tana (31,400 km<sup>2</sup>). Most of its terrain type consists of arid plains dotted with low hills and rock outcrops.

The Main Nile in Egypt is the last major stretch of the Nile before entering the Mediterranean. There are no flows generated below the Atbara confluence (Eltahir, 2004).

### 3. Climate of the Nile Basin

Climate characteristics and vegetation cover in the basin are closely correlated with the amount of precipitation (Fig. 2). Precipitation is to a large extent governed by the movement of the Inter-Tropical Convergence Zone (ITCZ) and the land topography. The main climate zones to be distinguished from North to South are: The Mediterranean climate, a narrow strip around the Nile Delta, followed by the very dry Sahara desert climate down to around 16° N, then a narrow strip of semi desert climate, followed by a wide Savannah climate (poor and tropical Savannah) down to the southern border of Sudan. On the extreme south and southwestern boundary of the basin (around Lake Victoria) tropical and rainforest climates are found. In general, precipitation increases southward, and with altitude (note the curvature of the rain isoheights parallel to the Ethiopian Plateau). Precipitation is virtually zero in the Sahara desert, and increases southward to about 1200–1600 mm/yr on the Ethiopian and Equatorial lake Plateaus. Two oceanic sources supply the atmospheric moisture over the Nile basin; the Atlantic and the Indian Oceans. The seasonal pattern of rainfall in the basin follows the movement of the ITCZ. The ITCZ is formed where the dry northeast winds meet the wet southwest winds. As these winds converge, moist air is forced upward, causing water vapor to condense (Mohamed *et al.*, 2005). El-Tom (1975) claimed that the highest precipitation falls in a region 300 to 600 km south of the surface position of the ITCZ in association with an upper tropospheric tropical easterly jet stream. The ITCZ moves seasonally, drawn toward the area of most intense solar heating or warmest surface temperatures. Normally by late August/early September it reaches its most northerly position up to 20° N. Moist air from both the equatorial Atlantic and the Indian Ocean flows inland and encounters topographic barriers over the Ethiopian



#### 4. Hydrology and Limnology of the Nile basin

The Nile starts from Lake Victoria (in fact from farther south at the Kagera River feeding the lake) and travels north, receiving water from numerous streams and lakes on both sides. In the Sudd, where it takes the name of Bahr el Jebel, the river spills its banks, creating huge swamps where more than half of the river inflow is evaporated. At Lake No, east of Malakal it is joined by the Bahr el Ghazal River draining the southwestern plains bordering the Congo Basin. The Bahr el Ghazal is a huge basin subject to high rainfall over the upper catchments, but with negligible contribution to the Nile flows. Almost all its gauged inflow ( $12\text{Gm}^3$ ) is evaporated in the central Bahr el Ghazal swamps. The Sobat tributary originating from the Ethiopian Plateau and partly from the plains east of the main river joins Bahr el Jebel at Malakal. Downstream this confluence (where it is called the White Nile), it travels downstream a mild slope up to the confluence with the Blue Nile at Khartoum. The Blue Nile originates from Lake Tana located on the Ethiopian Plateau at 1800m above MSL, and in a region of high summer rainfall (1500 mm/yr). The only main tributary of the Nile before it ends up at the Mediterranean Sea is the Atbara River, also originating from the Ethiopian Plateau. The flows originated from the Ethiopian Plateau are quite seasonal and with a more rapid response compared to the flow of the White Nile coming from the Equatorial lakes. Further details on the Nile hydrology can be found in Shahin (1985), Sutcliffe & Parks (1999). The relative contribution to the mean annual Nile water at Aswan of  $84.1\text{Gm}^3$  is approximately  $4/7$  from the Blue Nile,  $2/7$  from the White Nile (of which  $1/7$  from the Sobat), and  $1/7$  from the Atbara River. So the Ethiopian catchments (Sobat, Blue Nile and Atbara River) contribute to about  $6/7$  of the Nile water resources at Aswan. Ten countries share the Nile River: Burundi, Congo, Egypt, Eritrea, Ethiopia, Kenya, Rwanda, Sudan, Tanzania and Uganda. The percentage area of the Nile catchments within each country is: 0.4, 0.7, 10.5, 0.8, 11.7, 1.5, 0.6, 63.6, 2.7, and 7.4%, respectively. The Nile water is vital to the dry countries downstream (Egypt and Sudan), where historically intensive irrigation development exists, and still continues, imposing increasing demands on the Nile water. The upstream countries rely less on the Nile waters, (Mohamed *et al.*, 2005).

#### 5. Temperature

Temperature affects the speed of chemical reactions, the rate at which algae and aquatic plants photosynthesize, the metabolic rate of other organisms, as well as how pollutants, parasites, and other pathogens interact with aquatic residents. Over most of the Nile a prolonged or seasonal thermal stratification is absent, due to wind- and current-induced mixing in shallow waters. Among the headwater lakes, this is applicable to the relatively shallow L. Tana (Morandini, 1940) and L. George (Viner & Smith, 1973). Though the vertical temperature differences are small, in Lake Victoria, an annual cycle of thermal stratification can be distinguished (Fish, 1957). Thermal stratification appears briefly and irregularly in the downstream reservoir at Roseires (Hammerton, 1972a, b), but is annual and prolonged in Lake Nasser-Nubia (Entz, 1976) where it is eliminated by winter cooling and the entry of flood water. Prolonged stratification is lacking in the shallower reservoirs of Gebel (= Jebel) Aulia and Sennar in the Sudan. The physical and chemical characteristic features of the Nile water are shown in Table (1).



-  
-  
-

TO ACCESS ALL THE 36 PAGES OF THIS CHAPTER,  
<http://www.eolss.net/Eolss-sampleAllChapter.aspx> Visit:

### Bibliography

Abd El-Karim, M.S. (1999). Phytoplankton Dynamic and its productivity in Damietta Branch. M. Sc. Thesis, Fac. of Girls, Ain Shams University, 202pp [M Sc. Thesis dealing with chemical features and phytoplankton abundance, seasonal variation and community structure in Demietta Nile branch]

Abd El-Mola, H.R. (2009). Ecological studies on planktonic and epiphytic micro invertebrates in Lake Nasser, Egypt. Ph.D. Thesis, Fac. of Sci., Benha Univ. Egypt. 207 pp [This is a case study reporting the relation between water quality and zooplankton Dynamics in Lake Nasser]

Abdel-Hamid, M. I. (1991). Phytoplankton and water quality of the River Nile in Egypt. Ph. D. Thesis, El-Mansoura University, Egypt, 346 pp. [This is a case study reporting the relation between water quality and Phytoplankton Dynamics in River Nile at Egypt ]

Abdel-Monem, A.M.A. (1995). Spatial distribution of phytoplankton and primary productivity in Lake Nasser, Ph.D. Thesis, University Collage for Girls, Ain Sham Univ., 161pp. [Ph. D. Thesis dealing with chemical features, Primary productivity and phytoplankton abundance, seasonal variation and community structure in Lake Nasser]

Abdel Satar, A.M. (2005). Water quality assessment of river Nile from Idfo to Cairo. *Egypt. J. of aquatic res.* 31:200-223. [A recent assessment of the concentrations of oxygen, anion, cation and nutrient]

Abdin, G. (1948a). The conditions of growth and periodicity of the algal flora of the Aswan reservoir (Upper Egypt). *Bulletin of the Faculty of Sciences*, Egyptian University **27**: 157–175. [Detailed study of algal growth in Lake Nasser]

Abdin, G. (1948b). Physical and chemical investigations relating to algal growth in the River Nile, Cairo. *Bulletin de l' Institut d' Egypte* 29: 19–44. [Summarizes the effect of water quality on algal growth]

Abu-Gideiri, Y. B. and Ali, M. T. (1975). A preliminary biological survey of Lake Nubia. *Hydrobiologia* 46: 535–541.

Ali, M. M. and Sultan, M. A. (2006). Expansion of *Myriophyllum spicatum* (Eurasian water milfoil) into Lake Nasser, Egypt: Invasive capacity and habitat stability. *Aquatic Botany*, 84 : 239–244

Ali, M. M., Hamad, A. M., Springuel, I. V. and Murphy, K. J. (1995). Environmental factors affecting submerged macrophyte communities in regulated water bodies in Egypt. *Arch. Hydrobiol.* 133: 107-128

Ali, O. M. M. (2009). Aquatic Plants of the Sudan . 479-494. In H. J. Dumont (ed.), *The Nile. Monographiae Biologicae*, Vol. 89: 479-494. Springer, Dordrecht. [A multi-author monographs book, in which thirty-seven authors have taken up the challenge, and have written chapters cover the origin, environments, limnology and human use of the River Nile]

Allan, J. A., (2009). Nile Basin asymmetries: a closed fresh water resource, soil water potential, the political economy and Nile Tran boundary hydro politics. In H. J. Dumont (ed.), *The Nile. Monographiae Biologicae*, Springer, Dordrecht. [A multi-author monographs book, in which thirty-seven authors have taken up the challenge, and have written chapters cover the origin, environments, limnology and human use of the River Nile]

Aloo, P.A., (2003). Biological diversity of the Yala Swamp lakes, with special emphasis on fish species composition, in relation to changes in the Lake Victoria Basin (Kenya): threats and conservation measures. *Biodiversity and Conservation* 12: 905 – 920, 2003.

- Appleton, C. (2003). Alien and invasive fresh water Gastropoda in South Africa. *African Journal of Aquatic Science* 28: 69–81.
- Attia, B., (2008). Assessment of Vulnerability and Adaptation of Water Resources to Climate Change in Egypt. 13th IWRA world water congers, 1-4/9/2008 Montpellier, France.
- Bacci, G. (1951–1952). Elementi per una malacofauna del' Abissynia e delle Somalia. *Annales Museo Civico di Storia Naturale Giacomo Doria* 65: 1–44.
- Beyene, T., Lettenmaier, D.P. and Kabat, P. (2010). Hydrologic impacts of climate change on the Nile River basin: Implications of the 2007 IPCC climate scenarios, *Climatic Change* 100:433–461
- Bishai, H. A., Abdel-Malek S. A. and Khalil, M. T. (2000). *Lake Nasser*. Publications of National Biodiversity Unit, Egypt, pp 11–577.
- Bishai, H. M. (1962). The water characteristics of the Nile in the Sudan with a note on the effect of Eichhornia crassipes on the hydrobiology of the Nile. *Hydrobiologia* 19: 357–382.
- Brook, A. J. & J. Rzóska, (1954). The influence of the Gebel Aulyia dam on the development of Nile plankton. *Journal of Animal Ecology* 23: 101–114.
- Brown, D. S. (1994). *Freshwater Snails of Africa and Their Medical Importance*. Second revised edition. Taylor & Francis, London, pp. 609.
- Calder, R.I., Hall, R., Bastable, H., Gunston, H., Shela, O., Chirwa, A. and Kafundu, R. (1995). The impact of land use change on water resources in sub-Saharan Africa: a modeling study of Lake Malawi. *Journal of Hydrology* 170: 123–135.
- Camberlin, P. (2009). Nile Basin Climates. In H. J. Dumont (ed.), *The Nile. Monographiae Biologicae*, Vol. 89: 307–333. Springer, Dordrecht. [A multi-author monographs book, in which thirty-seven authors have taken up the challenge, and have written chapters cover the origin, environments, limnology and human use of the River Nile]
- Camberlin, P. (1997). Rainfall anomalies in the Source Region of the Nile and their connection with the Indian Summer Monsoon, *J. Climate*, 10, 1380–1392.
- Conway, D. and Hulme, M. (1993). Recent fluctuations in precipitation and runoff over the Nile sub-basins and their impact on main Nile discharge. *Climatic Change* 25:127 151.
- Conway, D. and Hulme, M. (1996). The Impacts of Climate Variability and Future Climate Change in the Nile Basin on Water Resources in Egypt. *International Journal of Water Resources Development*, 13(3): 277-296.
- Conway, D. (2005). From headwater tributaries to international river: observing and adapting to climate variability and change in the Nile Basin. *Global Environemental Change* 15: 99-114
- Cumberlidge, N. (2009). Freshwater crabs and shrimps (Crustacea: Decapoda) of the Nile basin. In H. J. Dumont (ed.), *The Nile. Monographiae Biologicae*, Vol. 89: 547–561. Springer, Dordrecht. [A multi-author monographs book, in which thirty-seven authors have taken up the challenge, and have written chapters cover the origin, environments, limnology and human use of the River Nile]
- De Nie, H.W. (1987). The decrease in aquatic vegetation in Europe and its consequences for fish populations. EIFAC/CECPI Occasional Paper No. 19. FAO, Rome. 52p.
- De Ridder, M. (1984). A review of the rotifer fauna of the Sudan. *Hydrobiologia* 110:113-130
- Dejen, E., Vijverberg, J. Nagelkerke L.A.J. and Sibbing, F.A. (2004). Temporal and spatial distribution of microcrustacean zooplankton in relation to turbidity and other environmental factors in a large tropical lake (L. Tana, Ethiopia). *Hydrobiologia* 513: 39–49.
- Dejen, E., M. de Graaf, L. A. L. Nagelkerke, T. Wudneh, J. W. M. and Sibbing, F. A. (2006). Lake Tana fishery and its sustainable development (in Amharic). Amhara Region Agricultural Research Institute & Department of Animal Sciences and Wageningen University. Tis Abay Printing Press, Bahir Dar, Ethiopia: 84 pp.
- Dumont, H. J., (1986a). Zooplankton of the Nile system. In B. R. Davies & K. F. Walker (eds), *The Ecology of River Systems*. Kluwer, Dordrecht, pp. 75–88.

- Dumont, H. J., (1986b). The Nile River system. In B. R. Davies & K. F. Walker (eds), *The Ecology of River Systems*. Kluwer, Dordrecht, pp. 61–74.
- Dumont, H. J., (2009). A description of the Nile basin, and a synopsis of its history, ecology, biogeography and natural resources. In H. J. Dumont (ed.), *The Nile. Monographiae Biologicae*, Vol. 89: 1–21. Springer, Dordrecht. [A multi-author monographs book, in which thirty-seven authors have taken up the challenge, and have written chapters cover the origin, environments, limnology and human use of the River Nile]
- El-Ayouty, E. Y., (1976). River phytoplankton. Annual Report and River Nile project, part II. River Ecosystem studies. *Egyptian Academy of Sciences and Technology*, Cairo.
- El-Shabrawy, G. M. and Dumont, H. J., (2003). Seasonal and spatial variation of zooplankton in the coastal zone of the main channel and main khors of Lake Nasser, with a note on the lake fisheries. *Hydrobiologia* 491: 119–132
- El-Shabrawy, G.M. and Khalifa, N. (2002), "Zooplankton abundance and community structure in the northern part and estuary of Rosetta Nile Branch in relation to some environmental variables", *Egyptian Journal of Aquatic Biology and Fisheries* Vol. 6 No.4: 69-90.
- Elshamy, M.E., (2000). Impacts of climate change on Nile flows. Diploma of Imperial College (DIC) Thesis, Imperial College London, London.
- Elster H. J. and Jensen, K. W. (1961). Limnological and Fishery Investigation of the Nozha Hydrodrome. Alexandria, Egypt. *Notes and Memoirs Hydrobiological Department*, Alexandria 43: 99 pp.
- Eltahir, E.A., (2004). "Sustainability Considerations of Big Dams: Merowe, Nile Basin," Paris, Anthony, Teresa Yamana, and Susan Young, Working paper
- Entz, B., (1976). Lake Nasser and Lake Nubia. In J. Rzóška (ed.), *The Nile, biology of an ancient river. Monographia Biologicae* 29. Junk, The Hague, pp. 271–298.
- FAO, (1997). Irrigation potential in Africa: A basin approach. The Nile basin FAO Land and Water Bulletin 4. Land and Water Development Division. FAO, Rome.
- Fish, G. R., (1957). A seethe movement and its effect on the hydrology of Lake Victoria. *Fisheries Publications*, London 10: 1–68.
- Fishar, M.R. and W. P. Williams (2006). A feasibility study to monitor the macroinvertebrate diversity of the River Nile using three sampling methods *Hydrobiologia* 556:137–147.
- Flohn, H., and Fraedrich, K., (1966). Tagesperiodische Zirkulation und Niederschlagsverteilung am Victoria-See (Ostafrika). *Meteorologische Rundschau* 19: 157–165.
- Goher, M., E. and Ali M. H. (2009). Monitoring of water quality characteristics and some heavy metals in water, sediment and macrophytes in Main Khors of Lake Nasser (Egypt). *J. Egypt. Acad. Soc. Environ. Develop.*, 10 (4):109-122
- Golterman, H. L. (1973). Natural phosphate sources in relation to phosphate budgets: a contribution to the understanding of eutrophication. *Water Research* 7: 3–17.
- Green, J. (2009). Birds of the Nile. In H. J. Dumont (ed.), *The Nile. Monographiae Biologicae* Vol. 89: 705–720. Springer, Dordrecht.
- Greenwood, P. H. (1974). The cichlid fishes of Lake Victoria, East Africa: the biology and evolution of a species flock. Bulletin of the British Museum (Natural History), *Zoology Series, Supplement* 6: 1–134.
- Habib, O. A. (2000). Primary production of the phytoplankton. In J. F. Craig (ed.), *Sustainable Fish Production in Lake Nasser: Ecological Basis and Management Policy*. ICLARM Conference Proceedings 61: 184 pp
- Hammerton, D. (1970). Water characteristics and phytoplankton production. Eleventh Annual Report 1963–1964, Hydrobiological Research Unit, University of Khartoum, pp. 5–12.
- Hammerton, D. (1972a). Survey of work in progress. Blue Nile survey. River Nile-Lake Nubia. Fifteenth Annual Report 1967–1968, Hydrobiological Research Unit, University of Khartoum, pp. 5–14.

- Hammerton, D. (1972b). Survey of work in progress. Blue Nile survey. White Nile survey. Sixteenth Annual Report 1968–1969, Hydrobiological Research Unit, University of Khartoum, pp. 4–9.
- Harper D.M. and Mavuti K.M. (1996). Freshwater wetlands. In: McClanahan T.R. and Young T.P. (eds), *East African Ecosystems and Their Conservation*. Oxford University Press, Oxford, UK, pp. 217– 239.
- Hassan, F. (1981). Historical Nile floods and their implications for climatic change. *Science* 212:1142–1145
- Hecky, R. E. (1993). The eutrophication of Lake Victoria. *Verhandlungen der Internationalen Vereinigung für Limnologie* 25: 39–48.
- Herdendorf, C. E. (1990). Distribution of the world's large lakes. In M. M. Tilzer & C. Serruya (eds), *Large Lakes Ecological Structure and Function*. Springer-Verlag, Berlin, pp. 3–38.
- Heywood V.H. and Watson R.T. (1995). *Global Biodiversity Assessment*. Cambridge University Press, Cambridge, UK, 1140 pp.
- Hulme, M. (1994). Global climate change and the Nile basin. In (Howell & Allan 1994)
- Hulme, M., Doherty R., Ngara T. and New M. (2005). *Global Warming and Africa: Climate Change and Africa: A reassessment*, Cambridge University Press, 338pp.
- Hurni, H. (1999). Sustainable management of natural resources in African and Asian mountains. *Ambio* 28: 382–389.
- Hynes, H. B. N. (1970). *The Ecology of Running Waters*, University of Toronto Press, Toronto, Canada, 555 pp.
- Ibrahim, A. M. (1984). The Nile: description, hydrology, control and utilization. *Hydrobiologia* 110: 1–13.
- Ibrahim, M.A. Mousa, A.A. and EL-Bokty, E.E. (1997). Environmental factors affecting the abundance and distribution of Macro-benthic organisms in Manzalah Lake , Egypt. *Bull. Nat. Inst. Oceanogr. & Fish.*, ARE 23: 315-331.
- Imoobe, T.O.T. and Christopher, A.O. (2010). Spatial variations in the composition and abundance of zooplankton in the Bahir Dar Gulf of Lake Tana, Ethiopia. *African Journal of Ecology* 48 (1): 72-77.
- Johnson, T. C., Kelts, K. and Odada, E. (2000). The Holocene history of Lake Victoria. *AMBIO* 29:2–11.
- Karani, P. W. (2005). Status of water quality monitoring in the Kenyan portion of Lake Victoria Basin. Nile Basin Initiative and Nile Trans-boundary Environmental Action Project, 2005. Nile Basin National Water Quality Monitoring Baseline Study Report for Kenya
- Kayombo S. and Jorgensen S.E. (2004). Experience and lessons learned brief for Lake Victoria. : [http://www.worldlakes.org/uploads/Victoria\\_draft\\_10.23.03.pdf](http://www.worldlakes.org/uploads/Victoria_draft_10.23.03.pdf)
- Kifle, D. and Belay, A. (1990). Seasonal variations in phytoplankton primary production in relation to light and nutrients in Lake Awassa, Ethiopia *Hydrobiologia* 196: 217-227.
- Korovchinsky, N. M. (2000). Species richness of pelagic Cladocera of large lakes in the eastern hemisphere. *Hydrobiologia* 434: 41–54.
- Kudhongania, A. W. and Cordone, A. J. (1974). Batho-spatial distribution patterns and biomass estimates of the major demersal fishes in Lake Victoria. *The African Journal of Tropical Hydrobiology and Fisheries* 3: 15–31
- Kurdin, V. P. (1968). Data on hydrological and hydrochemical observations on the White Nile. *Informatsionny Byulleten Biologiya Vnutrennikh Vol 2*: 49–56 (Russian)
- Lamb, H. F., Bates R., Marshall M. H., Umer M., Davies S. J. and Toland, H. H. (2007). Pleistocene desiccation of Lake Tana, source of the Blue Nile. *Quaternary International* 167–168, 226.
- Latif, A. F. A., and Elewa, A. A. (1980). Physico-chemical characteristics of Lake Nasser and Lake Nubia. Report on Surveys to Lake Nasser and River Nile Project. Academy of Scientific Research and Technology, Cairo, Egypt, 105 pp

- Lehman, J. T., R. Mugidde, R., & Lehman, D. A. (1998). Lake Victoria plankton ecology: Mixing depth and climate-driven control of lake condition. In J. T. Lehman (ed.), *Environmental Change and Response in East African Lakes*. Kluwer, Dordrecht, pp. 99–116.
- Lehman, J. (2009). Lake Victoria. In H. J. Dumont (ed.), *The Nile. Monographiae Biologicae*, Vol. 89: 215–241. Springer, Dordrecht. [A multi-author monographs book, in which thirty-seven authors have taken up the challenge, and have written chapters cover the origin, environments, limnology and human use of the River Nile]
- Merid, F., 2005. *Nile basin national water quality monitoring baseline study report for Ethiopia*. Nile Basin Initiative and Nile Trans-boundary Environmental Action Project.
- Meybeck M. (1996). River water quality. Global ranges, time and space variability's, proposal for some redefinitions., *Verh. Internat. Verein. Limnol.* 26: 81-96.
- Miracle, M. and Serra, M. (1989). Salinity and temperature influence rotifer life history characteristics. *Hydrobiologia* 186/187: 81-102.
- Mohamed, Y. A., van den Hurk, B. J. J. M., Bastiaanssen W. G. M., and Savenije, H. (2005). Hydroclimatology of the Nile: results from a regional climate model. *Hydrology and Earth System Sciences* 9: 263–278.
- Molla, M. and Menelik, T. (2004). Environmental impact assessment for unusual reduced water level of Lake Tana. In Proceedings of the Symposium on Lake Tana Watershed Management. Lake Net, USA: 35–48.
- Monakov, A. V. (1969). The zooplankton and zoobenthos of the White Nile and adjoining waters in the Republic of the Sudan. *Hydrobiologia* 33: 161–185.
- Monod, T., (1980). Décapodes. In J-R. Durand & C. Lévêque (eds), *Flore et Faune Aquatiques de l'Afrique Sahelo-Soudanienne* 1. ORSTOM, Paris, pp. 369–389.
- Mwebaza-Ndawula, L. (1994). Changes in relative abundance of zooplankton in northern Lake Victoria, East Africa. *Hydrobiologia* 272: 259–264.
- Newell, B. S. (1960). The hydrology of Lake Victoria. *Hydrobiologia* 15: 363–383.
- Nicholson, S. E., (1996). A review of climate dynamics and climate variability in eastern Africa, in: *The limnology, climatology and paleoclimatology of the East African lakes*, edited by: Johnson, T. C. and Odada, E., 25–56, Gordon and Breach, Amsterdam, 1996.
- Nyssen, J., Poesen J., Moeyersons J., Deckers J., Haile M. and Lang A. (2004). Human impact on the environment in the Ethiopian and Eritrean highlands – a state of the art. *Earth-Science Reviews* 64: 273–320.
- Ogari, J. (2001). Impact of exotic fish species and invasive water weeds such as the water hyacinth on Lake Victoria fisheries, Paper presented at LVEMP conference, Kisumu, Kenya.
- Ogutu-Ohwayo, R. (2003). The Fisheries of Lake Victoria; Harvesting Biomass at the expense of Biodiversity.
- Okonga, J. R., Sewagudde, S. M. , Mngodo, R. J. , Sangale, F. D. , Mwanuzi F. L., and Hecky, R. E. (2006). Water balance for Lake Victoria. In E. O. Odada & D. O. Olago (eds), Proceedings of the 11th World Lakes Conference, Nairobi, Kenya, Vol. 2, pp. 47–56.
- Otim, M., (2005). *Baseline study of the status of water quality monitoring in Uganda*. Nile Basin Initiative and Nile Trans-boundary Environmental Action Project. Nile Basin National Water Quality Monitoring Baseline Study Report for Ugand
- Piper, B. S., Plinston D. T., and Sutcliffe, J. V. (1986). The water balance of Lake Victoria. *Hydrological Sciences Journal* 31: 25–47.
- Rabeh, S. A. (2009). Bacteria and Viruses in the Nile In H. J. Dumont (ed.), *The Nile. Monographiae Biologicae*, Vol. 89: 407–430. Springer, Dordrecht [A multi-author monographs book, in which thirty-seven authors have taken up the challenge, and have written chapters cover the origin, environments, limnology and human use of the River Nile]

- Rzóska, J. (1974). The Upper Nile swamps, a tropical wetland study. *Freshwater Biology* 4: 1–30.
- Samaan, A. A., (1971). Report on trip to Lake Nasser to investigate its primary productivity during March 1971, Report to FAO/UN Lake Nasser Development Centre, Aswan, 11 pp.
- Shahin, M. (1985). *Hydrology of the Nile Basin* (Developments in Water Science 21). Elsevier, Amsterdam, Netherlands
- Shehata, S. A. (1968). Studies on plankton recovered from the Nile Water at Cairo district. M.Sc. Thesis, Cairo University, Egypt.
- Shehata, S.A. and Bader, S.A. (1985). Effect of Nile River water quality on algal distribution at Cairo, Egypt, *Environ. Internat.*, 11: 465-474.
- Shiel, R.J.; Walker, K.F. and Williams, W. (1982). Plankton of the lower River Morry, South Australia. *Ans. J. Mar. Freshwat. Res.*, 33: 210 – 227.
- Sibbing, F.A., Nagelkerke L.A.J., Stet R. J.M. and Osse, J.W.M. (1998). Speciation of endemic Lake Tana barbs (Cyprinidae, Ethiopia) driven by trophic resource partitioning; a molecular and ecomorphological approach. *Aquatic Ecology* 32: 217–227.
- Sinada, F. and Abdel Karim, A. G. (1984a). Physical and chemical characteristics of the Blue Nile and the White Nile at Khartoum. *Hydrobiologia* 110: 21–32.
- Sinada, F. and Abdel-Karim, A.G., (1984b). Quantitative Study of the Phytoplankton in the Blue and White Niles at Khartoum. *Hydrobiologia* 110: 47-55.
- Sinada, F. and Abdel-Karim, A.G., (1984c). Primary Production and Respiration of the Phytoplankton in the Blue and White Niles at Khartoum. *Hydrobiologia* 110: 57-59
- Sioli, H. (1984). *The Amazon: Limnology and landscape ecology of a mighty tropical river and its basin*. Dr. W. Junk Publishers, Dordrecht 761p.
- Sobhy, E. H. M. (1999). Effect of industrial waste of Iron and Steel Factories on Nile phytoplankton communities and productivity at Helwan. M.Sc. Thesis, Menofya University, Egypt.
- Springuel, I. V. and Murphy, K. J. (1990). Euhydrophytes of Egyptian Nubia. *Aquatic Botany*, 37: 17-25.
- Stern, N. (2007). *The Economics of Climate Change: The Stern Review*. Cambridge
- Sutcliffe, J. V. and Parks, Y. P. (1999). *The Hydrology of the Nile*. IAHS Special Publ. no. 5
- Talling J. F. (2009). Physical and Chemical Water Characteristics. In H. J. Dumont (ed.), *The Nile. Monographiae Biologicae*, Vol. 89: 367-395. Springer, Dordrecht. [A multi-author monographs book, in which thirty-seven authors have taken up the challenge, and have written chapters cover the origin, environments, limnology and human use of the River Nile]
- Talling, J. F. and Rzóska, J. (1967). The development of plankton in relation to hydrological regime I. the Blue Nile. *Journal of Ecology* 55: 637–662.
- Talling, J. F. (1976): Water characteristic in the Nile biology of an ancient river. (ed. Rzoska, J.) junk publisher, The Huage, 357-384.
- Talling, J.F., and Talling, I.B. (1965). The chemical composition of African lake waters. *International Revue der gesamten Hydrobiologie* 50: 421–463.
- Talling, J.F. (1957a). Diurnal changes in stratification and photosynthesis in some tropical African waters. *Proceedings of the Royal Society of London* 147: 57–83.
- Talling, J.F. (1957b). The longitudinal succession of water characteristics in the White Nile. *Hydrobiologia* 11: 73–89.
- Talling, J.F. (1957c). Some observations on the stratification of Lake Victoria. *Limnology and Oceanography* 2: 213–221
- Talling, J. F. (1963). Origin of stratification in an African Rift lake. *Limnology and Oceanography* 8: 68–78.

- Talling, J. F. (1966). The annual cycle of stratification and phytoplankton growth in Lake Victoria (East Africa). *Internationale Revue der gesamten Hydrobiologie* 51: 545–621.
- Talling, J. F., Sinada, F., Taha, O. E., and Sobhy E. M. (2009). Phytoplankton: Composition, Development and Productivity Jack. In H. J. Dumont (ed.), *The Nile. Monographiae Biologicae*, Vol. 89: 430-462. Springer, Dordrecht. [A multi-author monographs book, in which thirty-seven authors have taken up the challenge, and have written chapters cover the origin, environments, limnology and human use of the River Nile]
- Teshale, B. (2003). Influence of sediment on physico-chemical properties of Lake Tana. Workshop 'Fish and Fisheries of Lake Tana: *Management and Conservation*'. 6–8 October 2003, Bahir Dar, Ethiopia.
- Teshale, B., Lee, R. and Zawdie, G. (2001). Development initiatives and challenges for sustainable resource management and livelihood in the Lake Tana region of Northern Ethiopia. In A. B. Dixon, A. Hailu & A. P. Wood (eds), *Proceedings of the Wetland Awareness Creation and Activity Identification Workshop in Amhara National Regional State*. January 23rd 2001, Bahar Dar, Ethiopia, 33–43.
- Tewabe, D., Muhammed S. and Abdissa B. (2005). Distribution and abundance of macro-benthic and weed-based faunas in the northern part of Lake Tana. Internal Report ARARI, Bahir Dar, Ethiopia, 14 pp
- Toulibah, H. E., Abdel-Monem A. M., and Gaballah, M. M. (2000). Statistical analysis of phytoplankton structure and environmental variables in Lake Nasser – Egypt: 1 – Along the main channel. *Al-Azhar Journal of Microbiology* 49: 189–202.
- Tucker C. J. and Nicholson, S. E. (1999). Variations in the size of the Sahara Desert from 1980 to 1997. *AMBIO* 28: 587–91.
- Tudorancea, C., and Taylor, W. D. (2002). *Ethiopian Rift Valley Lakes*, 289 pp. Backhuys, Leiden.
- UNEP (1991). Environmental Data Report. 3rd edn. Basil Blackwell Scientific Publication, Oxford, UK.
- Van Damme, D. and Van Bocxlaer B. (2009). Freshwater molluscs of the Nile Basin, past and present. In H. J. Dumont (ed.), *The Nile. Monographiae Biologicae*, Vol. 89: 585–629. Springer, Dordrecht. [A multi-author monographs book, in which thirty-seven authors have taken up the challenge, and have written chapters cover the origin, environments, limnology and human use of the River Nile]
- Vijverberg, J., Sibbing F.A., and Dejen, E. (2009). Lake Tana: Source of the Blue Nile. In H. J. Dumont (ed.), *The Nile. Monographiae Biologicae*, Vol. 89: 163–192. Springer, Dordrecht. [A multi-author monographs book, in which thirty-seven authors have taken up the challenge, and have written chapters cover the origin, environments, limnology and human use of the River Nile]
- Viner, A. B. & Smith, I. R. (1973). Geographical, Historical and Physical aspects of Lake George. *Proc. R. Soc. Lond. B*. 184: 235-270
- Viner, A.B. (1972). Responses of a mixed phytoplankton population to nutrient enrichments of ammonia and phosphate, and some associated ecological implications. *Proceedings of the Royal Society of London B* 183: 351–370.
- Williams, M.A. (2002). Desertification. In I. Douglas (ed.), *Encyclopedia of Global Environmental Change, Volume 3: Causes and consequences of global environmental change*. Wiley, Chichester, pp. 282–290.
- Williams, M.A. (2003a). Desertification in Africa, Asia and Australia: human impact or climatic variability. *Annals of Arid Zone* 42: 213–230.
- Williams, M.A. (2003b). Changing land use and environmental fluctuations in the African savanna. In T. J. Bassett & D. Crummey (eds), *African Savannas: Global Narratives and Local Knowledge of Environmental Change*. James Currey, Oxford, pp. 31–52.
- Williams, M.A. (2009). Human Impact on the Nile Basin: Past, Present, Future In H. J. Dumont (ed.), *The Nile. Monographiae Biologicae*, Vol. 89: 771–779. Springer, Dordrecht. [A multi-author monographs book, in which thirty-seven authors have taken up the challenge, and have written chapters cover the origin, environments, limnology and human use of the River Nile]

Witte, F., Msuku B.S., Wanink J.H., Seehausen O., Katunzi E.F.B., Goudswaard P.C and Goldschmidt T. (2000). Recovery of cichlid species in Lake Victoria: an examination of factors leading to differential extinction. *Reviews in Fish Biology and Fisheries* 10: 233–241.

Witte, F., Oijen, M.J.P., and Van, Sibbing, F.A. (2009). Fish Fauna of the Nile. In H. J. Dumont (ed.), *The Nile. Monographiae Biologicae*, Vol. 89: 647–675. Springer, Dordrecht. [A multi-author monographs book, in which thirty-seven authors have taken up the challenge, and have written chapters cover the origin, environments, limnology and human use of the River Nile]

Witte, F., Goldschmidt T., Wanink J., van Oijen M., Goudswaard K., Witte-Maas E. and Bouton N. (1992). The destruction of an endemic species flock: quantitative data on the decline of the haplochromine cichlids of Lake Victoria. *Environmental Biology of Fishes* 34: 1–28.

Wondie, A., Mengistu S., Vijverberg J. and Dejen E. (2007). Seasonal variation in primary production of a large high altitude tropical lake (Lake Tana, Ethiopia): effects of nutrient availability and water transparency. *Aquatic Ecology* 41: 195–207.

Woodward, J.C., Macklin, M. G., Krom M.D., and Williams, M.A.J. (2007). The Nile: Evolution, Quaternary river environments and material fluxes. In A. Gupta (ed), *Large Rivers: Geomorphology and Management*. Wiley, New York, pp. 261–292.

Yin, X. G. and Nicholson, S. E. (2000). On the diurnal cycle of cloudiness over Lake Victoria and its influence on evaporation from the lake. *Hydrological Sciences Journal* 45: 407–424.

Zaghloul, F.A. (1985). Seasonal variation of plankton in Lake Nasser. Ph.D. Thesis, Fac. Science, Suez Canal Univ. Egypt. 364 pp

### Biographical Sketches

**Gamal M. El-Shabrawy**, who was born on 24 February 1964. He obtained his PhD degree from Faculty of Science, Mansoura University, Egypt in 1996. Working as research assistant, researcher and team leader in many research projects that have been carried out in the Egyptian lakes, wetland ecosystems since 1993 up till now. Contributed in producing 4 and 2 chapters in 2 international reference books published by Springer and Nova Publication and one chapter in local reference book for the National Biodiversity Unit of EEAA, Lake Bardawil (2005). Teaching post graduate lectures on aquatic ecology, limnology and lake management. Collaborating and consultant in the EIA studies of many tourist Villages and Petroleum Companies. Attending many training courses and scholarships in Egypt, Jordan and Belgium as well as several national and international symposia, conferences and congresses in Egypt and abroad. Supervisor of 14 M.Sc.'s and Ph.D.'s in the fields of population and community of zooplankton and Macrobenthos. Member of many national and international councils, committees and societies in the field of limnology and Aquatic Environmental Sciences. Consultant of many international and local journals. Forty two publications, in national and international specialized journals, covering many aspects of aquatic ecology such as: Long term changes, Spatial and seasonal variation of the zooplankton and macrobenthos in all Egyptian lakes, Fish Farms and River Nile.

**Goher M.E.** received B.Sc degree. in Biochemistry\Chemistry from Faculty of Science, Ain Shams Univ. Egypt in 1991, M.Sc. degree in Inorganic Chemistry from El-Menofyia Univ. Egypt, and Ph.D. in Inorganic and analytical chemistry from Al-Azhar Univ. Egypt in 2002. In 2010, he received an associated professor degree in chemistry of oceanography from NIOF Egypt. His interests include maintain and develop the aquatic environment especially fresh water streams and how to get rid the causes of water pollution. He has ten publications, in national and international specialized journals.