

## SYMBIOTIC BACTERIA AND FUNGI

**E.K. James and M. Fomina**

*College of Life Sciences, University of Dundee, Dundee DD1 5EH, UK*

**S.M. de Faria**

*EMBRAPA-Agrobiologia, km 47, Seropédica, 23851-970, RJ, Brazil*

**Keywords:** Nitrogen fixation, legumes, Actinorhizal plants, nodulation, *Rhizobium*, *Burkholderia*, *Frankia*, ecto-mycorrhizas (EcM), arbuscular mycorrhizas (AMs), phosphorus, bioremediation

### Contents

1. Introduction
  2. Nitrogen-fixing Legumes and their Symbionts
  3. Infection of Legumes by Rhizobia
  4. Other Nitrogen-Fixing Organisms
  5. The Role of Nitrogen Fixation in the Nitrogen Cycle
  6. Mycorrhizal Fungi and the Phosphorus Cycle
  7. Use of Symbiotic Fungi and Bacteria in Bioremediation
  8. Conclusion
- Glossary  
Bibliography  
Bibliographical Sketches

### Summary


Nitrogen (N) and phosphorus (P) are the two most important nutrients for plant growth, but in many environments both are in short supply and/or are unavailable to plants. This is particularly true in tropical forests, where high temperatures and humidity cause mineralisation of organic N, and the very high precipitation results in rapid leaching of the remaining soil N. Therefore, the fixation of atmospheric dinitrogen into ammonia and hence into organic compounds is a crucial part of the N- cycle of these forests. Nitrogenase, the enzyme responsible for biological nitrogen fixation (BNF), has a wide taxonomic distribution in the bacteria and archaea, but is confined to a limited number of species. Among these, soil bacteria called rhizobia, which form symbiotic nodules on the roots of legumes (Fabaceae), are recognised as the main contributors of biologically-fixed nitrogen to pristine environments, most particularly in tropical forests where they are so abundant and diverse, but there are other symbiotic, endophytic, associative and free-living diazotrophs that make a significant contribution, and this chapter describes these as well as the leguminous symbioses.

Similarly, the availability of the second most major essential nutrient for plants, P, is greatly limited by its occurrence and low availability in the soil, most particularly owing to the low solubility of phosphates, and hence plants very early in their evolutionary history developed a symbiotic relationship with fungi called mycorrhizas. There are two

types of mycorrhizas: ecto- and endo- (e.g. arbuscular and ericoid) mycorrhizas, and most plant families (> 95 %) are capable of forming a symbiosis with at least one type. In this chapter we review current knowledge about symbiotic mycorrhizal associations within tropical forests and assess their likely contributions to the P-cycle.

## 1. Introduction.

Nitrogen and phosphorous are the two main limiting nutrients in both natural and agricultural systems. In agriculture both nutrients are supplied in forms available to plants as fertilisers produced either through chemical processes (eg. the Haber-Bosch process that converts or “fixes” atmospheric nitrogen into ammonia using very high quantities of energy) or from mining mineral deposits (eg. from rock phosphate). Regardless of whether the fertilisers are produced chemically or by mining, the energetic and environmental costs of their production are prodigious. However, in most undisturbed non-agricultural ecosystems (as well as some agricultural ones) soluble nitrogen (ammonium) is made available to plants via a process known as biological nitrogen fixation (BNF), in which bacteria that contain the enzyme complex called nitrogenase (termed “diazotrophs”) can fix atmospheric N<sub>2</sub> into ammonia using energy from ATP and reductant (electrons) supplied by respiration (which can be aerobic or anaerobic):



This ammonia, which is potentially toxic to the organism, is usually then immediately converted into amino acids or amides for use by the diazotrophic bacterium in the production of proteins and peptides to facilitate its growth. The fixed N incorporated into diazotrophic bacteria is then released into the environment when they die, usually in the form of amino acids that then become mineralized and available for uptake by other bacteria and/or by plants. Interestingly, although nitrogenase, the enzyme responsible for nitrogen fixation, has a wide taxonomic distribution in the bacteria and archaea, it is actually confined to a relatively limited number of species (J.P.W. Young in PALACIOS & NEWTON, 2005). Nevertheless, many plants, particularly legumes, but also some other higher plants (eg. *Gunnera*, and “actinorhizal” plants, such as *Alnus*, *Casuarina* and *Myrica*), as well as some cycads and the fern *Azolla*, form mutualistic symbiotic relationships with nitrogen-fixing bacteria. In these systems, soil bacteria located in specialized organs take nitrogen from the atmosphere and convert it to soluble nitrogen that is available for plant growth. Nitrogen-fixing legumes are one of the main sources of nitrogen in both natural environments and many agricultural systems. However, although pathways controlling nitrogen-fixation and the bacterial symbionts involved are well understood in a handful of model and crop legume species, very little is known about the diversity of symbiotic systems in natural environments in terms of the bacteria involved and their host relationships. This is especially so in the tropics where most legume diversity is found.

-  
-  
-

TO ACCESS ALL THE 25 PAGES OF THIS CHAPTER,  
Visit: <http://www.eolss.net/Eolss-sampleAllChapter.aspx>

### Bibliography

Allen, O.N., and Allen, E.K. (1981) *The Leguminosae. A Source Book of Characteristics, Uses and Nodulation*. The University of Wisconsin Press, Madison Wisconsin. 812 p. [Seminal book on nodulation in legumes, and still highly relevant].

Cardoso, I.M., Kuyper, T.W. (2006). Mycorrhizas and tropical soil fertility. *Agriculture, Ecosystems and Environment* 116: 72-84. [Up to date review on the current state of mycorrhizal research in the tropics].

Dilworth, M.J., James, E.K., Sprent, J.I., and Newton, W.E. (eds). (2008) *Nitrogen-fixing legume symbioses*. Volume 7 of *Nitrogen Fixation: Origins, Applications and Research Progress*. Springer, Dordrecht. 402 p. [Arguably the most comprehensive set of text books currently available on nitrogen fixation comprising 7 volumes covering all aspects of the subject in great depth].

Gadd, G.M. (ed.). (2006) *Fungi in Biogeochemical Cycles*. Cambridge University Press, Cambridge, p. 236-266. [Fundamental text book covering the roles that fungi play in altering their environments and how this impacts upon the larger scale in biogeochemical terms].

Loureiro, M.F., James, E.K., and Franco, A.A. (1998). Nitrogen fixation by legumes in flooded regions. *Oecologia Brasiliensis* 4: 191-219. [A review that covers most aspects of legume N fixation research in the flooded regions of the tropics].

Palacios, R., and Newton, W.E. (2005). *Genomes and Genomics of Nitrogen-fixing Organisms*. Volume 3 of *Nitrogen Fixation: Origins, Applications and Research Progress*. Springer, Dordrecht. 246 p. [Arguably the most comprehensive set of text books currently available on nitrogen fixation comprising 7 volumes covering all aspects of the subject in great depth].

Pawlowski, K., and Newton, W.E. (2008). *Nitrogen-fixing Actinorhizal Symbioses*. Volume 6 of *Nitrogen Fixation: Origins, Applications and Research Progress*. Springer, Dordrecht. 310 p. [Arguably the most comprehensive set of text books currently available on nitrogen fixation comprising 7 volumes covering all aspects of the subject in great depth].

Pott, V.J., and Pott, A. (2000). *Plantas aquáticas do Pantanal*. Embrapa, Brasília. 400 p. [Beautifully illustrated book describing the aquatic plants that live in the largest wetland in the World].

Robson, G.D., Van West, P. and Gadd, G.M. (eds.). (2007). *Exploitation of Fungi*. Cambridge University Press, Cambridge, p. 236-254. [Several review papers covering all current aspects of research into beneficial fungi, including mycorrhizas].

Smith, S.E., and Read, D.J. (1997) *Mycorrhizal Symbiosis*. Academic Press, London, UK. [Excellent fundamental text book on mycorrhizas].

Sprent, J.I. (2001). *Nodulation in legumes*. London, UK: Royal Botanic Gardens, Kew. [The most important single volume of information about nodulation in legumes currently available. Comprehensive and easy to read].

Sprent, J.I., and James, E.K. (2007) Legume evolution: Where do nodules and mycorrhizas fit in? *Plant Physiology* **144**, 575-581. [Recent discussion paper examining the roles of the mycorrhizal and rhizobial symbioses in the evolution of legumes].

Werner D., and Newton, W.E. (eds). (2005). *Nitrogen fixation in agriculture, forestry, ecology, and the environment*. Volume 4 of *Nitrogen Fixation: Origins, Applications and Research Progress*. Springer,

Dordrecht. 347 p. [Arguably the most comprehensive set of text books currently available on nitrogen fixation comprising 7 volumes covering all aspects of the subject in great depth].

### **Biographical Sketches**

**Euan K. James** is a researcher at the University of Dundee and a consultant for the Scottish Crop Research Institute. His present research interests are focused on nodulation of legumes, particularly those from seasonally-dry tropical savannahs and forests.

**Sérgio Miana de Faria** is a researcher at Embrapa-Agrobiologia, km47, Seropédica, Rio de Janeiro, Brazil. His present research interests are plant-microbe interactions with an emphasis on the use of symbioses in the rehabilitation of degraded land.

**Marina Fomina** is a researcher at the University of Dundee. Her current research interests are the role of fungal communities and mycorrhizas in geochemical transformations of metals and minerals.