

## CLASSIFICATION OF WATER-RELATED DISEASE

### R Stanwell-Smith

*Royal Institute of Public Health, UK (formerly Consultant Epidemiologist, Water related disease surveillance, Communicable Disease Surveillance Centre, Public Health Laboratory Service, England, UK)*

**Keywords:** Water-related disease, water-borne disease, infectious disease, climate change, swimming pools, floods, epidemiology, surveillance, outbreaks

### Contents

1. Introduction
  2. Definitions and Systems for Classifying Water Related Disease
    - 2.1. Essential Components for the Classification
    - 2.2. Definition of Water-related Disease
    - 2.3. Categories of Water-related Disease
    - 2.4. Type of water exposure
    - 2.5. Cause and evidence: important issues in classifying water-related disease
    - 2.6. Burden of Proof in the Case of Water-related Disease
  3. Water-related infection
  4. Non-infectious Disease Related to Water
  5. Water Associated Disease
  6. Implications for water related surveillance
- Glossary  
Bibliography  
Biographical Sketch

### Summary

Water related disease encompasses illness resulting from both direct and indirect exposure to water, whether by consumption or by skin exposure during bathing or recreational water use. It includes disease due to waterborne or water-associated pathogens and toxic substances. A broader definition includes illness related to water shortage or water contamination during adverse climate events, such as floods and droughts, diseases related to vectors with part of their life cycle in water habitats; and disease related to inhalation of contaminated water aerosols. Water-related disease may be classified in terms of the agents responsible and the types of water exposure, but for surveillance purposes, the classification system must also include the probability of association with water, since multifactorial aetiology is involved in most, if not all, of the water-related diseases. In public health terms, the classification system should also include factors related to intervention, such as the type of water treatment or sanitation measures required to remove the disease-causing factors. Host factors are relevant to water related disease transmission, since many of the agents have more impact on individuals who are malnourished or already suffering from other disease, including diseases associated with immune deficiency. The inherent relationship between water and food also complicates water related disease classification: in many surveillance systems, statistics on water related diseases are subsumed within data on gastroenteritis

or on food poisoning.

## 1. Introduction

(i)The classification of water-related disease poses particular problems. Agents, diseases and types of exposures are well documented, but the relationship between them is subject to degrees of certainty, timing of exposure and the nature of the exposure. Water in one form or another is a universal exposure for human beings: water is a fundamental requirement for nutrition and overall health, with a corresponding wide range of diseases that can be classified as having a water component. All the agents involved have other means of transmission and the evidence that confirms a water association can be elusive, often depending on whether a large outbreak or documented incident of contamination has been identified. While it has been estimated that the worldwide burden of water-related disease includes millions of cases of illness and deaths, surveillance systems for detecting and confirming water related disease vary considerably between countries. In some countries, only severe and wide-scale outbreaks are recorded; in most, minor episodes of water-related disease are either not investigated or not recorded on national databases.

(ii)Historically, water related disease has been a major health problem for both developed and developing countries. Its priority as a surveillance issue has fallen partly as a result of improved water and sanitation in industrialized and northern countries of the world, associated with lower morbidity and mortality from diseases such as cholera and typhoid fever. Food related disease surveillance has overtaken it in importance and in some countries water-related disease is classified under food poisoning or gastrointestinal infection, further diminishing its importance in surveillance statistics. The emphasis on the infections affecting the gastrointestinal system is appropriate in terms of numbers of cases, since gastrointestinal infection is the commonest water-related illness, worldwide: but a gastrointestinal-based classification ignores respiratory infections linked to contaminated water, such as legionellosis, and also skin or systemic disorders. Non-infectious water-borne disease is increasingly recognized as a public health concern, requiring different types and complexity of surveillance from the traditional systems based on microbiological agents.

(iii)While surveillance of illness attributable to treated water is a particular concern in developed or industrialized countries, only a minority of the world's population receives continuous piped supplies. According to figures from the WHO and UNICEF in 2000, 2.4 billion people do not have acceptable means of sanitation and 1.1 billion are estimated to lack an improved water supply. For the 'water poor' populations of the world, the risks associated with untreated water or poor sanitation include both individual risk of illness and public health issues of monitoring, surveillance and prevention. Increased population mobility and travel to different countries has to some extent globalized the risk of water-related disease, and the corresponding requirement for national surveillance systems to include the wide range of potential illnesses. Water-related disease surveillance is likely to rise in priority with the impact of climate change, including the effects of global warming on water habitats and adverse weather events threatening the security of water supplies. Other threats to safe water include the effect of the world economy on the funds available for public health measures and the

possibility of deliberate release of pathogens or toxic agents into water intended for human consumption.

(iv) This chapter includes both a basic and an extended system for classifying water-related disease, as well as discussing how attribution of cause, in terms of levels of likelihood, can be incorporated into the classification.

## 2. Definitions and Systems for Classifying Water Related Disease

### 2.1. Essential Components for the Classification

There are three essential components for the classification: the pathogens and other agents involved in water-related disease; the type of water exposure; and the level of probability of a water cause. Host factors, such as nutritional status, are important in terms of the priority and detail required for surveillance systems in countries with high levels of malnutrition, immune deficiency, or significant mortality from water borne pathogens (see Figure 1)




Figure 1. Relationships involved in the classification of water-related disease

## **2.2. Definition of Water-related Disease**

Of the various terms for disease linked to water, water related disease is the most comprehensive. Water-related disease is defined as any significant or widespread adverse effects on human health, such as death, disability, illness or disorders, caused directly or indirectly by the condition, or changes in the quantity or quality of any waters.

The causes of water related disease include micro-organisms, parasites, toxins and chemical contamination of water. Other terms include ‘waterborne disease’, which implies direct spread and is used mainly to refer to disease caused by microbiological pathogens or chemical contaminants in water.

‘Water associated disease’ covers the wide range of diseases in which water plays a part, such as legionnaires’ disease, as well as diseases related to lack of water for washing and hygiene.

The advantage of the term water related disease is that it includes both water borne and water associated ill health, although diseases with an indirect association and another major mode of spread are usually excluded from specific surveillance systems.

An example of an indirectly related disease is trachoma: the predominant mode of spread is via poor hygiene and flies thriving in conditions of poor sanitation: clean water for hygiene is an important element in prevention, but the disease is not otherwise water-related.

Such diseases used to be known as ‘water-washed’, referring to the role of clean water in removing the agents, but the term is no longer widely used: McJunkin suggested the alternative of ‘water-hygiene diseases’.

## **2.3. Categories of Water-related Disease**

Seven categories of water-related disease can thus be identified (Table 1) waterborne microbiological disease; waterborne chemical disease; water hygiene disease; water contact disease; water vector habitat disease; excretal disposal disease and water aerosol disease. In addition to the subcategories in Table 1, other subcategories could be included such as those related to duration of exposure (acute/ prolonged).

## **2.4. Type of water exposure**

The commonest distinction regarding water exposure is between drinking water and recreational water. For some surveillance systems, disease is also classified according to whether the water supply involved is a public water system, usually regulated by national legislature and local by-laws, or other types of supply, such as private wells or untreated water sources. A more detailed classification requires specification of the type of water source, such as groundwater or surface water. For water-contact diseases, the classification sub-categories include fresh water and marine waters.

Category	Description of category	Type of water exposure	Subcategories	Example(s)
<b>Waterborne microbiological disease</b>	Diseases related to consumption of pathogens consumed in water; most due to human or animal faecal contamination of water	Drinking water	(i) Treated or untreated (raw) water (ii) Public (municipal) supplies or private supplies	Cholera, Typhoid fever, viral gastroenteritis e.g. due to Norovirus
<b>Waterborne chemical disease</b>	Disease related to ingestion of toxic substances in water	Drinking water	(i) Treated or untreated (raw) water (ii) Public (municipal) supplies or private supplies	Arsenicosis
<b>Water hygiene diseases</b>	Diseases whose incidence, prevalence or severity can be reduced by using safe (clean) water to improve personal and domestic hygiene	Any water used for washing/ personal hygiene	(i) Disease related to variations in water quality (ii) Disease related to water shortage	Scabies, shigellosis; trachoma
<b>Water contact diseases</b>	Caused by skin contact with pathogen-infested water or with chemical-contaminated water	Recreational water	(i) fresh water sources (ii) marine waters	Schistosomiasis (bilharzia); cyanobacteria
<b>Water vector habitat diseases</b>	Diseases where vector lives all or part of its life in or adjacent to a water habitat	Untreated freshwater sources	(i) rivers, streams (ii) small collections of stagnant water e.g. water butts	Malaria (mosquitoes); filariasis (mosquitoes); onchocerciasis (aquatic flies); schistosomiasis (snails); trypanosomiasis (tsetse flies)
<b>Excreta disposal diseases</b>	Diseases related to unsanitary disposal of human waste (faeces and urine)	Drinking water and untreated water sources	(i) diseases related to human/animal waste in drinking water (ii) diseases related to direct/ indirect contact with faeces/ urine	Ascariasis; faecal-oral infections e.g. shigellosis; schistosomiasis; trachoma
<b>Water aerosol diseases</b>	Diseases related to respiratory transmission, where a water aerosol containing suspended pathogens enters airway	Drinking or raw water sources	(i) water used in industrial/ residential buildings (ii) raw water sources	Legionellosis (legionnaires' disease; humidifier fever); Norwalk-like viral gastroenteritis

Table 1. Classification of water related disease

## 2.5. Cause and evidence: important issues in classifying water-related disease

For surveillance purposes, the definition of water related disease must include probability of a water cause as well as the agent responsible. Attributing cause to water is not straightforward, because of the factors and levels of probability to be considered. Most of the diseases related to water can also be transmitted in other ways and the evidence of water contamination is not always available by the time an outbreak of disease is identified. While epidemiological evidence, such as an analytical study showing a strong statistical association with water, is an important element in attributing cause, epidemiological data, without strong microbiological or chemical evidence, may not meet the requirements for legal evidence and may therefore be disputed by water providers. There are practical, political as well as epidemiological considerations involved in attributing cases of disease or an outbreak to water. These include the cost of arranging alternative water supplies at short notice and the damage to public confidence in water supplies when a water outbreak is suspected or announced. Hence surveillance systems usually include a system for classifying probability on the basis of the available evidence. For example, in the UK, outbreaks are classified as definite, probable or possible, depending on the quality of microbiological and epidemiological evidence (Table 2). A broadly similar system is used in the USA.

<b>1. UK Criteria</b>		
(a) the pathogen found in human case samples was also found in water samples;		
(b) documented water quality failure or treatment failure;		
(c) significant result from analytical epidemiological study (case control or cohort);		
(d) suggestive evidence of association from a descriptive epidemiological study, for the		
<b>Strength of Association</b>		
Strong	<b>(a)+(c), (a)+(d) or (b)+(c)</b>	
Probable	<b>(b)+(d), (c) only for (a)</b>	
Possible	<b>(b) + (d)</b>	
<b>2. USA (MMWR 1996)</b>		
<b>Class</b>	<b>Epidemiological data</b>	<b>Water quality data</b>
I Adequate	Adequate data provided in outbreak report about exposed vs unexposed persons; Relative risk or odds ratio $\geq 2$ or p-value $\leq 0.05$	Historical information or laboratory data e.g. history of malfunction of chlorine treatment; broken water main; no detectable free-chlorine residual; presence of coliforms in water
II Adequate	As above	Inadequate information
III Provided, but limited	Data provided but not meeting Class I criteria; no data provided to	As in class I.

	substantiate claim of no common exposures except water	
IV Provided, but limited	As above	Inadequate information

Table 2. Criteria for estimating strength of association between human illness and water

### 2.6. Burden of Proof in the Case of Water-related Disease

The need for evidence applies to all disease causation, but in the case of water-related disease the burden of proof has two important effects:

(i) Water related disease is under-estimated in most surveillance systems, since often only outbreaks with strong evidence are included, despite strong circumstantial evidence (such as descriptive epidemiological studies) for other incidents. The problem of attributing cause is even greater for small clusters and travel-related cases of disease, where the evidence may be impossible to collect. Under-estimation is related to the causative agent, the severity and duration of illness and the size of the outbreak: Under-estimation of the burden of water-related disease has a secondary effect on the priority given to surveillance systems and to preventive measures.

(ii) Attribution of cause to water is often delayed, while evidence is collected or until further cases occur. This can result in a much higher level of disease due to persistent or recurrent contamination. The mercury poisoning in Minamata, Japan, is a classic example of delayed attribution of a water-related cause. Delay in identifying a water cause is particularly likely to occur where the associated disease has a long incubation, for example in waterborne typhoid fever, or only occurs after prolonged accumulation in the body, as in water related arsenicosis. For some pathogens, the water exposure may be so far in the past that the link is missed for years, for example the possible water association only recently recognized for the emerging pathogen *Helicobacter pylori*.

-  
-  
-

TO ACCESS ALL THE 21 PAGES OF THIS CHAPTER,  
Visit: <http://www.eolss.net/Eolss-sampleAllChapter.aspx>

#### Bibliography

Altmann P, Cunningham J, Dhanesha U, Ballard M, Thompson J, Marsh F. (1999) Disturbance of cerebral function in people exposed to drinking water contaminated with aluminium sulphate: retrospective study of the Camelford water incident. *British Medical Journal* ; 319: 807-812

Abrahams MJ, Price J, Whitlock FA, Williams G The Brisbane Floods, January 1974: their impact on health. *Med J Aust* 1976; 2: 936-9

Andersson N. (1990) Disaster epidemiology: lessons from Bhopal. In: Major Chemical Disasters – medical aspects of management. Ed. V. Murray. London, UK: Royal Society of Medicine Services Ltd.

Anon. Public health consequences of a flood disaster – Iowa, (1993). *MMWR Morb Mortal Wkly Rep* 1993 (Sep 3); 42: 653-6

Axelrod C, Killam PP, Gaston MH, Stinson N. (1994) Primary health care and the Midwest flood disaster. *Public Health Rep* 1994 Sep-Oct; 109: 601-5

Bartram J, Rees G (eds). *Monitoring Bathing Waters: a practical guide to the design and implementation of assessments and monitoring programmes*. Published on behalf of the WHO. E& FN Spon, UK: 1999

Baxter P. Review of major chemical incidents and their medical management. In: Murray V (Ed) *Major Chemical Disasters- medical aspects of treatment* London: Royal Society of Medicine, 1990, pp 7-20;

Bedos C, Brodeur JM. Determinants of dental caries on Haitian school children and implications for public health (Déterminants de la carie dentaire parmi les écoliers haitiens et implications pour la santé publique). *Cahiers Santé* 2000; 10:161-8 (English summary).

Bennet G. Bristol floods 1968. Controlled survey of effects on health of local community disaster. *Br Med J* 1970; 22; 3(720): 454-8

Benton C, Forbes GI, Paterson GM, Sharp JCM, Wilson TS. The incidence of water-borne and water-associated disease in Scotland from 1945 to 1987. *Water Science and Technology* 1989; 21(3):125-129

Birley MH. *Guidelines for forecasting the vector-borne disease implications of water resources development*. Geneva: WHO, 1991

Bradford Hill A. The environment and disease: association or causation? *Proceedings of the Royal Society of Medicine*. 1965, 58: 295-300

Casemore D. Pseudo-outbreak of cryptosporidiosis. *Communicable Disease Report* 1992; R84-85

Checkley, W. et al., 2000. Effects of El Niño and ambient temperature on hospital admissions for diarrhoeal diseases in Peruvian children. *The Lancet*, vol. 355, no. 9202.

Chelala C *Environmental impact on child health*. Washington, DC: Pan American Health Organization (PAHO) and WHO, 2000 (ISBN 92 75 12265 2). See also [www.paho.org](http://www.paho.org)

Chinn J (ed). *Control of Communicable Diseases Manual*, 17<sup>th</sup> edition. American Public Health Association, 2000

Chorus I, Bartram J, eds. *Toxic cyanobacteria in water: a guide to their public health consequences, monitoring and management*. Published on behalf of WHO. London/New York: E&FN Spon (Routledge), 1998{?}

Codex Standard for Natural Mineral Waters and Code of Practice: <http://www.fao.org/WAICENT/FAOINO/ECONOMIC/ESN/codex>

Colwell RR. Global climate and infectious disease: the cholera paradigm. *Science*, 1996; 274: 2025-31

Communicable Disease Surveillance Centre. Strength of association between human illness and water: revised definitions for use in outbreak investigation. *Communicable Disease Weekly* 1996; 6: 65.

Communicable Disease Surveillance Centre. *CDR report* Jan 1999.

Cornwall and Isles of Scilly District Health Authority. *Water pollution at Lowermoor North Cornwall: Report of the Lowermoor incident health advisory group (Clayton Report)*. Cornwall and Isles of Scilly DHA, 1989.

Craun GF. *Waterborne disease outbreaks in the United States of America: causes and prevention*. *World Health Statistics Quarterly*, 1992; 45: 192-199

Department of the Environment and Department of Health. *Cryptosporidiosis in water supplies. Report of the group of experts (Badenoch report I)*. London: HMSO, 1990.

Department of the Environment and Department of Health. *Third Report from the Group of Experts on*



Cryptosporidium in Water Supplies (Bouchier report). London: HMSO, 1998.

Department of the Environment, Sport and Territories (Australia). Biodiversity and its value: biodiversity series no. 1. [http://www.erin.gov.au/life/general\\_info/op1.html](http://www.erin.gov.au/life/general_info/op1.html) Canberra, Australia: Department of EST, 1993

Dieter HH. The 'Red Rhine incident': a toxicological view. In: Golding AMB, Noah N, Stanwell-Smith R (eds) *Water and Public Health*. 1994, Smith-Gordon & Co: London pp 59-76

Eastcott HR. Swimmer's itch: a surfacing problem? *Communicable Disease Report* 1988; 88(12): 3-4.

Ecologically Sustainable Development (ESD) Working Group on Tourism. *Ecologically Sustainable Development Working Groups: Final Report –Tourism*. Canberra, Australia: AGPS, 1991

Engelman, R. (2000). *People in the balance: population and natural resources at the turn of the Millennium*. Washington, DC, USA, Population Action International. 36 p. <http://populationaction.org/balance/index.htm>

Evans J, Hyndman S, Stewart Brown S, Smith D, Petersen S. An epidemiological study of the relative importance of damp housing in relation to adult health. *J Epidemiol Community Health* 2000 Sep; 54(9): 677-86

Forss H. Efficiency of fluoride programs in the light of reduced caries levels in young populations. *Acta Odontol Scand* 1999; 57: 348-351

Furtado C, Adak GK, Stuart JM, Wall PG, Evans HS, Casemore DP. Outbreaks of waterborne infectious intestinal disease in England and Wales, 1992-5

Galbraith NS, Barrett NJ, Stanwell-Smith R. Water and disease after Croydon: a review of water-borne and water-associated disease in the UK 1937-86. *Journal of the Institute of Water and Environmental Management*, 1987; 1:7-21

Galbraith NS. A historical review of microbial disease spread by water in England and Wales. In: Golding AMB, Noah N, Stanwell-Smith R (eds) *Water and Public Health*. 1994, Smith-Gordon & Co: London pp 15-37

Githeko AK, Lindsay SW, Confalonieri UE, Patz JA. Climate change and vector-borne diseases: a regional analysis. *Bulletin of the World Health Organization*, 2000; 78(9)11361147

Gleick P H. *The World's Water 2000-2001: the biennial report on freshwater resources*. Washington DC USA: Island Press, 2000

Gwatkin DR and Guillot M, 2000. *The Burden of Disease among the Global Poor: Current Situation, Future Trends, and Implications for Strategy*. Washington, DC, The International Bank for Reconstruction and Development/The World Bank, p. 1-44.

Hrudey S. Drinking-water risk management principles for a total qualitative management framework. *J Toxicol Environ Health A*. 2004; 67(20-22):1555-66

Hrudey SE, Hrudey EJ. *Safe drinking water: lessons from recent outbreaks in affluent nations*. IWA Publishing, 2004 [ISBN 1 84339 042 6]

Hulme M, Jenkins G, Brooks N et al. What is happening to global climate and why? In: Department of Health, UK: *Health Effects of Climate Change in the UK. An Expert Review*. Department of Health, England, UK: 2002. Chapter 1.2 pp 21-25. ([www.doh.gov.uk](http://www.doh.gov.uk))

Hunter PR. *Waterborne Disease: epidemiology and ecology*. Chichester, England: John Wiley and Sons Ltd, 1997

Hunter PR. Drinking water and diarrhoeal disease due to *Escherichia coli*. *J Water Health* 2003; 1(2):65-72

Hunter P. *Waterborne disease: epidemiology and ecology*. Ch. 3. *Drinking Water and waterborne disease*. 1997, Chichester: John Wiley & Sons; pp 27-34.

Hunter P. *Waterborne disease: epidemiology and ecology*. Ch 11 *Cyanobacteria*. In: *Drinking Water and waterborne disease*. 1997, Chichester: John Wiley & Sons; pp 95-102.

Hunter P. Waterborne disease: epidemiology and ecology. Ch 15 *Campylobacteriosis* and Ch. 26 *Helicobacter infections*. In: *Drinking Water and waterborne disease*. 1997, Chichester: John Wiley & Sons; pp 132-142 and pp 202-205.

IPCS. Status reports on projects on endocrine disruptors: <http://endocrine.ei.jrc.it/>;

IPCS (International Programme on Chemical Safety) Fact Sheet no.10: Endocrine disruptors, May 1998 (web site <http://www.cho.int/pcs> [http://www.who.int/pcs/emerging\\_issues/end\\_disrupt.htm](http://www.who.int/pcs/emerging_issues/end_disrupt.htm))

Jobin WR. *Dams and disease*. London: Routledge, 1999

Jones PD, New M, Parker DE et al. Surface air temperature and its changes over the past 150 years. *Rev Geophys* 1999; 37: 173-199

Knobeloch L, Salna B, Hogan A, Postle J, Anderson H. Blue babies and nitrate-contaminated well water. *Environ Health Perspect*. 2000 Jul;108(7):675-8.

Kovats RS. El Nino and human health. *Bulletin of the World Health Organization*, 2000; 78(9):1127-1135

Kramer MD, Herwaldt BL, Craun GF, Calderon RL, Juranek DD. Surveillance for waterborne-disease outbreaks - United States, 1993-1994. *Morbidity and Mortality Weekly Report* 1996, 45 (SS-1):1-33

Lachance KR, Santos AB, Burns BJ. The response of an assertive community treatment program following a natural disaster. *Community Ment Health J* 1994; 30: 505-15

Lanz K 1995. *Watering the Land*. In: *The Greenpeace Book of Water*. Scotland: David & Charles, 1995. Chapter 7, pp 78-101

MacKenzie WR, Hoxie NJ, Proctor ME, Gradus MS, Blair KA, Peterson DE, Kazmierczak JJ, Addiss DG, Fox KR, Rose JB, Davis JP. A massive outbreak in Milwaukee of cryptosporidium infection transmitted through the public water supply. *New England Journal of Medicine*, 1994; 331: 161-167

McEvoy MB, Noah ND, Pilsworth R. Outbreak of fever caused by *Streptobacillus moniliformis*. *Lancet* 1987; 2:1361-1363

McEvoy M, Batchelor N, Hamilton g, MacDonald A, Faiers M, Sills A, Lee J, Harrison T. *Communicable Disease and Public Health* 2000; 3(1):43-5

McJunkin F E. *Water and human health*. National Demonstration Water Project for US Agency for International Development, USA, 1982 (AID/DSAN-C-0063)

McMichael AJ, Haines A, Slooff R, Kovats S (eds). *Climate Change and Human Health: an assessment prepared by a Task Group on behalf of the World Health Organization, World Meteorological Organization and the United Nations Environment Programme*. Geneva: WHO, 1996

Milgrom P, Riedy CA, Weinstein P, Tanner ACR, Manibausan L, Bruss J. Dental caries and its relationship to bacterial infection, hypoplasia, diet, and oral hygiene in 6- to 36-month old children. *Community Dentistry and Oral Epidemiology* 2000; 28: 295-306

Moore AC, Herwaldt BL, Craun GF, Calderon RL, Highsmith AK, Juranek DD Surveillance for waterborne disease outbreaks - United States, 1991-1992. *Morbidity and Mortality Weekly Report* 1993, 42 (SS-5):1-22.

Mulloy KB. Sewage workers: toxic hazards and health effects. *Occup Med* 2001 Jan; 16(1): 23-28

Nazareth B, Stanwell-Smith R, Rowland MGM, O'Mahony M C. Surveillance of waterborne disease in England and Wales. *Communicable Disease Review* 1994; 4: R93-95

Noji E. *The public health consequences of disasters*. Oxford: Oxford University Press, 1997

O'Donnell M, Platt C, Aston R Effect of a boil water notice on behaviour in the management of a water contamination incident. *Commun Dis Public Health* 2000 Mar; 3(1): 56-9

Ohl C A, Tapsell S. Flooding and human health. Editorial. *British Medical Journal* 2000; 321:1167-9

Okada M, Peterson SA. *Water Pollution Control Policy and Management: the Japanese Experience*. Japan: Gyosei, 2000. (ISBN4-324-06240-4)

Paneth N, Vinten-Johansen P, Brody H, Rip M. A rivalry of foulness: official and unofficial investigations of the London Cholera Epidemic of 1854. *American Journal of Public Health*, 1998; 88(10): 1545-1553

Parshionikar S U, Willian-True S, Shay Fout G, Robbins DE, Seys SA, Cassady JD, Harris R. Waterborne outbreak of gastroenteritis associated with a Norovirus. *Applied and Environmental Microbiology* 2003; 69(9): 5263-5268

Payment P, Franco E, Richardson L, Siemiatycki J. Gastrointestinal health effects associated with the consumption of drinking water produced by point-of-use domestic reverse-osmosis filtration units. *Applied Environmental Microbiology*, 1991; 57:945-948

Payment P, Richardson L, Siemiatycki J, Dewar R, Edwards M, Franco e. A randomised trial to evaluate the risk of gastrointestinal disease due to consumption of drinking water meeting currently accepted microbiological standards. *American Journal of Public Health*, 1991; 81: 703-708

Pool water treatment advisory group. *Swimming pool water treatment and quality standards*. LREO Books Ltd, Holly Lodge, Botesdale, Diss, Norfolk IP22, UK, 1999

Postel S. *Pillar of Sand: can the irrigation miracle last?* New York/London: WW Norton & Co, 1999

Public Health Laboratory Service. *Hygiene for hydrotherapy pools*. London, UK: PHLS, 1999

Public Health Laboratory Service. *Hygiene for spa pools*. London, UK: PHLS, 1994

Reid WV, Miller K R. *Keeping options alive: the scientific basis for biological diversity*. Washington DC, USA: World Resources Institute, 1989

Richardson A J, Frankenburg R A, Buck AC, Selkon JB, Colbourne JS, Parson JW, Mayon-White RT. An outbreak of waterborne cryptosporidiosis in Swindon and Oxfordshire. *Epidemiology and Infection*, 1991; 107:485-495

Riley J C, Lennon MA, Ellwood R P. The effect of water fluoridation and social inequalities on dental caries in 5-year old children. *International Journal of Epidemiology*, 1999; 28: 300-305.

Rooney RM, Bartram JK, Cramer EH, Mantha S, Nichols G, Suraj R, Todd EC. A review of outbreaks of waterborne disease associated with evidence for risk management. *Public Health Rep*. 2004; 119(4):435-42

Rowland A, Grainger R, Stanwell-Smith R, Hicks N, Hughes A. Water contamination in North Cornwall: a retrospective cohort study into the acute and short term effects of the aluminium sulphate incident in July 1988. *Journal of the Royal Society of Health* 1990; 166-172

Shearman D. Biodiversity and ecology are health issues. [http://www.nccnsw.org.au/member/cbn/projects/scientists\\_say/ss\\_shear.html](http://www.nccnsw.org.au/member/cbn/projects/scientists_say/ss_shear.html) (Community Diversity Network: <http://www.cbn.org.au>)

Smith AH. Contamination of drinking-water by arsenic in Bangladesh: a public health emergency. . *Bulletin of the World Health Organization*, 2000; 78(9)1093-1103

Smith W.E, Smith A M. *Minamata*. New York: Holt, Rinehart and Winston, 1975

Snow J. *On the mode of communication of cholera*, 2nd edition, much revised (London: Churchill, 1855; reprint, with an introduction by W H Frost, New York: Commonwealth Fund 1936).

Somerville N, Mackenzie I, Owen P, Miles D. Housing and health: does installing heating in their homes improve the health of children with asthma? *Public Health* 2000 Nov; 114 (6):434-9

Stanwell-Smith R, O'Brien S. Cholera in tourists returning from Kenya. *Eurobulletin* 1998;2:0416 ( <http://www.eurosurv.org>)

Stanwell-Smith R. In the steps of `King Cholera': from Mevagissey to Kenya. *Health and Hygiene*, 1999; 20: 1-4

Stanwell-Smith R. Water and public health in the United Kingdom: recent trends in the epidemiology of water-borne disease. In: Golding AMB, Noah N, Stanwell-Smith R (eds) *Water and Public Health*. 1994, Smith-Gordon & Co: London pp 39-57

Stanwell-Smith R. Public Health and waterborne diseases. *Public Health Medicine*, 1999; 2: 53-60

Stanwell-Smith R. Waterborne diseases and climate change. In: Department of Health. *Health effects of climate change in the UK*. London: Department of Health, 2002, pp120-132.

Stanwell-Smith R. National Surveillance systems. In: *Drinking Water and Infectious Disease: establishing the links* (Eds P R Hunter, M Waite, E Ronchi). CRC Press 2002 (ISBN 0-8493-1259-0)

Strachan DP Damp housing and childhood asthma: validation of respiratory reporting of symptoms. *BMJ* 1988; 297: 1223-6

United Nations Population Division, 2000. *World Population Nearing 6 Billion Projected Close to 9 Billion by 2050*. New York, United Nations Population Division, Department of Economic and Social Affairs (Internet communication of 21 September 2000 at web site <http://www.popin.org/pop1998/1.htm>).

United Nations Population Fund (UNFPA), 1997. *Population and sustainable development – Five years after Rio*. New York, UNFPA, p. 1-36.

Wechsberg J. *The lost world of the great spas*. New York: Harper & Row, 1979

Wheeler JG, Sethi D, Cowden JM, Wall PG, Rodrigues LC, Tompkins DS, Hudson MJ,

Roderick PJ. Study of infectious intestinal disease in England: rates in the community, presenting to general practice, and reported to national surveillance. *British Medical Journal* 1999; 318: 1046-1050

WHO (b). Disinfectants and disinfectant by-products. *Environmental Health Criteria* 216, WHO: Geneva, 2000

WHO (IPCS): *Hazardous chemicals in human and environmental health: a resource book for school, college and university students*. Geneva: WHO (IPCS), 2000

WHO 1999. *WHO Report on Infectious Diseases – Removing obstacles to healthy development*. Geneva, World Health Organization.

WHO 2000. *WHO Report on Global Surveillance of Epidemic-prone infectious diseases*. Department of Communicable Disease Surveillance and Response, WHO, Geneva, 2000.

WHO Fact Sheet No. 256. *Bottled Drinking Water*. Geneva: WHO, October 2000. <http://www.who.int/inf-fs/en/fact256.html>

WHO, 2000. *The world health report 2000 - Health systems: Improving performance*. Geneva, World Health Organization, 2000.

WHO. (Annapolis Protocol). *Health based monitoring of recreational waters: the feasibility of a new approach*. Geneva: WHO/SDE/WSH99.1, 1999

WHO. *Cholera in 1997*. *Weekly Epidemiological Record*, 1998; 73: 201-8

WHO. *El Niño and its health impacts*. *Weekly Epidemiological Record*, 1998; 73: 148-152

WHO. *Fluoride in drinking water*. *Water, Sanitation and Health Series/ draft/99.4*. Geneva: WHO, 1999

WHO. *Fluorides and oral health*. *WHO Technical Report Series* 846. Geneva: WHO, 1994

WHO. *Food safety and foodborne illness WHO fact sheet no 237*, March 2000. Geneva: WHO, 2000

WHO. *Guidelines for drinking water quality, 3rd edition. Volume 1: Recommendations. Volume 2: Health criteria and other supporting information*. Geneva: WHO, 2004

WHO. *Guidelines for safe recreational water environments. Vol 1 Recreational and bathing waters*. WHO: Geneva, 2003

WHO. *Guidelines for safe recreational water environments. Vol 2 Pools and Spas* WHO: Geneva, in preparation 2004

WHO. *Oral health surveys: basic methods. 4<sup>th</sup> edition*. Geneva: WHO

WHO. *Prevention of oral diseases*. WHO Offset Publication, no.103. Geneva: WHO, 1987

WHO. *Recent advances in oral health: report of a WHO Expert Committee*. WHO Technical Report Series, no. 826. Geneva: WHO, 1992

WHO. *Turning the tide of malnutrition: responding to the challenge of the 21<sup>st</sup> century*. *Nutrition of Health and Development, Sustainable Development and Healthy Environments*. Geneva: WHO, 2000

WHO. Human health and dams: The World Health Organization's submission to the World Commission on Dams. WHO: Sustainable Development and Healthy Environments, Geneva January 2000 (WHO/SDE/WSH/00 01)

WHO: El Niño and Health. Task Force on Climate and Health. Geneva: WHO, 1999. WHO/SDE/99.4

WHO: Fact sheet on Rift Valley Fever [www.who.int/inf-fs/en/fact207.html](http://www.who.int/inf-fs/en/fact207.html)

WHO: Food safety: an essential public health issue for the New Millennium. Geneva: WHO, 1999

WHO: Leptospirosis worldwide 1999. Weekly Epidemiological Record 1999; 74: 237-242

WHO: Leptospirosis, India. Weekly Epidemiological Record, 2000; 75: 217-223

WHO: Public Health impact of pesticides used in agriculture. Geneva: World Health Organization, 1990 (ISBN 92 4 156139 4); International Programme on Chemical Safety Fact sheet no 16: Epidemiology of pesticide poisoning (web site <http://www.who.int/pcs>).

WHO: The control of schistosomiasis. Second report of the WHO Expert Committee. WHO Technical Report Series 830. Geneva: WHO, 1993

WHO. Poor environmental quality, exposures and risks. In: Health and environment in sustainable development: five years after the Earth Summit. Geneva: World Health Organization, 1997. Chapter 4, 4.4 pp 96-104

WHO: Hazardous chemicals in human and environmental health: a resource book for school, college and university students. Geneva: WHO (IPCS), 2000 p10

WHO: Health consequences of the Chernobyl accident: results of IPHECA pilot projects and related national programmes. Summary report. Geneva: WHO, 1995; WHO. Nuclear accidents: harmonization of the public health response. Report on a WHO meeting. EURO Reports and Studies, no 110. Geneva: WHO, 1989.

WHO: Legionnaires' disease, Europe, 1999. Weekly Epidemiological Record, 2000; 75: 347-352

WHO: Malaria and complex emergencies. Weekly Epidemiological Record, 2000; 75: 217 ([http://mosquito.who.int/docs/fs1\\_e.htm](http://mosquito.who.int/docs/fs1_e.htm))

WHO: Onchocerciasis: Fact Sheet no 95 (Feb 2000). <http://www.who.int/inf-fs/ed/fact095.html>

WHO: Report of the WHO Informal Consultation on schistosomiasis control. Geneva: WHO, December 1998

WHO: Water for health: taking charge. Geneva: World Health Organization, 2001 (WHO/WSH/WWD/01.1)

WHO/UNICEF. The global water supply and sanitation assessment 2000. Geneva: WHO/UNICEF, May 2000

World Commission on Dams: Dams and Development: a New Framework for Decision-Making. Report, November 2000. Earthscan. <http://www.dams.org/report>

World Health Organization. Guidelines for drinking-water quality, 2nd edition: Volume 3. Surveillance and control of community supplies. 1997, WHO: Geneva.

World Meteorological Organization (WMO): A comprehensive assessment of the freshwater resources of the world. Produced on behalf of UN, UNDP, UNEP, FAO, UNESCO, WMO, World Bank, WHO, UNIDO and SEI. Stockholm: 1997

Yu S-Z. Drinking water and primary liver cancer. In Tang Z-Y, Wu M-C, Xia S-S (Eds) Primary Liver Cancer. Beijing: China Academic Publishers, 1989 pp 30-37

Yu Y, Taylor PR, Li J-Y et al. Retrospective cohort study of risk-factors for oesophageal cancer in Linzian, People's Republic of China. Cancer Causes Control 1993; 4:195-202

Yuan, 1993. Etiological Study of High Stomach Cancer Incidence Among Residents in Wastewater Irrigated Areas, Environmental Protection Science, 1993, 19(1):70-73, in: World Resources Institute, A guide to the global environment: environmental change and human health, New York, Oxford University

Press, p. 122

### **Biographical Sketch**

**Rosalind Stanwell-Smith** MB BCh, M.Sc, FRCOG, FFPHM is scientific adviser to the Royal Institute of Public Health and honorary senior lecturer at the London School of Hygiene and Tropical Medicine. Since 2000 she has worked as an independent public health consultant, following a long-term post as a nationally based consultant epidemiologist to the Communicable Disease Surveillance Centre of the Public Health Laboratory Service, now part of the Health Protection Agency. Her specialist areas at CDSC included water related disease epidemiology, health effects of climate change, imported infection and outbreak investigation. She has also conducted research studies in hospital infection and perinatal epidemiology and established the English confidential enquiry into still births and deaths in infancy (CESDI). She is the immediate Past President of the Section of Epidemiology and Public Health of the Royal Society of Medicine and is the honorary secretary to the John Snow Society, which commemorates the pioneering work of Dr John Snow in the epidemiology of cholera and anaesthesia.