

RECREATION IN NATURAL WATER RESOURCES

R. M. McNinch

Center for Water Sciences, Michigan State University, East Lansing, Michigan, USA

S. Singh

Department of Fisheries and Wildlife, Michigan State University, East Lansing, Michigan, USA

J. B. Rose

Department of Fisheries and Wildlife, Department of Crops and Soil Sciences, Michigan State University, East Lansing, Michigan, USA

Keywords: Recreational water, pollution, bacteria, pathogens, standards, guidelines, swimming, risks.

Contents

1. Introduction
 2. Recreational Waters
 3. Risks while Recreating
 - 3.1. Physical Risks
 - 3.2. Biological Risks
 4. Guidelines and Standards for Monitoring
 5. Pollution
 6. Sources of Fecal Pollution
 - 6.1. Humans
 - 6.2. Septic Tank Systems
 - 6.3. Wastewater Treatment Plants
 - 6.4. Combined Sewer Overflows (CSOs) and Storm Sewers
 - 6.5. Wildlife and Pets
 - 6.6. Animal Feeding Operations (AFOs)
 7. Reservoirs of Fecal Pollution and Toxins
 - 7.1. Sediments and Sand
 - 7.2. Seaweed and Algae
 - 7.3. Harmful Algal Blooms
 8. Impacts and Events
 - 8.1. Global Climate Change
 - 8.2. High Density and Overuse
 - 8.3. Events
 9. Protection of Recreational Water Sites
 - 9.1. Economic Importance
 - 9.2. Continued Monitoring
 - 9.3. Testing Methods and Advances
 - 9.4. Water pollution prevention
 - 9.5. Education
 10. Recreating Safely
- Glossary

Bibliography
Biographical Sketches

Summary

Water is the most valuable resource on this planet and is used for drinking, food supply, community development (industries) and recreational purposes. Population growth around the world, particularly near the coastlines has increased the demand for water and the services associated. Despite the technological and social advances in water engineering and management, there is currently an inadequate infrastructure to protect the coastal and recreational waters of the world from increasing pressures of pollution, overuse and climate change.

Anthropogenic activities have been instrumental in changing the climate and animal behavior; average seasonal temperatures and precipitation have shown different trends in recent years. Climate change can alter water quantity and quality, shoreline structure, wildlife habitat and waterfowl behavior. Existing waste treatment facilities may no longer be able to handle the increasing waste load which subsequently is released into recreational waters, causing eutrophication and algal blooms. In addition to water, sediments are now being recognized as a reservoir for pathogens and toxic metals. Once re-suspended, these organisms and materials can enter the water column and lead to illness in recreational users and aquatic animals.

Monitoring recreational waters is important in identifying the risk to recreational waters and public health. Research is being done to develop faster and more accurate measures for water quality testing. The need for these tools to be cost effective and accessible for monitoring agencies facing funding shortages is imperative. More effort needs to be undertaken for recreational water management and clean-up programs for already polluted waters. Educational programs can inform people of the risks associated with recreational waters and pollution prevention strategies. Inaccessible and polluted waters can affect local and national economies that rely on tourism. With increased public awareness and technology, recreational waters can be protected, maintained and restored for use by future generations.

1. Introduction

Hundreds of millions of people visit “recreational venues” such as lakes, rivers and ocean beaches each year. In the United States it is estimated that more than 180 million people visit the shoreline of the annually, including 60 million people who visit the international Great Lakes (Table 1). The quality and health of natural aquatic systems around the world are increasingly affected by population growth, over use, climate changes and large scale events such as algal blooms and oil spills. Water quality degradation is caused by numerous sources of pollution but particularly, fecal pathogens from human and animal waste associated with public health risks. Human waste inputs come from septic tank systems, waste water treatment plants and combined sewer overflows while animal waste inputs include wildlife, pets and agricultural and domestic animal feeding operations. It is also now known that sand and sediments are associated with reservoirs of these pollutants which can lead to increased physical and biological

threats to those recreating in these waters. Water quality studies, monitoring and standards are common global approaches for protecting and restoring recreational waters of the world.

2. Recreational Waters

Recreational areas including water are defined by the United Nations as: “Recreational land and associated surface water consist of land that is used as privately owned amenity land, parklands and pleasure grounds and publicly owned parks and recreational areas, together with associated surface waters.”

About 70% of the Earth’s surface is covered in water. 96.5 % of this water is contained in saline water (oceans). 1.74% is contained in glaciers and icecaps. 1.7% is groundwater and only 0.008% is freshwater lakes, rivers or swamp water. The remaining fraction of a percent of water is in saline lakes, soil moisture, permafrost and the atmosphere. Only freshwater can readily be used for consumption, however both salt and freshwater can be used for recreation (glaciers may be used for recreational activities such as hiking and viewing). However, most natural water recreation takes place along the coastlines of the world.

The total length of the world’s coastlines is 1,634,700.7 km (Table 2). There are also 19 great lakes (those that have a surface area >10,000 km²) in the world, representing a total surface area of 997,000 km². In terms of surface area, the Caspian Sea is the largest with 374, 000 km², followed by lakes Superior (82,100 km²), Aral (43,000 km²), Victoria (62,940 km²) and Huron (59,500 km²). In terms of volume, the Caspian Sea is the largest with 78,000 km³, followed by lakes Baikal (22,995 km³), Tanganyika (17,827 km³), Superior (12,230 km³) and Malawi (6,140 km³). Some of the most famous recreational waters include the Caspian Sea which has 7,000 km of shoreline (including islands), the Great Lakes which have a total of 17,017 km of shoreline (including the connecting channels and excluding the St. Lawrence River) and Lake Victoria which has 3,440 km of shoreline. (Table 3)

These shorelines around the world are host to thousands of public recreational beaches. In the United States there are 6,099 coastal beaches and Australia has approximately 11,000 beaches. The European Union has 14,345 coastal beaches and 6,749 freshwater bathing areas. This does not include the number of rivers, streams, and ponds that people visit each year that are not formally labelled as public recreational areas though. (Table 1)

| Location | Total Beaches | # Freshwater Beaches | # Marine Beaches | # Visitors/year |
|---------------------------|-----------------------|----------------------|------------------|-----------------|
| United States | 6,099 coastal beaches | No Data | No Data | >180 million |
| United States Great Lakes | No Data | 582 | None | 60 million |
| European Union | 21,094 | 6,749 | 14,345 | No Data |
| Africa | No data | 677 lakes | No Data | No Data |
| Australia | No data | No Data | ~11,000 | 80 million |

Table 1. Numbers of beaches and visitors World-wide

| Location | Total Coastline (km) |
|----------------------------------|----------------------|
| World | 1,634,700.7 |
| North America | 398,835.2 |
| Europe | 325,892.3 |
| Asia (excluding the Middle East) | 288,459.0 |
| South America | 144,566.8 |
| Oceania | 137,772.4 |
| Central America and Caribbean | 73,703.0 |
| Sub-Saharan Africa | 63,124.4 |
| Middle East and North Africa | 47,281.9 |

Table 2. Coastlines of the World

| Name of Lake | Surface Area (km ²) | Name of Lake | Volume (km ²) |
|--|---------------------------------|--|---------------------------|
| Caspian (Russia) | 374,000 | Caspian (Russia) | 78,200 |
| Superior (Canada & US) | 82,100 | Baikal (Russia) | 22,995 |
| Aral (Kazakhstan & Uzbekistan) | 43,000 | Tanganika (Burundi, Dem. Rep. Congo, Tanzania, & Zambia) | 17,827 |
| Victoria (Kenya, Tanzania & Uganda) | 62,940 | Superior (Canada & US) | 12,230 |
| Huron (Canada & US) | 59,500 | Malawi (Malawi, Mozambique & Tanganika) | 6140 |
| Michigan (US) | 57,750 | Michigan (US) | 4920 |
| Tanganika (Burundi, Dem. Rep. Congo, Tanzania, & Zambia) | 32,000 | Huron (Canada & US) | 3537 |
| Baikal (Russia) | 31,500 | Victoria (Kenya, Tanzania & Uganda) | 2518 |
| Great Bear | 31,326 | Great Bear | 2292 |

| | | | |
|-------------------------|--------|-------------------------|------|
| (Canada) | | (Canada) | |
| Tonle Sap (Cambodia) | 30,000 | Great Slave (Canada) | 2088 |

Table 3. Ten largest lakes of the World

3. Risks while Recreating

Recreational activities that take place in natural waters, such as swimming, boating and rafting, are popular across the world. However these activities are not always safe as there are both physical and biological risks involved.

3.1. Physical Risks

Physical risks at natural waters are usually associated with running and falling and diving in areas that are too shallow and contain rocks and other debris, which can cause cuts, bone breakage and head injuries. Drowning is also a risk at beaches, particularly where large waves may submerge swimmers heads and/or undertows occur (also known as rip currents and run-outs). These undertows can carry swimmers out away from the shore and into deeper, more dangerous waters quickly. Drowning is also a possibility when rafting, boating, tubing or water and jet skiing. During participation in these activities people can end up in deep or fast moving waters that are unsafe. In addition, exposure to the sun can cause damage to people's skin, with long term exposure associated with cancer. Heat stroke may occur where temperatures become too high. People who are most susceptible and should take extra precaution when exposed to the effects of the sun and heat include young children and elderly people.

3.2. Biological Risks

In addition to the physical risks of recreating in natural waters there are also biological risks, or risks which manifest as a recreational water illness caused by microbial pathogens. *[Note: biological risks associated with sharks and jellyfish for example are small in contrast to risks from microorganisms and are not included.]* Recreational water illnesses are acquired by swallowing, breathing or having physical contact with contaminated water. Contamination may be from anthropogenic waste materials produced from the activities of humans (e.g., sewage from cities) but can also occur from natural processes such as blooms of hazardous algae. Human and animal wastes are the most common sources of harmful pathogens which cause recreational water illnesses during recreational activities.

Humans and animals excrete microbes in their feces. Some of these microbes have the ability to survive for extended periods of time in the environment and are then transferred to other humans or animals through the oral route. The fecal-oral route refers to this pathway; excretion into the environment followed by ingestion by a susceptible individual, facilitating pathogen transfer between individuals and the environment. Microbes that are transferred from human to human by the fecal-oral route are referred to as human pathogens and include viruses and some specific bacteria such as *Shigella*. Microbes that are transferred from animal fecal material to humans are referred to as

enteric zoonotic pathogens and include *Cryptosporidium* and *Escherichia coli* (*E. coli*) O157:H7. People come into contact with these microbes by swallowing contaminated recreational waters, breathing aerosols or touching contaminated surfaces (e.g., sand) and then through their hands, transfer microbes to self and others.

Skin, wound, ear, eye, respiratory, neurological and gastrointestinal infections are among the most common recreational water illnesses. According to the World Health Organization (WHO) the most frequent adverse health outcome associated with exposure to fecally contaminated recreational water is enteric illness. Enteric illnesses are associated with the gastrointestinal tract (stomach and intestines) and most commonly associated with diarrhea, often referred to as AGI, acute gastrointestinal illness. Diarrhea illnesses can be caused by microbes such as bacteria including some *E. coli* strains (including *E. coli* O157:H7), *Shigella* and *Campylobacter*. Protozoan parasites associated with recreational water illness include *Cryptosporidium* and *Giardia* and some viruses include Adenoviruses and Noroviruses. Young children, elderly and immuno-compromised people are most at risk for becoming ill due to exposure to these harmful microbes. Table 4 lists these microbes and Table 5 presents some of the outbreaks that have been documented from fecal or urine contamination of lakes or from other microbial risks.

An outbreak occurs when many people visiting the lake or river at a similar time become ill. Most outbreaks are documented at lakes and are caused by unidentified microbes in which the people become ill with AGI. It is suspected that most of the pathogens such as Norovirus are coming from human feces but others could come from human or animal wastes such as the parasites *Cryptosporidium* and *Giardia* and bacteria such as *E. coli* O157:H7. The bacterium *Leptospira* comes from animal urine and causes Leptospirosis a disease associated with swimming, wading and whitewater rafting in contaminated lakes and rivers. It is found throughout the world but mostly in temperate or tropical climates. In addition to fecal pathogens, there are also widespread impacts due to the Schistosome parasites which are associated with snails and birds including ducks, which are part of the parasites normal life cycle. Exposure to these parasites, which are found in both marine and fresh waters, causes what is known as swimmers itch. In addition *Vibrio*, a naturally-occurring bacterium in marine systems, causes wound infections that can result in hospitalization and death.

| Microbe | Disease | Incubation time |
|--|---|---------------------|
| Bacteria | | |
| <i>Escherichia coli</i> (O157:H7) from cattle and human feces | Gastrointestinal | 12 hours-3 days |
| <i>Campylobacter</i> spp. From animal, bird, cattle and human feces | Gastrointestinal | 2-5 days |
| Cyanobacteria, known as blue green algae, naturally-occurring associated with nutrients. | Gastrointestinal , respiratory and nervous system | 30 minutes – 2 days |
| <i>Leptospira</i> from animal urine | Fever, headache, vomiting. | 2 days- 4 weeks |
| <i>Salmonella</i> spp. from humans and animal feces | Gastrointestinal | 1-3 days |
| <i>Shigella</i> spp. from human feces only | Bacillary dysentery | 2-4 days |
| <i>Vibrio</i> spp naturally occurring in marine systems, | Gastrointestinal, | 2-4 days |

| | | |
|--|--|--|
| | liver involvement, wound infections | |
| Parasites <i>Cryptosporidium parvum</i> from animal and human feces & <i>C. hominus</i> (oocysts) from human feces only. <i>Entamoeba histolytica</i> (cysts) from human feces only. <i>Giardia</i> (cysts) from animal and human feces <i>Schistosomes</i> from bird feces and found in snails. | Diarrhoea Amoebic dysentery Diarrhoea Swimmers Itch, cercarial dermatitis | 2-10 days 1-7 weeks 1-2 weeks Minutes |
| Virus from human feces only Adenovirus Hepatitis A Norwalk virus Rotavirus | Respiratory Hepatitis Diarrhoea, vomiting Diarrhoea, vomiting | <10 days 15-50 days 1-2 days |

Table 4. Common bacteria, parasites and viruses which may cause recreational illnesses

| Country/area | Years | Outbreaks/Cases | Pathogens |
|-------------------------|-----------|--|---|
| USA Lakes and rivers | 1999-2004 | 44 outbreaks 1091 cases | AGI (56%) Norovirus (14%) <i>Cryptosporidium</i> (10%) <i>E.coli</i> (with 0157) (9%) <i>Giardia</i> (5%) <i>Shigella</i> (5%) |
| Sweden Lake | 2004 | 2 lakes 163 cases | Norovirus |
| Australia Lakes | 1990-06 | 6 studies 1326 cases | Cyanobacteria |
| US Lakes/ponds | 1995 | Hundreds of cases | Shistosomiosis Swimmers Itch |
| Gulf of Mexico | 2003-04 | 142 cases 70 hospitalizations 9 deaths | <i>Vibrio</i> wound infections |
| Australia | 1998-2004 | 883 cases | Leptosporosis |

Table 5. Select recreational outbreaks

-
-
-

TO ACCESS ALL THE 31 PAGES OF THIS CHAPTER,
Visit: <http://www.eolss.net/Eolss-sampleAllChapter.aspx>

Bibliography

Bisson J.W. and V.J. Cabelli. (1979). *Membrane filter enumeration method for Clostridium perfringens*. Applied and Environmental Microbiology. 37(1):55-66. [This article explains enumeration methods for *C. perfringens*.]

Cabelli, V. (1983). *Health Effects Criteria for Marine Recreational Waters*. (EPA-600/ 1-80-031). USEPA. Office of Research and Development. Washington, D.C. [This report presents a mathematical relationship between quality of water, as determined by the fecal indicator bacteria, enterococci and swimming associated rates of gastrointestinal symptoms in bathers.]

Centers for Disease Control and Prevention. (2007). *Healthy Swimming*. Retrieved July, 2007 from <http://www.cdc.gov/healthyswimming/index.htm>. [This website provides information for the general public and health professionals regarding recreational water illnesses.]

Dufour, A.P. (1984). *Health Effects Criteria for Fresh Recreational Waters*. (EPA-600/1-84-004). USEPA. Office of Research and Development. Washington, D.C. [This report provides an assessment of the relationship between microbiological indicators of water quality and illnesses that may have resulted from swimming.]

Environment Canada. (2007). *Algal toxins and taste and odour*. Retrieved July, 2007 from <http://www.nwri.ca/threatsfull/ch2-1-e.html>. [This website provides information regarding harmful algal toxins.]

European Union. (2006). *Directive 2006/7/EC of the European Parliament and of the Council of 15 February 2006 concerning the management of bathing water quality and repealing Directive 76/160/EEC*. Official Journal of the European Union. Retrieved July, 2007 from http://europa.eu.int/eur-lex/lex/LexUriServ/site/en/oj/2006/l_064/l_06420060304en00370051.pdf. [This is the official revised bathing water directive of the European Union.]

Exxon Valdez Oil Spill Trustee Council. (2007). *Injured Resources and Services*. Retrieved August, 2007 from <http://www.evostc.state.ak.us>. [This official website for the Trustee Council of the Exxon Valdez oil spill provides information regarding clean-up and restoration efforts.]

Kalff, J. (2002). *Limnology; Inland Water Ecosystems*, 41-52 pp. Prentice Hall Inc. Upper Saddle River, New Jersey. [This chapter provides information on water resources, water pollution and lakes.]

Lipp, E.K., Kurz, R., Vincent, R., Rodriguez-Palacios, C., Farrah, S.R. and J.B. Rose. (2001). *The Effects of Seasonal Variability and Weather on Microbial Fecal Pollution and*

Enteric Pathogens in a Subtropical Estuary. Estuaries. 24(2): 266-276 [This study addressed distribution and seasonal changes in microbial indicators and human pathogens in shellfish and recreational waters.]

Natural Resources Defense Council (NRDC). (2007). *Testing the Waters 2007*. Retrieved August, 2007 from <http://www.nrdc.org/water/oceans/ttw/exesum.pdf>. [This website links to the 2007 Testing the Waters Report which details beach closures in the United States for 2006.]

Pearce, F. (2006). *Uganda pulls plug on Lake Victoria*. New Scientist. Retrieved August, 2007 from <http://environment.newscientist.com/channel/earth/mg18925384.100-uganda-pulls-plug-on-lake-victoria.html>. [This article details the use of Lake Victoria for peoples livelihood.]

Prescott, L.M., J.P. Harley and D.A. Klein. (1999). Microbiology, Fourth Edition. WCB/McGraw-Hill. [This is a microbiology textbook.]

Rabinovici, S.J.M., Bernknopf, R. L, Wein, A.M., Coursey, D.L. and R. L. Whitman. (2004). *Economic and Health Risk Trade-Offs of Swim Closures at a Lake Michigan Beach*. Environmental Science and Technology 38(10): 2737-2745. [This paper uses a case study to analyze the economic, health and recreational implications of swim closures.]

Republic of Azerbaijan. (2007). *The formation and physiography of the Caspian Sea*. Retrieved July, 2007 from http://www.country.az/_Geography/_Caspian/caspian_02_e.html. [This article explains the physiography of the Caspian Sea, including shoreline length.]

Shibata, T., Solo-Gabriele, H.M., Fleming, L.E., and S. Elmir. (2004). *Monitoring marine recreational water quality using multiple microbial indicators in an urban tropical environment*. Water Research 38: 3119-3131. [This study examined the relationship between indicator microbes and determined that exceedances of water quality guidelines and frequency of beach advisories is dependant on the indicator microbe used.]

Singh, S. (2007). *Investigation of Bacterial Fecal Indicators and Coliphage Virus in Sediment and Surface Water of Parks and Beaches along the Grand River (MI) and Lake Michigan (MI)*. Masters Thesis. Michigan State University. [This thesis report provides details on bacterial contamination in sand and sediments.]

Surfrider Foundation Australia. (1998). *Ocean outfalls-there is a solution to ocean pollution*. Retrieved July, 2007 from <http://www.surfrider.org.au/publications/media/ocean.php>. [This article details the number of ocean outfalls and associated issues in Australia.]

United Nations. (2007). *United Nations Statistics Division, Term: Recreational land and associated surface water*. Retrieved July, 2007. from <http://unstats.un.org/unsd/sna1993/glossform.asp?getitem=471>. [This website defines the term 'recreational land and associated surface water' according to the United Nations.]

United States Department of the Interior. (2003). *Ohio District's Microbiology Program*. United States Geological Survey. Retrieved July, 2007 from <http://oh.water.usgs.gov/microbiol.html>. [This website provides information on beach monitoring methods and predictive models.]

United States Environmental Protection Agency (USEPA). (1986). *Ambient water quality criteria for bacteria*. (EPA-440/5-84/002). Office of Water Regulations and Standards, Criteria and Standards Division. Washington, D.C. [This document details the water quality recommendations for ambient waters.]

United States Environmental Protection Agency (USEPA). (2001). *Total Coliform Rule: A quick reference guide*. (EPA 816-F-01-035). Office of Water. www.epa.gov/safewater. [This site provides total coliform bacteria guidelines.]

United States Environmental Protection Agency (USEPA). (2002). *Implementation Guidance for Ambient Water Quality Criteria for Bacteria*. (EPA-823-B-02-003). Office of Water. Washington, D.C. [This document provides water quality criteria for ambient waters.]

United States Environmental Protection Agency (USEPA). (2003). *Bacterial Water Quality Standards for Recreational Waters (Freshwater and Marine Waters)*. (EPA-823-R-03-008). Office of Water. Washington, D.C. [This document details the water quality standards for fresh and marine recreational waters.]

United States Environmental Protection Agency (USEPA). (2004). *National List of Beaches*. Retrieved July, 2007. <http://www.epa.gov/waterscience/beaches/list/list-of-beaches.pdf>. [This document lists the total number of beaches monitored and not monitored by state.]

United States Environmental Protection Agency (USEPA). (2006). *Evaluation Report; EPA provided quality and timely information regarding wastewater after Hurricane Katrina. Report No. 2006-P-00018*. Office of Inspector General [This document provides information regarding Hurricane Katrina and associated wastewater treatment facilities.]

United States Environmental Protection Agency (USEPA). (2007). *Great Lakes Fact Sheet*. Retrieved July, 2007 from <http://www.epa.gov/glnpo/factsheet.html>. [This site provides information regarding the physical size and surrounding populations of the Great Lakes.]

Van Asperen, I.A., Medema, G., Borgdorff, M.W., Sprenger, M.J.W, and A.H. Havelaar. (1998). *Risk of gastroenteritis among triathletes in relation to faecal pollution of fresh waters*. International Journal of Epidemiology 27: 309-315. [This study examined the cases of gastroenteritis among athletes who swam and those who did not.]

Weiskel, P., Howes, B. and G. Heufelder. (1996). *Coliform contamination of a coastal embayment: Sources and Transport pathways*. Environmental Science and Technology 30:1872-1881. [This article identifies waterfowl as one source of fecal pollution on beaches.]

Wither, A., M. Rehfisch and G. Austin. (2003). *The impact of bird populations on the microbial quality of bathing waters*. Diffuse Pollution Conference, Dublin. [This research focussed on quantifying the fecal indicator loads of birds in order to assess its significance on water quality.]

Whitman, R.L., Shively, D.A., Pawlik, H., Nevers, M.B. and M.N. Byappanahalli. (2003). *Occurrence of Escherichia coli and Enterococci in Cladophora (Chlorophyta) in nearshore water and beach sand of Lake Michigan*. Applied and Environmental Microbiology 69(8):4714-4719. [This study examined the survival time of *E. coli* and enterococci in *Cladophora* mats.]

World Health Organization (WHO). (1999). *Health-Based Monitoring of Recreational Water: The Feasibility of a New Approach (The 'Annapolis Protocol')*. (WHO/SDE/WSH/99.1). World Health Organization – Sustainable Development and Healthy Environments. [This document provides an assessment of the effectiveness of present approaches to monitoring and managing microbial risks at fresh and coastal recreational waters.]

World Health Organization (WHO). (2003). *Guidelines for safe recreational water environment Volume 1 - Coastal and fresh waters*. Retrieved July, 2007 from http://whqlibdoc.who.int/publications/2003/9241545801_contents.pdfhttp://www.who.int/water_sanitation_health/bathing/srwe1/en/. [This document provides the WHO guidelines for coastal and fresh waters.]

Biographical Sketches

Rachel M. McNinch recently graduated with her Masters degree from the Department of Fisheries and Wildlife at Michigan State University, after having earned her Bachelors degree from the same department. Her graduate research focused on waterfowl fecal bacteria content and waterfowl fecal bacteria loading effects on beach sand and water. She works for the Center for Water Sciences at Michigan State University and is currently using geographic information systems (GIS) to look at the state of Michigan's *Escherichia coli* database. Research interests include recreational water quality and human health.

Shikha Singh is a recent Masters graduate from Michigan State University in the department of Fisheries and Wildlife. Her research interests are focused upon water quality and environmental health. Currently, she is focused on identifying relationships with surface water and sediments near Lake Michigan using fecal indicator bacteria. She grew up in London, Ontario and graduated from the University of Western Ontario from the department of Biology before migrating to Michigan for further studies. When doing research, she has an interest in international aquatic issues and photography and can be seen traipsing through areas like Siberia, Russia, Portugal and India.

Dr. Joan B. Rose currently holds the Homer Nowlin Chair in Water Research at Michigan State University after receiving her PhD from the University of Arizona and spending 14 years at the University of South Florida. Dr. Rose is an international expert in water microbiology, water quality and public health safety publishing more than 250 manuscripts. She is considered one of the authorities on *Cryptosporidium*. She is currently co-director of the Center for Water Sciences which includes work with the Great Lakes and Human Health Center of NOAA. She is also Co-Director of the EPA/DHHS Center for Advancing Microbial Risk Assessment. She is currently the chair of the EPA Science Advisory Board Committee on Drinking Water and serves as an advisor to the US-Canada International Joint Commission.