

## **NEW OPPORTUNITIES REVEALED BY BIOTECHNOLOGICAL EXPLORATIONS OF EXTREMOPHILES**

**Mircea Podar and Anna-Louise Reysenbach**

*Department of Biology, Portland State University, Portland, OR 97201, USA.*

**Keywords:** extremophiles, genomics, biotechnology, enzymes, metagenomics.

### **Contents**

1. Introduction
2. Extremophiles and Biomolecules
3. Extremophile Genomics Exposing the Biotechnological Potential
4. Tapping into the Hidden Biotechnological Potential through Metagenomics
5. Unexplored Frontiers and Future Prospects

Acknowledgements

Glossary

Bibliography

Biographical Sketches

### **Summary**

Over the past few decades the extremes at which life thrives has continued to challenge our understanding of biochemistry, biology and evolution. As more new extremophiles are brought into laboratory culture, they have provided a multitude of new potential applications for biotechnology. Furthermore, more recently, innovative culturing approaches, environmental genome sequencing and whole genome sequencing have provided new opportunities for biotechnological exploration of extremophiles.

### **1. Introduction**

Organisms that live at the extremes of pH (>pH 8.5,< pH 5.0), temperature (>45°C, <15°C), pressure (>500 atm), salinity (>1.0M NaCl) and in high concentrations of recalcitrant substances or heavy metals (extremophiles) represent one of the last frontiers for biotechnological and industrial discovery. As we learn more about the extremes at which life can survive and thrive, more of these extremophiles are brought into culture and their genomes sequenced. In many cases, biotechnological applications of extremophiles and their biomolecules (e.g. enzymes) have been the driving force in both academic and industrial research of these organisms. Extremophiles and extremozymes occupy an important place in the multibillion dollars environmental biotechnology industry, with applications spanning agricultural, biomedical and industrial sectors. Due to the highly competitive nature of industrial R&D, in most cases the path from an extremophile to a successful commercial application is not documented in peer reviewed scientific publications and can be partially followed through patents, biotechnology meetings and company websites. In this review, we will consider a few new developments as several recent reviews focus on the biotechnological applications of extremophiles

## 2. Extremophiles and Biomolecules.

The most direct application of extremophiles in biotechnological processes involves the organisms themselves. Among the most established is biomining (bioleaching), in which microbial consortia are used to extract metals such as copper, cobalt, gold and uranium from ores. This has received considerable interest lately, propelled by developments in microbial isolation and the application of genomic approaches for studying the individual organisms and their community. These processes involve iron or sulphur oxidizing acidophilic microorganisms adapted to different temperature ranges, from mesophiles (bacteria such as *Acidithiobacillus*, *Leptospirillum* and the archaea *Ferroplasma*) to thermophiles (archaea from the genera *Sulfolobus*, *Metallopharea* and *Acidianus*) [see also – *Biotechnology of Archaea*]. When the activity of such extreme acidophiles is not controlled in biomining operations, they can lead to acid mine drainage (AMD), which causes considerable environmental damage. While the microbes responsible for bioleaching and AMD are inherently adapted to extremely low pH and high concentration of metals, they are part of an open and dynamic consortia and continuously adapt to the particular conditions they are facing. In commercial bioleaching operations this can result in the selection of more robust and efficient strains, with reduced sensitivity to metal toxicity. For example, arsenic resistance genes have been horizontally transferred via a transposon from an unidentified bacterium to *Acidithiobacillus caldus* and *Leptospirillum ferriphilum*, resulting in substantially increased resistance to arsenic in gold-bearing arsenopyrite bioreactors. This demonstrates that taxonomy alone is not necessarily predictive of the physiological fitness of an individual organism and that the community gene pool can impact the adaptability of the constituent members across taxonomic barriers. Such strain engineering and selection for improved biotechnological characteristics may be applied to a wide range of bioremediation projects [see also– *Bioremediation*] as has been done for contaminating petroleum hydrocarbons accidentally released in arctic environments.

Most of the applications involving extremophiles are based on their biomolecules, primarily enzymes but also other proteins (e.g. cryoprotectant antifreeze proteins), lipids, various small molecules. The most well known example of a successful application of an extremophile-derived product is *Taq* DNA polymerase which was isolated from *Thermus aquaticus*, first isolated from a geothermal spring from Yellowstone National Park. This enzyme approaches sales of about half a billion dollars per year. The molecular biology research tool enzymes not only include a wide range of other thermostable polymerases and ligases, but also enzymes isolated from psychrophilic organisms, such as the “Antarctic phosphatase” from New England Biolabs. Genencor commercialized one of the first industrial extremozymes for use in textiles detergents, a cellulase isolated from an alkaliphilic bacterium from an east African soda lake. Numerous other examples are reviewed by Antranikian [see – *Biotechnology of Archaea*].

Enzymes that have optimal activity at extreme temperatures and pH are widely used in household detergents and in the food, textile, pulp and paper, leather processing, chemical intermediates industries. For each application, the enzymes have to fulfill numerous requirements related to such features as activity and stability, substrate

specificity and enantioselectivity. As a result, natural enzymes often are not optimal for the desired biotechnological application. Consequently, a variety of approaches have been used to modify enzyme properties such as error-prone PCR, saturation mutagenesis, structure-based protein engineering and in vitro evolution approaches. Such approaches are best combined with genetic selection or high throughput screening, to identify the rare mutants that approach the target characteristics, followed by an iterative process of building fitness to the resulting variants. For example, thermostability was built into a suite of other optimal enzymatic characteristics of pectinases used in cotton fabric processing and increased alkaline stability was obtained for a previously engineered xylanase from the fungus *Trichoderma reesei*. Furthermore, based on genomics, structural data and computational modeling, certain protein architecture and amino acids usage trends provide clues to mechanisms of protein thermostability and promise to lead to predictive protein thermostabilization.

### 3. Extremophile Genomics Exposing the Biotechnological Potential

Extremophiles, including eukarya (e.g. *Alvinella pompejana*, *Tetrahymena thermophila*, *Dunaliella salina*), have been prime subjects for genomic sequencing projects in an effort to understand the fundamental mechanisms of adaptation to specific environments and for practical applications. Recently, four psychrophilic bacteria have been published: two from arctic sediments, *Desulfotaleas psychrophila* and *Colwellia psycherythraea*, one from coastal Antarctic waters, *Pseudoalteromonas haloplankis* and *Photobacterium profundum*, a deep sea bacterium that is both adapted to low temperatures and high pressure. While there does not appear to be a distinct genomic trait unifying these cold adapted organisms, they share several characteristics, most notably a membrane with increased proportion of polyunsaturated and branched fatty acids to increase fluidity at low temperatures (and high pressure), and the presence of cold shock proteins which are believed to increase translation efficiency by destabilizing secondary structures in mRNA. Additionally, cryoprotectants increase the capacity for nutrient uptake. Due to higher solubility of oxygen at low temperatures and hence potential for oxidative damage within the cell, metabolic reactions that generate reactive oxygen are reduced and molybdopterin-dependent metabolism is eliminated (in *P. haplokantis*) or the number of catalase and superoxide dismutase genes increased (in *C. psycherythraea* and *D. psychrophila*). While there are specific amino acid usage trends that appear to correlate with the psychrophilic proteomes, the most important adaptation for cold adapted enzymes appears to be a high specific activity at low temperatures. This is achieved by a highly flexible catalytic center at the expense of overall reduced protein stability and susceptibility for thermal denaturation. Such enzymatic characteristics are currently exploited in food biotechnological applications [see also– *Fermented Foods and their Processing*] that require low temperatures (e.g. milk and fruit juice processing). Beside cold adaptation, *P. profundum* is also able to withstand high pressure and has become a model organism for understanding piezophily. For example, transport processes and energy generation by cytochrome respiration are impacted by high pressure.

Similar to the situation in cold adaptation, intrinsic properties of nucleic acids, lipids and enzymes/proteins allow thermophiles to flourish in high temperature environments, and specific composition biases and structural adaptations have been identified. The

recent publications describing the genomes of two hyperthermophilic archaea, *Nanoarchaeum equitans* and *Thermococcus kodakaraensis*, one archaeal thermoacidophile (*Sulfolobus acidocaldarius*) and one thermophilic bacterium (*Carboxydotherrmus hydrogenoformans*) while not significantly changing the perception on high temperature adaptation, bring together a number of exciting novelties on various aspects of the biology, genome evolution and metabolic versatility in specific thermophilic environments.

The genome of *N. equitans* came shortly after the discovery of this unusual organism, which represents the first case of an archaeon that is an obligate symbiont or parasite on another archaeon, the marine crenarchaeote *Ignicoccus* sp.. The genome abounds in oddities, from being at the smallest spectrum of cellular genome sizes and encoding virtually no metabolic pathways, to containing a large number of split genes, including uniquely split tRNA genes. One of the split genes is represented by two separate ORFs encoding DNA polymerase B (also fused with a split intein) which has is functional *in vitro* and *trans* spliced at high temperature thus restoring a functional polymerase.

The *Sulfolobus acidocaldarius* genome sequence represents an important landmark in archaeal genomics as this organism is one of the few laboratory genetic systems in Archaea. *S. acidocaldarius* genome has a number of similarities and differences with two other *Sulfolobus* species, and will ease future experimental studies and facilitate overall biotechnological applications of these Crenarchaeota. The bacterial thermophile, *Carboxydotherrmus hydrogenoformans* (35), has a remarkable efficiency to carry out carbon monooxidase metabolism due to the presence of five anaerobic CO dehydrogenase complexes.

This genomic blueprint should allow detailed studies of hydrogenogenesis, which could become an important industrial process for generating hydrogen. The most thermoacidophilic organism known, *Picrophilus torridus* inhabits solfataric environments with pH below 1 and about 60°C, and its low pH adapted enzymes will most likely be found useful for biotechnological applications requiring acidic conditions. To cope with such conditions, *P. torridus* has evolved a membrane with low proton permeability and special lipid composition as well as efficient transport mechanisms to maintain the internal pH at values compatible with biochemical functions. Unlike other organisms, which use sodium ion and ATP-driven primary transporters, *P. torridus* predominantly uses the internal high proton concentration (pH 4) to power a large number of solute secondary transporters.

At the other extreme, the genome of a haloalkaliphilic archaeon from highly alkaline soda lakes, *Natronomonas pharaonis*, has revealed several adaptations to this environment. These include an overall modification of the proteome to increase the fraction of acidic amino acids and reduce protein hydrophobicity, a coating of the cell membrane with glycoproteins and secreted enzymes attached by lipid anchors as well as an efficient transport system for heavy metals and nitrogen compounds which are scarce in hypersaline environments. The halophilic bacterium *Salinibacter ruber* displays similar adaptation mechanisms to hypersaline environments and some of these may have been acquired via lateral gene transfer. No doubt, many more surprises are hidden in these genomes that can be exploited for biotechnological purposes.

-  
-  
-

TO ACCESS ALL THE 11 PAGES OF THIS CHAPTER,  
Visit: <http://www.eolss.net/Eolss-sampleAllChapter.aspx>

### Bibliography

Allen EF and Banfield JF. Community genomics in microbial ecology and evolution. *Nat rev Microbiol.* 2005, 3:489-498. [A review of recent advances in metagenomics, with emphasis on ecological and evolutionary questions that can be addressed using this approach].

Antranikian, G, Vorgias, GE and Bertoldo, C. Extreme environments as a source for microorganisms and novel biocatalysts. *Adv Biochem Engin/Biotechnol*, 2005, 96:219-262. [A detailed review compiled on a large body of literature on the biotechnological aspects of extremophiles].

Arnold, F.H. and Georgiou, G (Eds) *Directed Enzyme Evolution: Screening and Selection Methods.* Methods in Molecular Biology Series, Humana Press, 2003. [A collection of methods and protocols that can be used to generate and identify engineered enzymes that have desired novel characteristics].

Atomi H. Recent progress towards the application of hyperthermophiles and their enzymes. *Curr Opin Chem Biol.* 2005 Apr;9(2):166-73. [A review of novel applications of organisms adapted to high temperatures and enzymes derived from them to industrial processes].

Atomi, H, Matsumi, R and Imanaka, T. Reverse gyrase is not a prerequisite for hyperthermophilic life. *J. Bacteriol.* 2004, 186:4829-4833. [An experimental genetic study that shows that the reverse gyrase gene is not absolutely required for the growth of an archaeon at high temperatures].

Berezovsky, IN and Shakhnovich EI Physics and evolution of thermophilic adaptation. *Proc. Natl. Acad. Sci. USA*, 2005, 102:12742-12747. [A study of structure and sequence determinants of protein adaptation to function at high temperatures].

Berezovsky, IN, Chen, WW, Chi, PJ and Shakhnovich, EI. Entropic stabilization of proteins and its proteomic consequences. *PLoS Comput Biol.* 2005, 1:e47. [The discovery of a novel entropic mechanism of protein thermostability which results in lysine increase and arginine decrease in proteins from hyperthermophiles].

Brakstad OG, Bonaunet K. Biodegradation of Petroleum Hydrocarbons in Seawater at Low Temperatures (0-5 degrees C) and Bacterial Communities Associated with Degradation. *Biodegradation.* 2006 17:71-82. [Application of cold adapted bacteria to biodegradation of oil in polluted arctic marine environments].

Campanaro S, Vezzi A, Vitulo N, Lauro FM, D'Angelo M, Simonato F, Cestaro A, Malacrida G, Bertoloni G, Valle G, Bartlett DH. Laterally transferred elements and high pressure adaptation in *Photobacterium profundum* strains. *BMC Genomics.* 2005, 6:122. [Genomic investigation of mechanisms of adaptation to high pressure in a gamma proteobacterium].

Chen, L, Brugger, K, Skovgaard, M, Redder, P, She, Q, Torarinsson, E, Greve, B, Awayez, M, Zibat, A, Klenk, HP and Garrett RA. The genome of *Sulfolobus acidocaldarius*, a model organism of the Crenarchaeota. *J Bacteriol.* 2005, 187:4992-4999. [Sequencing and annotation of the genome of a *Sulfolobus* species].

Choi JJ, Nam KH, Min B, Kim SJ, Soll D and Kwon S . Protein trans-splicing and characterization of a split family B-type DNA polymerase from the hyperthermophilic archaeal parasite *Nanoarchaeum equitans*. *J Mol Biol.* 2006, 356:1093-1106. [Biochemical characterization of a trans-splicing intein in a *Nanoarchaeum* gene]

- DeLong, EF and Karl, DM. Genomic perspectives in microbial oceanography. *Nature*, 2005, 437:336-342. [Review of recent in marine microbial genomics].
- Egorova, K. and Antranikian, G, Industrial relevance of thermophilic Archaea, *Current Opp Microbiol* 2005, 8:1-7. [Review of biotechnological applications of high temperature-adapted enzymes from Archaea].
- Falb M, Pfeiffer F, Palm P, Rodewald K, Hickmann V, Tittor J and Oesterhelt D. Living with two extremes: conclusions from the genome sequence of *Natronomonas pharaonis*. *Genome Res.* 2005, 15:1336-1343. [Sequencing and annotation of the genome of an archaeon adapted to high salinity and alkaline pH].
- Feller G, Gerday C. Psychrophilic enzymes: hot topics in cold adaptation. *Nat Rev Microbiol.* 2003 1:200-208. [Review of understanding the mechanisms of enzymatic adaptation to low temperature].
- Fenel F, Zitting AJ, Kantelinen A. Increased alkali stability in *Trichoderma reesei* endo-1, 4-beta-xylanase II by site directed mutagenesis. *J Biotechnol.* 2006 121:102-107. [Bioengineering of fungal enzymes to degrade xylan under alkaline conditions].
- Ferrer M, Martinez-Abarca F, Golyshin PN. Mining genomes and 'metagenomes' for novel catalysts. *Curr Opin Biotechnol.* 2005, 16:588-593. [Review on the discovery of biotechnological catalysts from individual microbial genomes and from environmental samples].
- Ferrer M, Golyshina OV, Chernikova TN, Khac hane AN, Martins Dos Santos VA, Yakimov MM, Timmis KN, Golyshin PN. Microbial enzymes mined from the Urania deep-sea hypersaline anoxic basin. *Chem Biol.* 2005, 12:895-904. [Bioprospecting for enzymes adapted to high salinity]
- Fukui, T, Atomi, H, Kanai, T, Matsumi, R, Fujiwara, S and Imanaka, T. Complete genome sequence of the hyperthermophilic archaeon *Thermococcus kodakaraensis* KOD1 and comparison with *Pyrococcus* genomes. 2005, *Genome Res.* 15:352-363. [Comparative genomics of several *Pyrococcus* species].
- Futterer O, Angelov A, Liesegang H, Gottschalk G, Schleper C, Schepers B, Dock C, Antranikian G and Liebl W. Genome sequence of *Picrophilus torridus* and its implications for life around pH 0. *Proc Natl Acad Sci U S A.* 2004, 101:9091-9096. [Mechanisms of adaptation to extremely acid conditions revealed by genomic sequencing and analysis of an acidophilic archaeon].
- Galperin, MY. The secret of being cool. *Environmental Microbiology*, 2005, 7:1499-154. [A retrospective of genomic projects during 2004-2005].
- Gomes, J. and Steiner, W. The biocatalytic potential of extremophiles and extremozymes. *Food Technol and Biotechnol.* 2004, 42:223-235. [Review of biotechnological applications of enzymes from extremophiles, with emphasis on the food industry].
- Hallam SJ, Putnam N, Preston CM, Detter JC, Rokhsar D, Richardson PM, DeLong EF. Reverse methanogenesis: testing the hypothesis with environmental genomics. *Science.* 2004, 305:1457-1462. [The authors used selective enrichment techniques and metagenomic sequencing and analysis to identify genes and infer metabolic characteristics of deep-sea archaeal methanotrophs. ]
- Handelsman, J. Metagenomics: application of genomics to uncultured microorganisms. *Microbiol Mol Biol Rev.* 2004, 68:669-685. [A detailed review on the types of metagenomic approaches used to study microbial communities and mine for useful genes.]
- Head, I.M., Jones, M.D. and Röling, W.F.M. Marine microorganisms make a meal of oil. *Nature Reviews Microbiol.* 2006, 4:173-182. [An update on the microbiology of petroleum degradation in the sea].
- Hoppe, C and Schomburg, D. Prediction of protein thermostability with a direction- and distance-dependent knowledge-based potential. *Protein Sci.* 2005 14:2682-2692.[A computational study to predict protein thermostability].
- Huber H, Hohn MJ, Rachel R, Fuchs T, Wimmer VC and Stetter KO. A new phylum of Archaea represented by a nanosized hyperthermophilic symbiont. *Nature*, 2002, 417:63-67. [The discovery of a new group of Archaea, the first obligate parasite/symbiont on another archaeon.]
- Khachane, AN, Timmis, KN and Dos Santos, VA. Uracil content of 16S rRNA of thermophilic and psychrophilic prokaryotes correlates inversely with their optimal growth temperatures. *Nucleic Acids*

Res. 2005, 33:4016-4022. [Discovery of a correlation between the optimum life temperature and the uracil content in the ribosomal RNA of microbes].

Korkegian, A, Black, ME, Baker, D and Stoddard, BL. Computational thermostabilization of an enzyme. *Science*, 2005, 308:857-860. [A predictive protein thermostabilization study combined with structural and metabolic analyses.]

Lorenz P and Eck J. Metagenomics and industrial applications. *Nat Rev Microbiol*. 2005, 3:510-516. [A chronological and approach-specific review on how genomics and metagenomics have impacted the biotechnology industry.]

Makarova, KS, Wolf, Y and Koonin, EV. Potential genomic determinants of hyperthermophily. *Trends Genet*. 2003, 19:172-176. [An attempt to identify the genes that are characteristic to organisms that live at high temperature].

Medigue C, Krin E, Pascal G, Barbe V, Bernsel A, Bertin PN, Cheung F, Cruveiller S, D'Amico S, Duilio A, Fang G, Feller G, Ho C, Mangenot S, Marino G, Nilsson J, Parrilli E, Rocha EP, Rouy Z, Sekowska A, Tutino ML, Vallenet D, von Heijne G, Danchin A. Coping with cold: the genome of the versatile marine Antarctica bacterium *Pseudoalteromonas haloplanktis* TAC125. *Genome Res*. 2005 15:1325-35. [Genomic adaptation to low temperature]

Methe BA, Nelson KE, Deming JW, Momen B, Melamud E, Zhang X, Moulton J, Madupu R, Nelson WC, Dodson RJ, Brinkac LM, Daugherty SC, Durkin AS, DeBoy RT, Kolonay JF, Sullivan SA, Zhou L, Davidsen TM, Wu M, Huston AL, Lewis M, Weaver B, Weidman JF, Khouri H, Utterback TR, Feldblyum TV, Fraser CM. The psychrophilic lifestyle as revealed by the genome sequence of *Colwellia psychrerythraea* 34H through genomic and proteomic analyses. *Proc Natl Acad Sci U S A*. 2005 102:10913-10918. [Genomic adaptation to low temperature]

Mongodin EF, Nelson KE, Daugherty S, Deboy RT, Wister J, Khouri H, Weidman J, Walsh DA, Papke RT, Sanchez Perez G, Sharma AK, Nesbo CL, MacLeod D, Baptiste E, Doolittle WF, Charlebois RL, Legault B, Rodriguez-Valera F. The genome of *Salinibacter ruber*: convergence and gene exchange among hyperhalophilic bacteria and archaea. *Proc Natl Acad Sci U S A*. 2005, 102:18147-18152. [Genomics of microorganisms adapted to high salt environments]

Olson GJ, Brierley JA, Brierley CL. Bioleaching review part B: progress in bioleaching: applications of microbial processes by the minerals industries. *Appl Microbiol Biotechnol*. 2003 63:249-57. [The microbiology and biochemistry of microbial biomining].

Podar M, Eads JR, Richardson TH. Evolution of a microbial nitrilase gene family: a comparative and environmental genomics study. *BMC Evol Biol*. 2005, 5:42. [A combination of genomic, metagenomic and biochemical data with phylogenetic analysis leading to new hypotheses on the evolution and function of nitrilases.]

Rabus R, Ruepp A, Frickey T, Rattei T, Fartmann B, Stark M, Bauer M, Zibat A, Lombardot T, Becker I, Amann J, Gellner K, Teeling H, Leuschner WD, Glockner FO, Lupas AN, Amann R, Klenk HP. The genome of *Desulfotalea psychrophila*, a sulfate-reducing bacterium from permanently cold Arctic sediments. *Environ Microbiol*. 2004, 6:887-902. [Genome sequencing and analysis of a bacterium adapted to low temperatures].

Randau L, Munch R, Hohn MJ, Jahn D and Soll D. Nanoarchaeum equitans creates functional tRNAs from separate genes for their 5'- and 3'-halves. *Nature*, 2005, 433:537-541. [The computational prediction and experimental confirmation for the presence of split tRNA genes in *N. equitans*, the only organism known so far in which such tRNA trans-splicing occurs.]

Ram RJ, Verberkmoes NC, Thelen MP, Tyson GW, Baker BJ, Blake RC 2nd, Shah M, Hettich RL, Banfield JF. Community proteomics of a natural microbial biofilms. *Science*. 2005, 308:1915-20. [A metaproteomic characterization of a biofilm from an acid mine drainage site].

Rawlings, DE. Characteristics and adaptability of iron- and sulfur-oxidizing microorganisms used for the recovery of metals from minerals and their concentrates. *Microb Cell Fact*. 2005 4:13. [A research review on the physiology and genetic adaptability of biomining microbes supplemented with new data.]

Rice G, Stedman K, Snyder J, Wiedenheft B, Willits D, Brumfield S, McDermott T, Young M. Novel viruses from extreme thermal environments. *Proc Natl Acad Sci U S A* 2001, 98:13341-5. [Discovery of

novel viruses from high temperature environments in the Yellowstone National Park].

Robertson, D.E. and Noel, J.P. (Eds), *Methods in Enzymology: Protein Engineering*, vol. 388, Elsevier Inc. 2004. [A volume dedicated to various methods for in vitro protein engineering].

Robertson DE, Chaplin JA, DeSantis G, Podar M, Madden M, Chi E, Richardson T, Milan A, Miller M, Weiner DP, Wong K, McQuaid J, Farwell B, Preston LA, Tan X, Snead MA, Keller M, Mathur E, Kretz PL, Burk MJ, Short JM. Exploring nitrilase sequence space for enantioselective catalysis. *Appl Environ Microbiol.* 2004, 70:2429-2436. [A metagenomic project for the discovery of novel nitrilases with activity against selected compounds of industrial importance].

Rohwerder T, Gehrke T, Kinzler K, Sand W. Bioleaching review part A: progress in bioleaching: fundamentals and mechanisms of bacterial metal sulfide oxidation. *Appl Microbiol Biotechnol.* 2003 63:239-48. [The microbiology and biochemistry of environmental side effects of microbial biomining].

Solbak AI, Richardson TH, McCann RT, Kline KA, Bartnek F, Tomlinson G, Tan X, Parra-Gessert L, Frey GJ, Podar M, Luginbuhl P, Gray KA, Mathur EJ, Robertson DE, Burk MJ, Hazlewood GP, Short JM, Kerovuo J. Discovery of pectin-degrading enzymes and directed evolution of a novel pectate lyase for processing cotton fabric. *J. Biol. Chem.* 2005, 11:9431-9438. [A metagenomic study on the discovery of pectin degrading enzymes followed by protein engineering to improve the characteristics of selected hits].

Steven B, Leveille R, Pollard WH, Whyte LG. Microbial ecology and biodiversity in permafrost. *Extremophiles* 2006, 10: Mar 21; [Epub ahead of print]. [A review on the microbiology of permafrost].

Tringe SG, von Mering C, Kobayashi A, Salamov AA, Chen K, Chang HW, Podar M, Short JM, Mathur EJ, Detter JC, Bork P, Hugenholtz P, Rubin EM. Comparative metagenomics of microbial communities. *Science.* 2005, 308:554-7. [A metagenomic study of three whale fall microbial communities and a soil sample. Comparative analysis of the data with other public metagenomic sequences revealed specific gene signatures and metabolic features that can be used to identify and characterize different consortia.

Tung HC, Bramall NE, Price PB. Microbial origin of excess methane in glacial ice and implications for life on Mars. *Proc Natl Acad Sci U S A.* 2005, 102:18292-18296.

Tyson GW, Chapman J, Hugenholtz P, Allen EE, Ram RJ, Richardson PM, Solovyev VV, Rubin EM, Rokhsar DS, Banfield JF. Community structure and metabolism through reconstruction of microbial genomes from the environment. *Nature.* 2004, 428:37-43. [The first genomic assembly and detailed microbial Community analysis using random sequencing of environmental DNA libraries.]

Tyson GW, Lo I, Baker BJ, Allen EE, Hugenholtz P, Banfield JF. Genome-directed isolation of the key nitrogen fixer *Leptospirillum ferrodiazotrophum* sp. nov. from an acidophilic microbial community. *Appl Environ Microbiol.* 2005, 71:6319-24. [The first microorganism to be isolated using experimental condition designed based on metagenomic analysis of the microbial community.]

Valenzuela L, Chi A, Beard S, Orell A, Guiliani N, Shabanowitz J, Hunt DF, Jerez CA. Genomics, metagenomics and proteomics in biomining microorganisms *Biotechnol Adv.* 2006 24:197-211. [An update on biomining and the genomic-driven advances in understanding the biochemical processes associated with biomining microbes].

van den Burg B. Extremophiles as a source for novel enzymes. *Curr Opin Microbiol.* 2003 Jun;6(3):213-8. Review. [A review of how enzymes derived from extremophiles can be used for biotechnological processes].

Vezi A, Campanaro S, D'Angelo M, Simonato F, Vitulo N, Lauro FM, Cestaro A, Malacrida G, Simionati B, Cannata N, Romualdi C, Bartlett DH, Valle G. Life at depth: *Photobacterium profundum* genome sequence and expression analysis. *Science.* 2005 307:1459-1461. [A study on the genomic determinants of bacterial adaptation to high pressure and low temperature.]

Ward N. and Fraser C.M. How genomics has affected the concept of microbiology. *Current Opinion Microbiol,* 2005 8:564-571. [An overview on the implications of genomic approaches to the field of microbiology].

Waters, E, Hohn, MJ, Ahel, I, Graham, DE, Adams, MD, Barnstead, M, Beeson, KY, Bibbs, L, Bolanos, R, Keller, M, Kretz, K, Lin, X, Mathur, E, Ni, J, Podar, M, Richardson, T, Sutton, GG, Simon, M, Soll,



D, Stetter, KO, Short, JM and Noordewier M. The genome of *Nanoarchaeum equitans*: insights into early archaeal evolution and derived parasitism. 2003, Proc Natl Acad Sci U S A, 100:12984-12988. [*The first genome of an obligate archaeal parasite/symbiont with numerous unique genomic features. The authors propose that Nanoarchaea are a basal archaeal lineage.*]

Wu M, Ren Q, Durkin AS, Daugherty SC, Brinkac LM, Dodson RJ, Madupu R, Sullivan SA, Kolonay JF, Nelson WC, Tallon LJ, Jones KM, Ulrich LE, Gonzalez JM, Zhulin IB, Robb FT and Eisen JA. Life in Hot Carbon Monoxide: The Complete Genome Sequence of *Carboxydotherrmus hydrogenoformans* Z-2901. PLoS Genet., 2005, 1:e65. [The genomic sequencing and annotation for the bacterium *Carboxydotherrmus hydrogenoformans* and its metabolic reconstruction].

### Biographical Sketches

**Anna-Louise Reysenbach** obtained her PhD in Microbiology from the University of Cape Town in 1987, where she studied the physiology of the obligate anaerobe *Clostridium acetobutylicum*. She then joined Dr. Jody Demings laboratory at the University of Washington where her research career with extremophiles from deep-sea vents started. In 1990, Anna-Louise joined Dr. Norman Pace's laboratory, where she worked on the natural phylogenetic diversity of hot spring environments in Yellowstone National Park. She then took a faculty position at Rutgers University and later at Portland State University where her research interests cover the ecology, physiology and genomics of novel and hard to grow thermophilic Archaea and Bacteria. She has served on numerous NASA, US- National Research Council, US-National Science Foundation committees and panels. Her research takes her to remote hot spring environments and she has studied thermophiles from deep-sea vents in the Indian, Pacific and Atlantic oceans.

**Mircea Podar** obtained a diploma in Biology at the University of Cluj, Romania. He then continued doctoral studies at the University of Texas Southwestern Medical Center in Dallas, Texas, where he studied the biochemistry and genetics of self-splicing group II introns in yeast mitochondria and obtained a PhD in 1997. He then pursued evolutionary studies of ctenophores (comb jellies) as well as microbial diversity and mobile genetic elements in deep sea hydrothermal vents, as a postdoctoral scholar at the Woods Hole Oceanographic Institution and the Marine Biological Laboratories in Woods Hole, Massachusetts. After a short second postdoctoral appointment at the Salk Institute in La Jolla, California, in 2000 he took a scientist position at Diversa Corporation in San Diego. There, he studied the evolution of enzyme gene families, comparative microbial genomics and metagenomics, from fundamental and biotechnological perspectives. He left Diversa in 2006 and is presently an adjunct professor at Portland State University and a scientist at Oak Ridge National Laboratory. His interests range from ctenophores biology to microbial genomics and metagenomics and their impact on understanding ecology and evolution.