

## BIOTRANSFORMATIONS

**Hiltrud Lenke**

*Chemingeering Business Design AG, Switzerland*

**Andreas Schmid**

*Technische Universität Dortmund and ISAS Leibniz Institute of Analytical Sciences, Germany*

**Keywords:** biotransformation, biodegradation, biocatalysis, environment, screening, industrial applications, enzymes, bacteria, microbes, chemical reactions, catalysis, bioprocess, synthesis

### Contents

1. Introduction
  2. Screening of Biocatalysts
    - 2.1. Classical Screening Approaches
    - 2.2. Screening by Considering Biodiversity
  3. Biodegradative Pathways for Biotransformations
    - 3.1. Aromatic Compounds
    - 3.2. Aliphatic Compounds
    - 3.3. Heterocyclic Aromatic Compounds
    - 3.4. Nitroaromatic Compounds
  4. Biocatalyst Characterization and Design
  5. Bioprocessing
    - 5.1. Biocatalyst Production
    - 5.2. Biotransformation Application
      - 5.2.1. Whole Cell Biotransformations
      - 5.2.2. Use of Isolated Enzymes
    - 5.3. Downstream Processing
  6. Technical Applications of Biotransformations
- Acknowledgements  
Glossary  
Bibliography  
Biographical Sketches

### Summary

Biotransformations are the basis of life. All natural organic and most inorganic molecules, the building blocks of organic and living matter, are subject to constant change and turnover.

The turnover time is determined by the reactivity and the reaction partners of the respective compounds, and varies from millions of years to microseconds. Fossil hydrocarbons are stored unchanged over millions of years in subsurface environments hostile for microbial enzymatic activities, whereas they are transformed to CO<sub>2</sub> and water within days or only hours in the presence of oxygen or other electron acceptors,

salts, and specialized microbes. The same is true for all other natural and most human-engineered chemicals.

These multistep transformations are catalyzed by enzymes that are mostly specific for the reaction type and starting compound. Yet, very frequently enzymes catalyze not only the reaction they were evolved for by nature over thousands or millions of years, but also the conversion of structurally and/or electronically similar derivatives.

This feature of enzymes can be used for numerous technical purposes like the clean up of contaminated environments (bioremediation) or the production of high value compounds for chemical, agricultural, and pharmaceutical industries.

In this article, we describe biotransformations used for the production of high value chemicals. We give an overview of different aspects such as the discovery and sources of enzyme based catalysts, and their design and application in bioprocesses on not only a small scale but also on an industrial scale.

There is a focus on oxygenenases but most concepts and principles described are also valid for all other enzyme classes.

## 1. Introduction

Because of the high potential of microorganisms to biodegrade natural organic compounds, they play a fundamental role in the global recycling, and thus in the maintenance of the ecological balance. Some microorganisms can utilize nearly every natural organic compound as a source of energy and/or cell building blocks.

Even many synthetic and non-natural organic compounds, so-called “xenobiotics” (see also *Biodegradation of Xenobiotics*), were shown to be biodegradable, demonstrating the enormous potential of metabolic activities in microorganisms (see also *Microorganisms as Catalysts for the Decontamination of Ecosystems and Detoxification of Chemicals*).

The variety of microorganisms able to degrade natural and synthetic organic compounds can be used for applications in environmental biotechnology as well as in industrial synthetic chemistry. In particular, the latter approach to use enzymes for biotransformations is of growing interest.

Biotransformations are chemical reactions that are catalyzed by microorganisms in terms of growing or resting cells or that are catalyzed by isolated enzymes. Because of the high stereo- or regioselectivity combined with high product purity and high enantiomeric excesses, biotransformations can be technically superior to traditional chemical synthesis.

If these features can be combined with economic benefits, biotransformations become the functional part of new chemical processes for organic synthesis. Further advantages of biocatalytical processes are the mild and ecologically harmless reaction conditions (normal pressure, low temperature, neutral pH), which are one important requirement

for sustainability. Table 1 shows the different enzyme classes and their reaction types used for biotransformations that are applied in the pharmaceutical, agrochemical, chemical, fragrance and flavor, and nutritional industries.

The use of biotransformations for industrial synthetic chemistry is an interdisciplinary, and therefore very exciting, field that needs the close cooperation of microbiologists, molecular biologists, chemists, and engineers.

As is shown in Figure 1, several steps are necessary before a biotransformation process can successfully be implemented for an industrial application. After identification of a target reaction, which may be an already existing industrial process that can be substituted by an enzymatic process, finding a suitable biocatalyst is the first crucial step in process development.

Besides classical methods, new technologies including the screening for non-culturable microorganisms (see also *Viable but Non-Culturable Bacteria in the Marine Environment and the Biotechnological Tools to Detect Them*) and high throughput screening techniques are speeding up the discovery of new biocatalysts.

As the next step, the biocatalyst has to be characterized by well-known biochemical techniques in order to identify key parameters like substrate range, reaction conditions, and kinetic data (see also *Microbial Cell Culture*).

This allows a first estimation of the reaction yield and process costs (see also *Process Optimization Strategies for Biotechnology Products: From Discovery to Production*).

Modern technologies allow an improvement of the desired biocatalyst by several engineering tools such as heterologous gene expression and protein and metabolic engineering (see also *Methods in Gene Engineering*, and *Protein Engineering*).

Enzyme class	Number		Reaction type
	classified (EC-number)	commercially available	
Oxidoreductases	650	90	Oxidation $\begin{array}{c} \text{H} \\   \\ \text{R}-\text{C}-\text{H} \\   \\ \text{H} \end{array} \xrightarrow[\text{-H}_2\text{O}]{\text{O}_2, 2[\text{H}]} \begin{array}{c} \text{H} \\   \\ \text{R}-\text{C}-\text{OH} \\   \\ \text{H} \end{array}$ Reduction $\begin{array}{c} \text{O} \\    \\ \text{R}_1-\text{C}-\text{R}_2 \end{array} \xrightarrow{2[\text{H}]} \begin{array}{c} \text{HO} \quad \text{H} \\ \diagdown \quad / \\ \text{C} \\ / \quad \diagdown \\ \text{R}_1 \quad \text{R}_2 \end{array}$ Oxygenation of C-C, C=C, (de)hydrogenation
Transferases	720	90	Transfer of complete groups: —CH <sub>3</sub> , —CH <sub>2</sub> OH, —CHO, —CH <sub>2</sub> -COOH, acyl, sugar, or phosphoryl
Hydrolases	636	150	Hydrolysis or formation of esters, amides, lactones, epoxides, nitriles, anhydrides, glycosides, and organohalides $\begin{array}{c} \text{O} \\    \\ \text{R}_1-\text{C}-\text{O}-\text{R}_2 \end{array} \rightleftharpoons \text{R}-\text{COOH} + \text{HO}-\text{R}_2$
Lyases	255	35	Addition or elimination of small molecules on C=C, C=N, C=O bonds $\begin{array}{c} \text{O} \\    \\ \text{R}_1-\text{C}-\text{H} \end{array} + \text{HCN} \rightleftharpoons \begin{array}{c} \text{OH} \\   \\ \text{R}-\text{C}-\text{CN} \\   \\ \text{H} \end{array}$

Isomerases	120	6	Isomerization such as racemization, epimerization, and rearrangement
Ligases	80	5	Bond formation or cleavage of C-O, C-S, C-N, C-C under energy consumption

Source: adapted from Faber (2000).

Table 1. Classification of enzymes valuable for biotransformations. Source: adapted from Faber, 2000.

UNESCO – EOLSS  
SAMPLE CHAPTERS

Process development has to meet economic as well as environmental constraints, and mostly includes the following steps: the production of the biocatalyst (whole cells or isolated enzymes) has to be optimized with respect to growth conditions, stability, induction, and storage (see also *Enzyme Production*). In an additional step the actual biotransformation process is optimized. Depending on the desired reaction, immobilization of the biocatalyst, enzyme catalysis in organic solvents, two liquid reactions, or cofactor regeneration are considered. Finally, a method for product recovery has to be developed using different extraction techniques, precipitation, crystallization, or *in situ* recovery. The article gives an overview of the state of the art and current developments necessary for developing a biotransformation process for industrial applications (Figure 1). Taking into account that numerous publications are available on the topic (see Bibliography) this article focuses on the screening of biocatalysts and on the use of enzymes involved in degradative pathways of xenobiotics for industrial applications. Additionally, industrial biotransformations that have successfully been applied in the synthetic chemical industry are discussed.




Figure 1. Key steps in developing a biotransformation process

## 2. Screening of Biocatalysts

Biocatalysts are based on microbes or isolated enzymes. A large number of microbes are available from national type culture collections like ATCC (USA), DSMZ (Germany), NCIB (UK), JCM (Japan) and others (see <http://www.dsmz.de/species/abbrev.htm> and see also *The Importance of Microbial Culture Collections and Gene Banks in Biotechnology*). Isolated enzymes are also commercially available from various sources (listed, for example, in Faber, 2000). Over 35 000 different reactions catalyzed by enzymes are described and can be easily

screened in databases with respect to substrate, product, productivity, and so on ([www.accelrys.com](http://www.accelrys.com) and <http://umbbd.ahc.umn.edu/>).

On the other hand, finding a powerful new biocatalyst is successful only after parallel screening of large numbers of strains in collections or after screening of mostly environmental samples—despite the impressive number of reactions described in databases. This is because of the often broad, but still limited, substrate spectrum of enzymes, and the difficulties in actually getting individual strains without restrictions for commercial use. In future, the availability of whole genome sequences and the possibility of direct cloning and heterologous expression of genes of interest will simplify this.

## 2.1. Classical Screening Approaches

The most common approach for isolating bacteria able to degrade or transform a specific organic compound is the simple use of the compound of interest as a substrate for enrichment cultures. For example, a great number of bacteria were isolated by their ability to utilize aliphatic or aromatic hydrocarbons as a growth substrate, which means as the sole source of carbon and energy. Based on the toxic behavior and on the physical or chemical properties of the respective compound, different strategies for offering the substrate are used:

- Gaseous compounds like n-butane have to be supplied via the gas phase. The substrate is continuously dissolved into the aqueous phase (the growth rate is limited by the mass transfer rate of the substrate).
- Volatile and toxic compounds are often placed in a separate glass bulb (for example, benzene or toluene). This provides the bacteria with a continuous supply of the substrate without generating a two-phase system that might kill the bacteria by destroying the functions of their membrane.
- Water insoluble compounds (for instance, naphthalene) are supplied as solids at concentrations well above their water solubility, or are offered in a second organic phase. In both cases the substrate is continuously dissolved into the aqueous phase as it is degraded, and thus the compound is rarely toxic to the bacteria.

In order to overcome problems with toxicity, nitrogen-containing compounds like nitroaromatics or nitriles are often supplied at low concentrations as the sole source of nitrogen in the presence of a readily degradable carbon and energy source. Because this enrichment strategy is often based on partial degradation of the nitrogen-containing compound, it can be particularly helpful for complex molecules.

For biotransformations, the screening by selection can be carried out with direct use of the substrate of interest or by the use of a readily degradable analogue of the substrate (cometabolism).

When screening by selection is impossible, bacteria from strain collections can be screened for the desired biotransformation, using chromometric or fluoremetric detection methods in agar or microplate experiments. Fluorogenic assays are described

for the screening of hydrolases, aldolases, and alcohol dehydrogenases. Remarkably, the latter is even an enantioselective assay. In principle, a more time-consuming screening is possible using GC, GC-MS, LC, or LC-MS methods for the analysis of the biotransformation (see also *Chemical Methods Applied to Biotechnology*, and *Physical Methods Applied to Biotechnology*). In this context it has to be mentioned that two approaches are possible. On the one hand it makes sense to screen a library of microbial catalysts using a wide range of different genera. On the other hand, if an interesting biocatalytic reaction is already described in a certain organism, it is reasonable to screen related genera and families since there are often similarities in enzymatic equipment.

In specific cases, a combination of screening by selection and screening by detection is a promising approach for biotransformations. For example, a *Rhodococcus rhodochrous* strain J1, which is used as catalyst for acrylamide production, was isolated by a combination of these methods. Acrylonitrile-utilizing bacterial strains were enriched and afterwards tested for acrylonitrile-hydrolyzing activity. In a similar manner, a collection of toluene and naphthalene degrading bacteria were screened in order to find a strain able to transform D-limonene to (+)-*trans* carveol.

A miniaturized growth system was recently developed for this purpose because even after preselection large numbers of organisms have to be screened for promising enzyme activities. This system allows maintenance, replication, and growth of microbial strains in microtiter plates without cross-contamination. After sufficient growth of the microbial strains, enzyme activities, determined by product formation, can be measured after two hours of incubation. A successful screen for oxygenases with 2000 microbial strains possessing a wide variety of catabolic activities was achieved using this method.

## 2.2. Screening by Considering Biodiversity

In the last few years, extreme environments regarding temperature, pH, pressure, and salt concentration are increasingly used as sources for new enzymes. New technologies were developed for cultivating extremophiles, but the favored approach is cloning and expressing genes from such strains into conventional host strains in order to produce enzymes with extremophilic properties.

A crucial disadvantage of the screening by selection approach described above is the fact that most bacteria in the environment are assumed to be nonculturable by using traditional isolation and cultivation techniques. To overcome this drawback a new technology in enzyme screening is becoming increasingly important. DNA extracted from environmental, and thus uncultivated, sources can be cloned and expressed into domesticated host strains. The recombinant clones are screened for new enzyme activities with high-throughput screening methods. This recombinant approach can also be used for enzymes from known microorganisms that are difficult to handle in the laboratory. An increasing number of new enzymes have been found in environmental DNA libraries. DeSantis and coworkers isolated 120 unique nitrilases using this approach. Additionally, Lorenz and coworkers were able to isolate several new lipases, esterases, and metalloproteases as well as oxygenases. Today, this fast availability of new enzymes applies pressure for the development of high-throughput biochemical enzyme characterization methods to learn about kinetics, stabilities, and morphology.

-  
-  
-

TO ACCESS ALL THE 40 PAGES OF THIS CHAPTER,  
Visit: <http://www.eolss.net/Eolss-sampleAllChapter.aspx>

## Bibliography

Adlercreutz P. (1996). Cofactor regeneration in biocatalysis in organic media. *Biocatalysis and Biotransformation* **14**, 1–30. [The review describes numerous technologies for the regeneration of cofactors / coenzymes necessary for cost efficient application of isolated cofactor dependent enzymes in organic solvents.]

Arnold F.H. (2001). Combinatorial and computational challenges for biocatalyst design. *Nature* **409**, 253–257. [The article gives an overview about enzyme engineering at a molecular level.]

Bailey J.E. and Ollis D.F. (1986). *Biochemical Engineering Fundamentals*, Columbus, Ohio: McGraw-Hill Higher Education. [The textbook gives a comprehensive introduction to biochemical engineering.]

Banwell M.G., Blakey S., Harfoot G., and Longmore R.W. (1999). *cis*-1,2-dihydrocatechol in chemical synthesis: first synthesis of L-ascorbic acid (vitamin C) from a non-carbohydrate source. *Australian Journal of Chemistry* **52**, 137–142. [The synthesis of vitamin C from chlorobenzene-*cis*-dihydrodiol is described.]

Bickerstaff G.F. (1997). *Immobilization of Enzymes and Cells*, New Jersey: Humana Press. [The textbook is one of the most comprehensive collections of immobilization methodologies.]

Blanch H.W. and Clark D.S. (1997). *Biochemical Engineering*, New York: Marcel Dekker. [The textbook gives a comprehensive overview about fundamental and applied aspects on biochemical engineering including bioreactor design and immobilized biocatalysts.]

Bosetti A., Van Beilen J.B., Preusting H., Lageveen R.G., and Witholt B. (1992). Production of primary aliphatic alcohols with a recombinant *Pseudomonas* strain, encoding the alkane hydroxylase enzyme system. *Enzyme and Microbial Technology* **14**, 702–708. [The article describes the production of 1-alkanols from the corresponding n-alkanes in two liquid bioreactors.]

Boyd D.R. and Sheldrake G.N. (1998). The dioxygenase-catalyzed formation of vicinal *cis*-diols. *Natural Product Reports* **15**, 309–324. [The article gives an overview about dioxygenase-catalyzed reactions and the classification of substrates including structure, enantiopurity, and absolute configuration of 158 vicinal *cis*-dihydrodiols.]

Brink L.E.S. and Tramper J. (1985). Optimization of organic solvent in multiphase biocatalysis. *Biotechnology and Bioengineering* **27**, 1258–1269. [Fundamental investigations of two liquid-phase biocatalytic conversions are described studying the epoxidation of propene and 1-butene by free or immobilized cells.]

Bruggink A. (2001). *Synthesis of Beta-Lactam Antibiotics: Chemistry, Biocatalysis and Process Integration*, Dordrecht, Netherlands: Kluwer. [The textbook presents industrial developments of the chemical and biochemical processes used to manufacture of beta-lactam antibiotics.]

Bruhn C., Lenke H. and Knackmuss H.J. (1987). Nitrosubstituted aromatic compounds as nitrogen source for bacteria. *Applied and Environmental Microbiology* **53**, 208–210. [Bacteria are described that were isolated by their ability to utilize nitroaromatic compounds as sole source of nitrogen.]

Bühler B., Schmid A., Hauer B., and Witholt B. (2000). Xylene monooxygenase catalyzes the multistep oxygenation of toluene and pseudocumene to corresponding alcohols, aldehydes, and acids in *Escherichia coli* JM101. *Journal of Biological Chemistry* **275**, 10085–10092. [An *E. coli* expressing the

monooxygenase genes *xylM* and *xylA* under the control of the *alk* regulatory system of *Pseudomonas oleovorans* Gpo1 is described to catalyze the multistep oxygenation of toluenes.]

Bühler B., Witholt B., Hauer B., and Schmid A. (2002). Characterization and application of xylene monooxygenase for multistep biocatalysis. *Applied and Environmental Microbiology* **68**, 560–568. [Xylene monooxygenase is used in *E. coli* for sequential hydroxylation of high concentrations of xylenes.]

Burton S.G., Cowan D.A., and Woodley J.M. (2002). The search for the ideal biocatalyst. *Nature Biotechnology* **20**, 37–45. [The review discusses recent developments of molecular technologies to search for the ideal biocatalyst for industrial applications.]

Canada K.A., Iwashita S., Shim H., and Wood T.K. (2002). Directed evolution of toluene ortho - monooxygenase for enhanced 1- naphthol synthesis and chlorinated ethene degradation. *Journal of Bacteriology* **184**, 344–349. [The article describes a molecular approach to increase the enzyme activity of a toluene-2-monooxygenase able to oxidize naphthalene to 1-naphthol.]

Carrea G., Ottolina G., and Riva S. (1995). Role of solvents in the control of enzyme selectivity in organic media. *Tibtech* **13**, 63–69. [The review discusses the effects of changing the reaction medium on enzyme enantio- and regioselectivity.]

Carrea G. and Riva S. (2000). Properties and synthetic applications of enzymes in organic solvents. *Angewandte Chemie Int.Ed.Engl*, **39**, 2226–2254. [The first part of the review discusses the factors that affect activity, stability, structure, and selectivity of enzymes in organic solvents. The second part describes a number of the synthetic applications of enzymes in organic media.]

Cheetham P.S.J. (2000). Case studies in the application of biocatalysts for the production of (bio)chemicals. *Applied Biocatalysis, 2nd edn*, (eds. A.J.J. Straathof and P. Adlercreutz), pp. 93–152. Harwood Scientific Publishers, Reading, UK. [The article lists various industrial bioprocesses for the production of (fine) chemicals.]

Chenault H.K., Simon E.S., and Whitesides G.M. (1988). Cofactor regeneration for enzyme-catalyzed synthesis. *Biotechnology and Genetic Engineering Reviews* **6**, 221–270. [A classical review describing practically feasible regeneration systems for different cofactors/coenzymes in aqueous reaction media. ]

Cherry J.R. (2000). Directed evolution of microbial oxidative enzymes. *Current Opinion in Biotechnology* **11**, 250–254. [The article describes improved oxidative enzymes obtained via directed evolution.]

Chotani G., Dodge T., Hsu A., Kumar M., LaDuca R., Trimbur D., Weyler W., and Sanford K. (2000). The commercial production of chemicals using pathway engineering. *Biochimica et Biophysica Acta* **1543**, 434–455. [The article gives an overview about pathway engineering to improve factors necessary for the successful commercial production of chemicals.]

de Bont J.A.M. (1998). Solvent-tolerant bacteria in biocatalysis. *Tibtech* **16**, 493–499. [The article presents mechanisms of bacterial solvent tolerance and highlights the use of solvent tolerant bacteria for biocatalysis.]

DeSantis G., Zhu Z., Greenberg W.A., Wong K., Chaplin J., Hanson S.R., Farwell B., Nicholson L.W., Rand C.L., Weiner D.P., Robertson D.E., and Burk M.J. (2002). An enzyme library approach to biocatalysis: development of nitrilases for enantioselective production of carboxylic acid derivatives. *Journal of American Chemical Society* **124**, 9024–9025. [A variety of nitrilases were successfully screened from environmental DNA-libraries.]

Duda M., Kuehnle A., Lenke H., Linja L., and Sieglén U. (2000) Process for the oxidation of hydrocarbons by the use of microorganisms. (EP1149918, <<http://12.espacenet.com/espacenet/viewer?PN=EP1149918&CY=ep&LG=en&DB=EPD>>. [Butane-degrading bacteria are described that can be used for the production of 1-butanol.]

Duetz W.A., Fjällman A.M., Ren S.Y., Jourdat C., and Witholt B. (2001). Biotransformation of D-limonene to (+)trans-carveol by toluene-grown *Rhodococcus opacus* PWD4 cells. *Applied and Environmental Microbiology* **67**, 2829–2832. [The article presents an example where a collection of toluene- and naphthalene-degrading bacteria was successfully screened by HPLC detection]

Duetz W.A., Ruedi L., Hermann R., O'Connor K., Buchs J., and Witholt B. (2000). Methods for intense aeration, growth, storage, and replication of bacterial strains in microtiter plates. *Applied and Environmental Microbiology* **66**, 2641–2646. [A miniaturized growth system is described that can be used for biocatalyst screening.]

Dufour E., Tam W., Nagler D.K., Storer A.C., and Menard R. (1998). Synthesis of amidrazones using an engineered papain nitrile hydratase. *FEBS Letters* **433**, 78–82. [The article describes an example where site direct mutagenesis led to an improved nitrile hydratase.]

Ellis L.B., Hershberger C.D., Bryan E.M., and Wackett L.P. (2001). The University of Minnesota Biocatalysis/Biodegradation Database: emphasizing enzymes. *Nucleic Acids Research* **29**, 340–343. [The article gives an overview about a database that provides information on microbial catabolic enzymes and their organization into metabolic pathways.]

Ensign S.A., Small F.J., Allen J.R., and Sluis M.K. (1998). New roles for CO<sub>2</sub> in the microbial metabolism of aliphatic epoxides and ketones. *Archives of Microbiology* **169**, 179–187. [The articles describes investigations on the degradative pathway of short-chain aliphatic epoxides and ketones by *Xanthobacter* strain Py2.]

Faber K. (2000). *Biotransformations in Organic Chemistry, 4th edn.* Berlin, Heidelberg, and New York: Springer-Verlag. [The textbook gives an excellent introduction to the use of biocatalysts including principles of stereoselective transformations, kinetics, and enzyme nomenclature. Different types of reactions according to the reaction principle are described.]

Flickinger M.C. and Drew S.J. (1999). *Encyclopedia of Bioprocess Technology: Fermentation Biocatalysis and Bioseparation.* New York: Wiley. [The textbook presents the applications and established theories in biotechnology focusing on industrial applications of fermentation, biocatalysis and bioseparation.]

Garcia A.A., Kim D.H., Whited G., Kwart L., Anthony W. and Downie C. (1994). Recovery of a cyclic diol produced via biocatalysis. *Isolation and Purification* **2**, 19–25. [The article outlines the use of hydrolyzed polymer adsorbents for the recovery of a *cis*-dihydrodiol from fermentation broth.]

Ghisalba O. (2000). Biocatalyzed reactions. *New Trends in Synthetic Medicinal Chemistry* (ed. F. Gualtieri), pp. 189–219. Weinheim, Germany: Wiley-VCH. [This excellent review describes the scope of biocatalysis from the point of view of a pharmaceutical company, written with great insight in limitations and perspectives. Examples are given to illustrate the arguments.]

Hagedorn S. (1985) Strain of *Pseudomonas putida* for producing an intermediate compound in the production of *para*-cresol. (US 4,542,100; <<http://patft.uspto.gov/netacgi/nph-Parser?Sect1=PTO1&Sect2=HITOFF&d=PALL&p=1&u=/netahtml/srchnum.htm&r=1&f=G&l=50&s1=4,542,100.WKU.&OS=PN/4,542,100&RS=PN/4,542,100>>). [A process is presented where 4-methylcyclohexa-3,5-dien-1,2-diol-1-carboxylic acid produced with a *Pseudomonas putida* strain is converted to *p*-cresol.]

Harrop A.J., Woodley J.M., and Lilly M.D. (1992). Production of naphthalene-*cis*-glycol by *Pseudomonas putida* in the presence of organic solvents. *Enzyme and Microbial Technology* **14**, 725–730. [The effect of water-immiscible organic solvents were investigated for the oxidation of naphthalene by *Pseudomonas putida*.]

Hashimoto S. and Ozaki A. (1999). Whole microbial cell processes for manufacturing amino acids, vitamins or ribonucleotides. *Current Opinion in Biotechnology* **10**, 604–608. [A description of processes for basic and classical fine chemicals with a focus on Japanese companies.]

He Z. and Spain J.C. (2000). Reactions involved in the lower pathway for degradation of 4- nitrotoluene by *Mycobacterium* strain HL 4-NT-1. *Applied and Environmental Microbiology* **66**, 3010–3015. [The paper describes the ring cleavage reaction in the degradation of 4-nitrotoluene that can be used for the production of picolinic acids.]

Held M., Suske W., Schmid A., Engesser K.H., Kohler H.P., Witholt B., and Wubbolts M.G. (1998). Preparative scale production of 3-substituted catechols using a novel monooxygenase from *Pseudomonas azelaica* HBP 1. *Journal of Molecular Catalysis B: Enzymatic* **5**, 87–93. [The articles gives an overview about catechols as synthons for pharmaceuticals and describes a monooxygenase useful for the production of 3-substituted catechols.]

Held M., Schmid A., Kohler H.P.E., Suske W., Witholt B., and Wubbolts M.G. (1999). An integrated process for the production of toxic catechols from toxic phenols based on a designer biocatalyst. *Biotechnology and Bioengineering* **62**, 641–648. [The production of 3-substituted catechols is described using a recombinant *E. coli* strain expressing a monooxygenase. Solid absorption materials are used for *in situ* product recovery.]

Hirai T., Fukumasa M., Nishiyama I., Yoshizawa A., Shiratori N., Yokoyama A., and Yamana M. (1991). New ferroelectric crystal having 2-methylalkanoyl group. *Ferroelectrics* **114**, 251–257. [The article gives an example that aliphatic epoxides as chiral intermediates can be used for technical applications.]

Hirose Y., Kariya K., Sasaki J., Kurono Y., Ebike H., and Achiwa K. (1992). Drastic solvent effect on lipase-catalyzed enantioselective hydrolysis of prochiral 1,4-dihydropyridines. *Tetrahedron Letters* **33**, 7157–7160. [The article gives an example where the enantioselectivity of the reaction is dependent on the solvent used.]

Hollmann F., Schmid A., and Steckhan E. (2001). First synthetic application of a monooxygenase employing indirect electrochemical NADH regeneration. *Angewandte Chemie Int.Ed.Engl.* **40**, 169–171. [A method for electrical coupling the reduction of NAD<sup>+</sup> for highly selective enzymatic hydroxylations to a cathode.]

Hudlicky T., Gonzales D., and Gibson D.T. (1999). Enzymatic dihydroxylation of aromatics in enantioselective synthesis: expanding asymmetric methodology. *Aldrichimica Acta* **32**, 35–62. [Excellent overview about the formation of vicinal *cis*-dihydrodiols and their use in asymmetric synthesis including a list of 164 *cis*-dihydrodiols produced by dioxygenase catalyzed reactions.]

Kaiser J.P., Feng Y., and Bollag J.M. (1996). Microbial metabolism of pyridine, quinoline, acridine, and their derivatives under aerobic and anaerobic conditions. *Microbiological Reviews* **60**, 483–498. [The article reviews the biodegradation of heterocyclic aromatic compounds under aerobic and anaerobic conditions with emphasis on metabolic pathways.]

Karumanchi R.S., Doddamani S.N., Sampangi C., and Todd P.W. (2002). Field-assisted extraction of cells, particles and macromolecules. *Trends in Biotechnology* **20**, 72–78. [The article describes the potential of field-assisted separations providing a major improvement in bioseparation.]

Kataoka M., Sri Rohani L.P., Wada M., Kita K., Yanase H., Urabe I., and Shimizu S. (1998). *Escherichia coli* transformant expressing the glucose dehydrogenase gene from *Bacillus megaterium* as a cofactor regenerator in a chiral alcohol production system. *Bioscience Biotechnology and Biochemistry* **62**, 167–169. [The article presents an example of the use of a recombinant *E. coli* co-expressing reductase and NADPH-regenerating glucose dehydrogenase.]

Keller K., Friedmann T., and Boxman A. (2001). The bioseparation needs for tomorrow. *Trends in Biotechnology* **19**, 438–441. [Developments of efficient, economical, and selective separation methods are discussed for successful commercialization of bioprocesses.]

Kiener A. (1992). Enzymatic oxidation of methyl groups on aromatic heterocycles: a versatile method for the preparation of heteroaromatic carboxylic acids. *Angewandte Chemie Int.Ed.Engl.* **31**, 774–775. [The article describes the biocatalytic part of an industrial process run by Lonza AG (CH) for the production of heterocyclic aromatic acids as synthons for pharmaceuticals.]

Klibanov A.M. (2001). Improving enzymes by using them in organic solvents. *Nature* **409**, 241–246. [The article summarizes the knowledge of the past 15 years about improving enzymes by using them in organic solvents.]

Kumamaru T., Suenaga H., Mitsuoka M., Watanabe T., and Furukawa K. (1998). Enhanced degradation of polychlorinated biphenyls by directed evolution of biphenyl dioxygenase. *Nature Biotechnology* **16**, 663–666. [The article demonstrated an improved activity towards various substrates after DNA shuffling of homologous genes.]

Layh N., Stolz A., Böhme J., Effenberger F., and Knackmuss H.J. (1994). Enantioselective hydrolysis of racemic naproxen nitrile and naproxen amide to *S*-naproxen by new bacterial isolates. *Journal of Biotechnology* **33**, 175–182. [Bacteria are described that were isolated by the ability to utilize nitrile and amides as sole source of nitrogen.]

Lendenmann U. and Spain J.C. (1996). 2-aminophenol 1,6-dioxygenase: a novel aromatic ring cleavage enzyme purified from *Pseudomonas pseudoalcaligenes* JS45. *Journal of Bacteriology* **178**, 6227–6232. [The enzyme characterized can be used for the production of picolinic acids.]

Leonida M.D. (2001). Redox enzymes used in chiral syntheses coupled to coenzyme regeneration. *Current Medical Chemistry* **8**, 345–369. [The review discusses the objectives of cofactor regeneration for the applicability of enzymatic redox reactions for organic synthesis.]

Li Z., Feiten H.J., Van Beilen J.B., Duetz W., and Witholt B. (1999). Preparation of optically active N-benzyl-3-hydroxypyrrolidine by enzymatic hydroxylation. *Tetrahedron-Asymmetry* **10**, 1323–1333. [The article demonstrates that alkane-degrading bacteria can be used for biohydroxylations.]

Liese A. and Filho M.V. (1999). Production of fine chemicals using biocatalysis. *Current Opinion in Biotechnology* **10**, 595–603. [The article discusses recent advances in industrial biocatalysis including the use of cofactor-dependent oxidoreductases as isolated enzymes.]

Liese A., Seelbach K., and Wandrey C. (2000). *Industrial Biotransformations*, Weinheim, Germany: Wiley-VCH. [The excellent textbook presents applied biotransformations including enzyme characteristics and process descriptions.]

Loida P.J. and Sligar S.G. (1993). Engineering cytochrome P-450cam to increase the stereospecificity and coupling of aliphatic hydroxylation. *Protein Engineering* **6**, 207–212. [The article describes an example where site direct mutagenesis led to an improved haem monooxygenase.]

Lorenz P., Liebton K., Niehaus F., Köhler B., Wolf M., Eck J., and Zinke H. (2001). *The mMetagenome: - a Challenging Source for eEnzyme dDiscovery.*, Darmstadt, Germany: BioTrans 2001, Darmstadt, Germany. [Lipases, esterases, metalloproteases, and oxygenases were successfully screened from environmental DNA-libraries.]

Lorenz P., Liebton K., Niehaus F., and Eck J., (2002). Screening for novel enzymes for biocatalytic processes: assessing the metagenome as a resource of novel functional sequence space. *Current Opinion in Biotechnology* **13**, 572-577. [The article focuses on the recognition of non-cultivated and, in particular, on prokaryotic microorganisms as dominant forms of life on earth and resource of novel valuable enzymes.]

Lye G.J. and Woodley J.M. (1999). Application of *in situ* product-removal techniques to biocatalytic processes. *Trends in Biotechnology* **17**, 395–402. [The article discusses the need for *in situ* product-removal and the process research for its implementation.]

Madigan M.T. and Mairs B.L. (1997). Extremophiles. *Scientific American* **276**, 82–87. [Extreme environments can be used as sources for new enzymes.]

Manzoni M. and Rollini N. (2002). Biosynthesis and biotechnological production of statins by filamentous fungi and application of these cholesterol-lowering drugs. *Applied Microbiology and Biotechnology* **58**, 555–564. [The biotechnological application, productivity and perspectives of fungi which are not growing in suspension are presented and discussed.]

May O., Nguyen P.T., and Arnold F.H. (2000). Inverting enantioselectivity by directed evolution of hydantoinase for improved production of L-methionine. *Nature Biotechnology* **18**, 317–320. [The hydantoinase process for production of L-methionine was improved by the use of random mutagenesis and saturation mutagenesis.]

McClay K., Fox B.G., and Steffan R.J. (2000). Toluene monooxygenase-catalyzed epoxidation of alkenes. *Applied and Environmental Microbiology* **66**, 1877–1882. [An *E. coli* strain expressing a cloned toluene-4-monooxygenase was shown to oxidize a variety of alkenes to the corresponding epoxides.]

Meulenbergh R. and de Bont J.A.M. (1995). Microbial production of catechols from nitroaromatic compounds. *Biodegradation of Nitroaromatic Compounds* (ed. J.C. Spain), pp. 37–52. New York, London: Plenum Press. [The article summarizes the research aimed at the production of catechols from nitroaromatic compounds.]

Michels P.C. and Rosazza J.P.N. (1999). Methods for biocatalysis and biotransformation. *Manual of Industrial Microbiology and Biotechnology, 2nd edn* (eds. A.L. Demain and J.E. Davies), pp. 165-180 Washington DC: ASM Press. [The article gives an excellent overview about methodologies required to develop a successful industrial biocatalytic process.]

Miura A. and Dalton H. (1995). Purification and characterization of the alkene monooxygenase from *Nocardia corallina* B-276. *Bioscience Biotechnology and Biochemistry* **59**, 853–859. [The alkene monooxygenase is capable of stereospecific epoxidation of alkenes.]

Nickerson D.P., Harford-Cross C.F., Fulcher S.R., and Wong L.L. (1997). The catalytic activity of cytochrome P450cam towards styrene oxidation is increased by site-specific mutagenesis. *FEBS Letters* **405**, 153–156. [Improving the activity of a haem monooxygenase towards an unnatural substrate showed the effectiveness of site-specific mutagenesis.]

Nikolova P. and Ward O.P. (1993). Whole cell biocatalysis in nonconventional media. *Journal of Industrial Microbiology* **12**, 76–86. [The article reviews biocatalytic reactions carried out by whole cells in nonconventional media.]

Nishino S.F. and Spain J.C. (1993) Degradation of nitrobenzene by a *Pseudomonas pseudoalcaligenes*. *Applied and Environmental Microbiology* **59**, 2520–2525. [The enzymes that can be used for the production of aminophenols are described in detail.]

Nishino S.F., Spain J.C. and He Z. (2000). Strategies for aerobic degradation of nitroaromatic compounds by bacteria: process discovery to field application. *Biodegradation of Nitroaromatic Compounds and Explosives* (eds. J.C. Spain, J.B. Hughes, and H.J. Knackmuss), pp. 7–62. Boca Raton, USA: CRC Press LLC. [Degradative pathways of nitroaromatic compounds are presented including biocatalytic applications.]

Nishino S.F., Spain J.C., Duetz W., and Witholt B. (2001). Production of phenylacetylene picolinic acid from diphenylacetylene by a toluene-degrading *Acinetobacter*. ASM Conference on Biodegradation, Biotransformation, and Biocatalysis (B3), Puerto Rico, USA, 2001. [The successful screening of toluene degrading organisms led to the isolation of a bacterial strain that can be used for the production of picolinic acids.]

Ogawa J. and Shimizu S. (1999). Microbial enzymes: new industrial applications from traditional screening methods. *Trends in Biotechnology* **17**, 13–21. [The review summarizes the industrial enzymes derived from traditional screening including the screening strategies.]

Ogawa J. and Shimizu S. (2002). Industrial microbial enzymes: their discovery by screening and use in large-scale production of useful chemicals in Japan. *Current Opinion in Biotechnology* **13**, 367–375. [The review presents large-scale production processes recently established in Japan that use microbial enzymes obtained through screening from nature.]

Olsen R.H., Kukor J.J., and Kaphammer B. (1994). A novel toluene-3-monooxygenase pathway cloned from *Pseudomonas pickettii* PKO1. *Journal of Bacteriology* **176**, 3749–3756. [The article describes the characterization of a monooxygenase that catalyzes the regioselective oxidation of toluene to *m*-cresol.]

Oppenheim S.F., Studts J.M., Fox B.G., and Dordick J.S. (2001). Aromatic hydroxylation catalyzed by toluene 4-monooxygenase in organic solvent/aqueous buffer mixtures. *Applied Biochemistry and Biotechnology* **90**, 187–197. [The catalytic function and stability of a purified toluene 4-monooxygenase was investigated in the presence of various organic solvents.]

Panke S., de L., V, Kaiser A., Witholt B., and Wubbolts M.G. (1999). Engineering of a stable whole-cell biocatalyst capable of (*S*)-styrene oxide formation for continuous two-liquid-phase applications. *Applied and Environmental Microbiology* **65**, 5619–5623. [Recombinant strains of *Pseudomonas putida* carrying genetic expression cassettes with xylene oxygenase- and styrene monooxygenase-encoding genes were shown to oxidize styrene in a continuous two-liquid-phase system.]

Panke S., Held M., Wubbolts M.G., Witholt B., and Schmid A. (2002). Pilot-scale production of (*S*)-styrene oxide from styrene by recombinant *Escherichia coli* synthesizing styrene monooxygenase. *Biotechnology and Bioengineering* **80**, 33–41. [The article presents a successful pilot-scale application of recombinant *E. coli* as the catalyst for epoxidation in an organic/aqueous reaction medium.]

Panke S. and Wubbolts M.G. (2002). Enzyme technology and bioprocess engineering. *Current Opinion in Biotechnology* **13**, 111–116. [The review discusses recent developments in enzyme technology for chemical synthesis focusing on dynamic kinetic resolution, enzyme reactions in organic solvents and two liquid-phase systems, and on the use of ionic liquids as reaction medium.]

Parales R.E., Lee K., Resnick S.M., Jiang H., Lessner D.J., and Gibson D.T. (2000). Substrate specificity of naphthalene dioxygenase: effect of specific amino acids at the active site of the enzyme. *Journal of Bacteriology* **182**, 1641–1649. [Site-directed mutagenesis was used to determine the contributions of several active-site residues of naphthalene dioxygenase.]

Patel R.N. (2001). Biocatalytic synthesis of intermediates for the synthesis of chiral drug substances. *Current Opinion in Biotechnology* **12**, 587–604. [The article provides examples of the use of enzymes for the synthesis of chiral intermediates for drugs in development.]

Pikus J.D., Studts J.M., McClay K., Steffan R.J., and Fox B.G. (1997). Changes in the regioselectivity of aromatic hydroxylation produced by active site engineering in the diiron enzyme toluene 4-monooxygenase. *Biochemistry* **36**, 9283–9289. [Site-directed mutagenesis was used to study the contributions of active site residues of toluene 4-monooxygenase.]

Rasor J.P. and Voss E. (2001). Enzyme-catalyzed processes in pharmaceutical industry. *Applied Catalysis A: General* **221**, 145–158. [The review lists and discusses selected application examples of biotransformations for pharmaceutical applications.]

Reddy J., Lee C., Neeper M., Greasham R., and Zhang J. (1999). Development of a bioconversion process for production of *cis*-1*S*,2*R*-indandiol from indene by recombinant *Escherichia coli* constructs. *Applied Microbiology and Biotechnology* **51**, 614–620. [A recombinant *E. coli* strain expressing the genes for toluene-2,3-dioxygenase and dihydrodiol dehydrogenase of *Pseudomonas putida* can be used for the production of *cis*-1*S*,2*R*-indandiol.]

Reetz M.T., Zonta A., Schimossek K., Liebton K., and Jaeger K.E. (1997). Creation of enantioselective biocatalysts for organic chemistry by *in-vitro* evolution. *Angewandte Chemie Int.Ed.Engl.* **36**, 2830–2832. [The article describes an improved *S*-enantioselectivity of a lipase that was altered using error-prone PCR.]

Resnick S.M., Lee K., and Gibson D.T. (1996). Diverse reactions catalyzed by naphthalene dioxygenase from *Pseudomonas* sp. strain NCIB 9816. *Journal of Industrial Microbiology* **17**, 438–457. [The diverse oxidation reactions catalyzed by naphthalene dioxygenase are described that can be used for the production of chiral synthons.]

Riesenberg D. and Guthke R. (1999). High-cell-density cultivation of microorganisms. *Applied Microbiology and Biotechnology* **51**, 422–430. [The article gives an overview about high-cell-density fermentations with different microorganisms.]

Salter G.J. and Kell D.B. (1995). Solvent selection for whole cell biotransformations in organic media. *Critical Reviews in Biotechnology* **15**, 139–177. [The review focuses on whole cell biotransformations in organic media with special emphasis on the mechanisms and physical bases of solvent toxicity and on the use of appropriate indicators and predictors for solvent selection.]

Schenzle A., Lenke H., Spain J.C., and Knackmuss H.J. (1999a). Chemoselective nitro group reduction and reductive dechlorination initiate degradation of 2-chloro-5-nitrophenol by *Ralstonia eutropha* JMP134. *Applied and Environmental Microbiology* **65**, 2317–2323. [The chemoselective reduction of nitro groups leading to the formation of hydroxylaminoaromatic compounds is described.]

Schenzle A., Lenke H., Spain J.C., and Knackmuss H.J. (1999b). 3-Hydroxylaminophenol mutase from *Ralstonia eutropha* JMP134 catalyzes a Bamberger rearrangement. *Journal of Bacteriology* **181**, 1444–1450. [The rearrangement of hydroxylaminoaromatic compounds by a mutase is an important step in the bioproduction of aminophenols.]

Schiraldi C. and De Rosa M. (2002). The production of biocatalysts and biomolecules from extremophiles. *Trends in Biotechnology* **20**, 515–521. [The article describes recent developments in the production of biomass and related enzymes and biomolecules from extremophile sources.]

Schleper C., Swanson R.V., Mathur E.J., and DeLong E.F. (1997). Characterization of a DNA polymerase from the uncultivated psychrophilic archaeon *Cenarchaeum symbiosum*. *Journal of Bacteriology* **179**, 7803–7811. [The article describes a successful example where a new enzyme activity was screened from non-culturable microorganisms.]

Schmid A., Kollmer A., Mathys R.G., and Witholt B. (1998). Developments toward large-scale bacterial bioprocesses in the presence of bulk amounts of organic solvents. *Extremophiles* **2**, 249–256. [The article

discusses fundamental questions using two-liquid processes and describes the development of a high-pressure, explosion proof bioreactor system.]

Schmid A., Dordick J.S., Hauer B., Kiener A., Wubbolts M., and Witholt B. (2001). Industrial biocatalysis today and tomorrow. *Nature* **409**, 258–268. [The article gives an overview about current use and future trends of biocatalysis in chemical industry discussed by the use of process examples of BASF, DSM, and Lonza.]

Schulze B. and Wubbolts M.G. (1999). Biocatalysis for industrial production of fine chemicals. *Current Opinion in Biotechnology* **10**, 609–615. [Main industrial applications of biocatalysts and recent developments to perform resolutions of racemic mixtures, dynamic kinetic resolutions, or asymmetric syntheses are outlined.]

Shields M.S., Montgomery S.O., Chapman P.J., Cuskey S.M., and Pritchard P.H. (1989). Novel pathway of toluene catabolism in the trichloroethylene-degrading bacterium G4. *Applied and Environmental Microbiology* **55**, 1624–1629. [The article describes the characterization of a monooxygenase that catalyzes the regioselective oxidation of toluene to *o*-cresol.]

Shiratori N., Yoshizawa A., Nishiyama I., Fukumasa M., Yokoyama A., and Hirai T. (1991). New ferrolytic liquid crystals having 2-fluoro-2-methyl alkanoyloxy group. *Molecular Crystals and Liquid Crystals* **199**, 129–140. [The article gives an example that aliphatic epoxides as chiral intermediates can be used for technical applications.]

Short J.M. (1997). Recombinant approaches for accessing biodiversity. *Nature Biotechnology* **15**, 1322–1323. [The article gives an overview about the opportunity to consider biodiversity by analyzing environmental DNA libraries.]

Small F.J. and Ensign S.A. (1997). Alkene monooxygenase from *Xanthobacter* strain Py2. Purification and characterization of a four-component system central to the bacterial metabolism of aliphatic alkenes. *Journal of Biological Chemistry* **272**, 24913–24920. [The article focuses on the characterization of an alkene monooxygenase from *Xanthobacter* strain Py2 that catalyzes the epoxidation of short chain alkenes.]

Smith M.R. (1990). The biodegradation of aromatic hydrocarbons by bacteria. *Biodegradation* **1**, 191–206. [The articles present degradative pathways of aromatic compounds by bacteria with special emphasis on benzene, toluene, naphthalene, biphenyl, and selected fused aromatic hydrocarbons.]

Spieß T., Desiere F., Fischer P., Spain J.C., Knackmuss H.J., and Lenke H. (1998). A new 4-nitrotoluene degradation pathway in a *Mycobacterium* strain. *Applied and Environmental Microbiology* **64**, 446–452. [Enzymes that can be used for the bioproduction of aminophenols are described.]

Straathof A.J.J. and Adlercreutz P. (2000). *Applied Biocatalysis, 2nd edn.* Netherlands: Harwood Academic Publishers. [The textbook presents fundamentals of biocatalysis with technological experience and commercial case studies. This includes information on reactor engineering and immobilization of biocatalysts.]

Takahashi O., Umezawa J., Furuhashi K., and Takagi M. (1989). Stereocontrol of a tertiary hydroxyl group via microbial epoxidation. A facile synthesis of prostaglandine  $\gamma$ -chains. *Tetrahedron Letters* **30**, 1583–1584. [The article described a bioprocess to synthesize useful precursors of tertiary alcohols.]

Thomas S.M., DiCosimo R., and Nagarajan V. (2002). Biocatalysis: applications and potentials for the chemical industry. *Trends in Biotechnology* **20**, 238–242. [A recent review discussing the application of biotransformations in the traditional chemical industry, including a critical view on challenges for biocatalyst and bioprocess development.]

Trudgill P.W. (1990a). Cyclopentanone 1,2-monooxygenase from *Pseudomonas* NCIMB 9872. *Methods Enzymology* **188**, 77–81. [The article focuses on the characterization of cyclopentanone 1,2-monooxygenase that catalyzes a Baeyer-Villiger oxidation.]

Trudgill P.W. (1990b). Cyclohexanone 1,2-monooxygenase from *Acinetobacter* NCIMB 9871. *Methods Enzymology* **188**, 70–77. [The article focuses on the characterization of cyclohexanone 1,2-monooxygenase that catalyzes a Baeyer-Villiger oxidation.]

van de Sandt E.J.A.X. and De Vroom E. (2000). Innovations in cephalosporin and penicillin production: painting the antibiotics industry green. *Chimica Oggi-Chemistry Today* **18**, 72–75. [Biocatalytic routes to produce these antibiotics are described with respect to sustainability.]

Vicenzi J.T., Zmijewski M.J., Reinhard M.R., Landen B.E., Muth W.L., and Marler P.G. (1997). Large-scale stereoselective enzymatic ketone reduction with *in-situ* product removal via polymeric resins. *Enzyme and Microbial Technology* **20**, 494–499. [The article describes a process where polymeric hydrophobic resins were used to supply the toxic substrate and to remove the toxic product from the reaction mixture.]

Wackett L.P. and Hershberger C.D. (2001). *Biocatalysis and Biodegradation: Microbial Transformation of Organic Compounds*, Washington D.C.: ASM Press. [The textbook presents fundamental concepts of the microbial transformation of organic compounds and its application for biotechnology and biodegradation.]

Wahler D. and Reymond J.L. (2001). Novel methods for biocatalyst screening. *Current Opinion in Chemical Biology* **5**, 152–158. [The article gives an overview on recent developments in high-throughput enzyme assays for the screening of biocatalysts.]

Watkinson R.J. and Morgan P. (1990). Physiology of aliphatic hydrocarbon-degrading microorganisms. *Biodegradation* **1**, 79–92. [The article reviews aspects of the physiology and biochemistry of the microbial degradation of aliphatic hydrocarbons.]

Weijers C.A., de Haan A., and de Bont J.A. (1988). Microbial production and metabolism of epoxides. *Microbiological Science* **5**, 156–159. [The article describes several bacteria that form optically active epoxides or are able to degrade epoxides enantioselectively allowing the bioproduction of chiral building blocks in organic synthesis.]

Wendeborn S., De Mesmaeker A.D., and Brill W.K.D. (1998). Polymer bound 3,5-cyclohexadiene-1,2-diols as core structures for the development of small molecule libraries. *Synlett* **1998**, 865–868. [The article outlines the highly efficient sequence for the synthesis of complex chiral and highly functionalized molecules on solid phase bound cyclohexadienediols.]

Werlen C., Kohler H.P., and van der Meer, Jr. (1996). The broad substrate chlorobenzene dioxygenase and *cis*-chlorobenzene dihydrodiol dehydrogenase of *Pseudomonas* sp. strain P51 are linked evolutionarily to the enzymes for benzene and toluene degradation. *Journal of Biological Chemistry* **271**, 4009–4016. [The article describes four dioxygenase families that were identified by amino acid sequence comparison of the catalytic oxygenase subunit.]

Wery J., Mendes da Silva D.I., and de Bont J.A. (2000). A genetically modified solvent-tolerant bacterium for optimized production of a toxic fine chemical. *Applied Microbiology and Biotechnology* **54**, 180–185. [The production of 3-methylcatechol is described in a two liquid-phase system using a solvent-tolerant *Pseudomonas putida* strain as biocatalyst.]

Whited G.M. and Gibson D.T. (1991). Toluene-4-monooxygenase, a three-component enzyme system that catalyzes the oxidation of toluene to *p*-cresol in *Pseudomonas mendocina* KR1. *Journal of Bacteriology* **173**, 3010–3016. [The article describes the characterization of a monooxygenase that catalyzes the regioselective oxidation of toluene to *p*-cresol.]

Wieser M., Heinzmann K., and Kiener A. (1997). Bioconversion of 2-cyanopyrazine to 5-hydroxypyrazine-2-carboxylic acid with *Agrobacterium* sp. DSM 6336. *Applied Microbiology and Biotechnology* **48**, 174–176. [A fermentation process for the two-enzyme-step bioconversion is described.]

Witholt B., De Smet M.J., Kingma J., Van Beilen J.B., Kok M., Lageveen R.G., and Eggink G. (1990). Bioconversions of aliphatic compounds by *Pseudomonas oleovorans* in multiphase bioreactors: background and economic potential. *Tibtech* **8**, 46–52. [The article outlines the potential of bioconversions by the alkane degrading *Pseudomonas oleovorans* in two-liquid-phase bioreactors for the production of intermediate value compounds]

Wubbolts M.G., Hoven J., Melgert B., and Witholt B. (1994). Efficient production of optically active styrene epoxides in two-liquid phase cultures. *Enzyme and Microbial Technology* **16**, 887–894. [The article outlines the potential of biooxidations in two-liquid-phase systems for the production of styrene epoxides.]

Yoshikawa N., Ohta K., Mizuno S., and Ohkishi H. (1993). Production of *cis,cis*-muconic acid from benzoic acid. *Bioprocess Technology* **16**, 131–147. [A technical process for the production of *cis,cis*-muconic acid is described using whole cells of an *Arthrobacter* strain as biocatalyst.]

Zaks A. (2001). Industrial biocatalysis. *Current Opinion in Chemical Biology* **5**, 130–136. [Industrial process examples are described to highlight beneficial aspects of biocatalysis.]

Zelder O. and Hauer B. (2000). Environmentally directed mutations and their impact on industrial biotransformation and fermentation processes. *Current Opinion in Microbiology* **3**, 248–251. [The article presents traditional and industrial biocatalyst improvement.]

Zhang J. and Greasham R. (1999). Chemically defined media for commercial fermentations. *Applied Microbiology and Biotechnology* **51**, 407–421. [The review focuses on the application development and practical considerations to use chemically defined media for industrial bioprocesses.]

### Biographical Sketches

**Hiltrud Lenke**, born in 1961, studied chemistry at the University of Wuppertal, Germany. She finished her studies in 1986 with a diploma thesis that dealt with the biodegradation of 2,6-dinitrophenol. She took her Ph.D. at the Institute for Microbiology at the University of Stuttgart, Germany, in the subject of the microbial degradation of polynitrophenols. She continued this work for another year as a post-doctoral student. Between 1991 and 2002 she worked as a group leader at the Fraunhofer Institute for Interfacial Engineering and Biotechnology in Stuttgart, Germany. Initially, her main task was in the area of environmental microbiology. Since 1998, her main emphasis has been in biotransformation and biocatalysis. During the summer of 2001 she finished her habilitation to gain the *venia legendi* for microbiology. Since January 2002, she has been a visiting lecturer for microbiology at the University of Stuttgart. In 2003, she started work as a consultant for biotechnology at Chemengineering GmbH in Wiesbaden, Germany.

**Andreas Schmid**, born in 1966, studied Biology with a focus on immunology and microbiology at the University of Stuttgart where he received his Diploma in 1992. During his studies in Stuttgart and at the Swiss Federal Institute of Environmental Science and Technology (EAWAG) in Dübendorf he worked as an academic assistant at the Fraunhofer Society (Stuttgart) and at the Institute of Microbiology (University of Stuttgart) until 1992. In February 1997 he received a PhD in microbiology from the University of Stuttgart. After research activities in applied environmental biotechnology (waste air treatment), at the faculty of civil engineering, University of Stuttgart he joined the Institute of Biotechnology (ETH Zurich) as post doctoral fellow concentrating on the development of microbial biocatalysts for organic synthesis reactions. Since January 1999 he is lecturer and head of the research group 'Technical Enzymology' at the Institute of Biotechnology, ETH Zurich. His research group is specialized on the development and application of bacterial enzymes as biocatalysts with respect to basic biochemistry, protein engineering and process development.